

PII: S0040-4039(96)01795-9

Syntheses of Highly Constrained β -Aryl Isohexanoic Acid Derivatives Via Asymmetric Michael Addition

Subo Liao, Yinglin Han,[†] Wei Qiu, Michael Bruck and Victor J. Hruby *

Department of Chemistry, The University of Arizona, Tucson, Arizona 85721

Abstracts: A series of enantiomerically pure highly sterically hindered β -branched isohexanoic acid derivatives have been synthesized with high diastereoselectivity via asymmetric Michael addition. The X-ray crystal structure of (4S,3'S)-3-[3'-(2,6-dimethylphenyl)isohexanyl]-4-phenyl-2-oxazolidinone demonstrated that the β -configuration was induced from the Si-face, and that the torsional angle χ_2 was restricted by the bulky β -isopropyl group to the range expected from molecular modeling. Copyright © 1996 Elsevier Science Ltd

The stereoselective Michael addition of organometallic reagents to α,β -unsaturated acyl derivatives which contain a chiral auxiliary is one of the most reliable methods to synthesize optically pure β -branched carbonyl derivatives.¹⁻⁴ Although various products have been prepared with high diastereoselectivity via asymmetric Michael additions to chiral oxazoline,² ester,³ amide,⁴ or imide derivatives,⁵ to our knowledge there still have not been any examples of the syntheses of highly sterically hindered β -branched carboxylic acids derivatives by this method. As part of our efforts to design and synthesize conformationally constrained novel β -branched amino acids,⁶ we have developed an efficient procedure to prepare these important intermediates (*i.e.* enantiomerically pure β -methyl carboxylic acid derivatives) via asymmetric Michael addition by using optically pure 4-phenyloxazolidinone as an auxilliary.⁷ Using this methodology, the asymmetric Michael addition of



various non-hindered Grignard reagents in the presence of $\text{CuBr}\cdot\text{SMe}_2$ and the co-solvent dimethylsulfide could be carried out with high stereoselectivity and in reasonable yields.⁷ However, in our initial attempts to

[†] Permanent address: Nanjing University of Chemical Technology, Nanjing 210009, P. R. China

synthesize much more sterically-hindered β -isopropyl aromatic amino acids, the Michael addition of isopropyl magnesium chloride to the Michael acceptor, (4R,2E)-3-[3'-(4-methoxyphenyl)propenoyl]-4-phenyl-2-oxazolidinone, did not work well using the previously developed reaction conditions. Instead, we applied a chiral resolution method to synthesize the desired optically pure intermediates of β -phenylisohexanoic acid derivatives.⁸

Recently, we have re-examined the asymmetric Michael reaction in order to develop a general approach for the asymmetric syntheses of hindered unusual β -isopropyl aromatic amino acids. A new Michael reaction acceptor, (4S,2E)-3-(1-oxo-2-isohexenyl)-4-phenyl-2-oxazolidinone, was tested, and it was found that it could react with a variety of aromatic Grignard reagents, including the highly sterically hindered 2',6'-dimethylphenyl magnesium bromide under mild conditions.



Table 1. Diastereoselective Syntheses of β -Aryl Isohexanoic Acid Derivatives

Entry	R ₁	R ₂	R ₃	Yield ^a	de⁵	[α] _D ^{20°} ^c	$\Delta \upsilon (Hz)^d$
1	Н	Н	Н	82%	>90%	+9.7°	57.2
2	OCH ₃	н	н	97%	>90%	+6.0°	56.8
3	ОСН3	СН,	н	94%	>90%	+7.3°	69.9
4	н	CH ₃	CH ₃	68%	>90%	+7.6°	98.9
5	OCH ₃	CH ₃	CH ₃	86%	>90%	+6.6°	98.6

a. The yield was calculated from the weight of product after column chromatography purification; b. The de was determined from 'H NMR of crude product,¹¹ de>90% means that no diasteroisomer was observed from 'H NMR; c [c] = 1 mg/mL in CDCl₃. d. Δv is the chemical shift difference between the two methyl groups of the β -isopropyl group.

As compared to the previously reported reaction conditions,⁷ we have found the reactions of the phenyl or substituted phenyl Grignard reagents only require catalytic amounts of $CuBr \cdot SMe_2$ and no dimethylsulfide as co-solvent. However, the para-methoxylated phenyl Grignard reagents needed one equivalent of $CuBr \cdot SMe_2$ to secure the complete conjugation addition reaction. Otherwise, 1,2-addition reactions could be observed, and even became the major reaction pathway.⁹ The reason why phenyl and para-methoxyphenyl Grignard reagents require different amounts of Cu(I) catalyst for 1,4-conjugation addition is still not clear, and is under investigation. In any case, all products were obtained in good to excellent yields with high optical purity (Table 1). These β -branched isohexanoic acid derivatives could be easily converted into their corresponding unusual β -isopropyl aromatic amino acids using chemistry previously elaborated in our laboratory.⁷

An X-ray crystal structure of the compound (4S,3'S)-3-[3'-(2,6-dimethylphenyl)isohexanyl]-4-phenyl-2oxazolidinone 4 has been determined and the results are shown in Figure 1. These results suggested that in the asymmetric synthesis, the β -carbon configuration was induced from the Si-face presumably because the Re-face was shielded by the phenyl ring of the oxazolidinone in the Michael reaction transition state,⁹ and the rotation of aromatic ring is found to be χ_2 =+112.6° which is in the angle range expected from computer modeling.¹⁰ The chemical shift differences ($\Delta \upsilon$) of the two isopropyl methyl groups could be used to assess the degree of constraint in these β -aryl isohexanoic acid derivatives. As seen from the Table 1, $\Delta \upsilon$ increases with the number of methyl substituents on the aromatic ring. Compounds 4 and 5 have the largest $\Delta \upsilon$ values, compounds 1 and 2 the smallest $\Delta \upsilon$ value. The para-methoxy substituent on the aromatic ring has no effects on $\Delta \upsilon$ values. Thus, the presence of the 2' and 6'-dimethyl groups and of the β -isopropyl group produce torsional constraints which reduce greatly the flexibility of β -aryl isohexanoic acid derivatives, which are important precursors for preparation of β -isopropyl aromatic amino acids.



Figure 1. The Stereo view of the crystal structure of (4S, 3'S)-3-[3'-(2,6-dimethyl-phenyl)isohexanyl]-4-phenyl-2-oxazolidinone 4

Experimental Procedures: To a mixture of RMgBr (1.7 mmol, 1.5 equivalents) and CuBr-SMe₂ (36 mg, 0.15 equivalents except for para-methoxyphenyl magnesium bromide which needs 1.0 equivalent) in 5 mL THF at -20°C was added dropwise a solution of (4S,2E)-3-(1-oxo-2-isohexenyl)-4-phenyl-2-oxazolidinone (1.16 mmol, 1.0 equivalent) in 3 mL of freshly distilled THF. A yellow color was observed during the addition process. The resulting mixture was kept stirring at -15°C for two hours, then slowly warmed to room temperature during one hour. The reaction was then quenched with 20 mL of saturated ammonium chloride, and the product was extracted with ether (3 x 20 mL). The combined organic extracts were washed with brine (2 x 20 mL), water (20 mL), and dried over anhydrous magnesium sulfate. After evaporation of dried organic phases, the crude product was purified by silica gel chromatography.

Acknowledgments: The authors wish to acknowledge Prof. R. S. Glass for very helpful discussions about this research, and the financial support of grants from NIDA DA 06284 and DA 08657, and U.S. Public Health Service DK 21085.

References and Notes

- a) Tomioka, K. and Koga, K. in "Asymmetric Synthesis", Ed., Morrison, J.D. Academic Press, 1983, Vol. 2, 201-224. b) Barner, B.A. and Meyers, A.I. J. Am. Chem. Soc. 1984, 106, 1865-1866. c) Oppolzer, W.; Poli, G.; Kingma, A.J.; Starkemann, C. and Bernardinelli, G. Helv. Chim. Acta. 1987, 70, 2201-2214.
- 2. a) Meyers, A.I. and Whitten, C.E. J. Am. Chem. Soc. 1975, 97, 6266-6267. b) Meyers, A.I. and Shipman, M. J. Org. Chem. 1991, 56, 7098-7102.
- a) Fang, C.; Ogawa, T.; Suemune, H. and Sakai, K. *Tetrahedron: Asymmetry* 1991, 2, 389-398.
 b) Kawana, M. and Emoto, S. *Bull. Chem. Soc. Jpn.* 1966, 39, 910-916.
- 4. a) Mukaiyama, T. and Iwasawa, N. Chem. Lett. 1981, 913-916. b) Touet, J.; Baudouin, S. and Bown, E. Tetrahedron: Asymmetry 1992, 3, 587-590.
- 5. a) Melnyk, O.; Stephan, E.; Pourcelot, C. and Cresson, P. Tetrahedron 1992, 48, 841-850. b) Tomioka, K.; Suenaga, T. and Koga, K. Tetrahedron lett. 1986, 27, 369-372.
- (a) Hruby, V. J. Biopolymers 1993, 1073-1082; (b) Hruby, V. J.; Al-Obeidi, F. and Kazmierski, W. Biochem. J. 1990, 268, 249-262.
- (a) Dharanipragada, R.; VanHulle, K.; Bannister, A.; Bear, S.; Kennedy, L. and Hruby, V. J. *Tetrahedron* 1992, 48, 4733-4748; (b) Nicolas, E.; Russell, K.C.; Hruby, V.J. J. Org. Chem. 1993, 58, 7766-7770; (c) Nicolas, E.; Russell, K.C.; Knollenberg, J. and Hruby, V.J. J. Org. Chem. 1993, 58, 7565 -7571; (d) Jiao, D.; Russell, K.C. and Hruby, V.J. Tetrahedron 1993, 49, 3511- 3520; (e) Li, G.; Patel, D.; and Hruby, V.J. J. Chem. Soc. Perkin Trans. 1 1994, 3057-3059; (f) Qian, X.; Russell, K. C.; Boteju, L.W. and Hruby, V. J. Tetrahedron 1995, 51, 1033-1054.
- 8. Liao, S. and Hruby, V. J. Tetrahedron Lett.. 1996, 37, 1563-1566.
- 9. Low, B-S.; Li, G.; Lung, F-D. and Hruby, V.J. J. Org. Chem. 1995, 60, 5509-5514.
- 10. Shenderovich, M.; Liao, S. and Hruby, V.J. unpublished results.
- ¹H NMR: δ ppm 2. 7.20-6.74(m, 9H, aromatic protons), 5.31(dd, J=8.7, 4.Hz, 1H, oxazolidinone, -11. CHAr-), 4.58(t, J=8.7Hz, 1H, oxazolidinone, -CH₂-/proR), 4.08(dd, J=7.1, 4.8Hz, 1H, oxazolidinone, $-CH_2$ -/proS), 3.80(s, 3H, -ArOCH₃), 3.74(dd, J=15.5, 5.2Hz, 1H, -C_aH-/proS), 3.07(dd, J=15.5, 5.1Hz, 1H, -C_aH-/proR), 2.87(m, 1H, -C_bH-), 1.83[m, 1H, -CH(CH₃)₂], 0.95(d, J=6.7Hz, 3H, -CH₃), 0.72(d, J=6.7Hz, 3H, -CH₃); 3. 7.20-6.52(m, 8H, aromatic protons), 5.28(dd, J=8.7, 4.1Hz, 1H, oxazolidinone, -CHAr-), 4.53(t, J=8.8Hz, 1H, oxazolidinone, -CH₂-/proR), 4.04(dd, J=8.7, 4.1Hz, 1H, oxazolidinone, -CH,/proS), 3.80(dd, J= 15.0, 10.0Hz, 1H, -C,H-/proS), 3.79(s, 3H, -ArOCH₂), 3.13(m, 1H, -C₆H-), 2.97(dd, J=15.0, 4.7Hz, 1H, -C₆H/proR), 2.07(s, 3H, -ArCH₃), 1.81[m, 1H, -CH(CH₃)₂], 1.00(d, J=6.6Hz, 3H, -CH₃), 0.72(d, J=6.7Hz, 3H, -CH₃); 4. 7.25-6.83(m, 8H, aromatic protons), 5.31(dd, J=8.74, 4.0Hz, 1H, oxazolidinone -CHAr-), 4.60(t, J=8.7Hz, 1H, oxazolidinone, -CH₂-/proR), 4.13(dd, J=8.7, 4.1Hz, 1H, oxazolidinone, -CH₂-/proS), 3.64-3.35(m, 3H, -C_aH₂C_BH-), 2.45(s, 3H, -CH₃), 2.25-2.18[m, 1H, -CH(CH₃)₂], 2.17(s, 3H, -CH₃), 1.06(d, J=6.4Hz, 3H, -CH₃), 0.66(d, J=6.7Hz, 3H, -CH₃); 5. 7.26-6.39(m, 7H, aromatic protons), 5.31(dd, J=8.7, 4.0Hz, 1H, oxazolidinone, -CHAr-), 4.59(t, J=8.7Hz, 1H, oxazolidinone, -CH₂/proR), 4.13(dd, J=8.8, 4.0Hz, H, oxazolidinone, -CH₂-/proS), 3.74(s, 3H, -ArOCH₃), 3.65(dd, J=15.0, 7.8Hz, 1H, -C_aH-/proS), 3.50- $3.27(m, 2H, -C_{g}H-/proR and -C_{g}H-)$, $2.43(s, 3H, -ArCH_{3})$, $2.14[m, 1H, -CH(CH_{3})_{2}]$, 2.13(s, 3H, -ArCH3), 1.05(d, J=6.5Hz, 3H, -CH₃), 0.66(d, J=6.7Hz, 3H, -CH₃).

(Received in USA 9 July 1996; revised 3 September 1996; accepted 9 September 1996)