

Organic Preparations and Procedures International: The New Journal for Organic Synthesis

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/uopp20>

SYNTHESIS OF 6H-bis-[1]-BENZOPYRANO[2,3-b:3',4'-e]PYRIDIN-8(8H)ONES AND 3-(2'-HYDROXYBENZOYL)-5H-[1]BENZOPYRANO[4,3-b]PYRIDINES AND THEIR DERIVATIVES

K. Venkat Reddy^a, G. Sabitha^a & A. V. Subba Rao^a

^a Department of Chemistry, P. G. College of Science Saifabad, Osmania University, Hyderabad, 500 004, INDIA

Published online: 18 Feb 2009.

To cite this article: K. Venkat Reddy, G. Sabitha & A. V. Subba Rao (1996): SYNTHESIS OF 6H-bis-[1]-BENZOPYRANO[2,3-b:3',4'-e]PYRIDIN-8(8H)ONES AND 3-(2'-HYDROXYBENZOYL)-5H-[1]BENZOPYRANO[4,3-b]PYRIDINES AND THEIR DERIVATIVES, Organic Preparations and Procedures International: The New Journal for Organic Synthesis, 28:3, 325-332

To link to this article: <http://dx.doi.org/10.1080/00304949609356538>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.tandfonline.com/page/terms-and-conditions>

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae, and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand, or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.


**SYNTHESIS OF 6H-bis-[1]-BENZOPYRANO[2,3-b:3',4'-e]PYRIDIN-8(8H)ONES
AND 3-(2'-HYDROXYBENZOYL)-5H-[1]BENZOPYRANO[4,3-b]PYRIDINES
AND THEIR DERIVATIVES**

K. Venkat Reddy, G. Sabitha[†] and A. V. Subba Rao*

*Department of Chemistry, P. G. College of Science
Saifabad, Osmania University, Hyderabad -500 004, INDIA*

4-Chromanone (**1**)¹ is an attractive synthon for the elaboration of a variety of heterocycles fused at 3,4 of benzopyran moiety.² Such a feature is encountered in several physiologically important heterocyclic systems.³ We now describe, the synthesis of 6H-bis-[1]-benzopyrano[2,3-b:3',4'-e]pyridin-8(8H)-ones and 3-(2'-hydroxybenzoyl)-5H-[1]-benzopyrano[4,3-b]pyridines, making use of 4-chromanone (**1**) by reaction with 3-cyanochromone (**2**),⁴ and 3-formylchromone (**4**).⁵ These compounds (**2** and **4**) possess active reaction sites in the pyrone ring suitable for the construction of fused heterocyclic molecules by nucleophilic addition-cyclization reaction.⁶⁻⁸

Reaction of **1** with 3-cyanochromone (**2**) in refluxing ethanol in the presence of a catalytic amount of piperidine proceeds as shown in Scheme 1. Michael addition of **1** on the pyrone ring, with concomitant opening of the pyrone ring, followed by double cyclization and loss of a molecule of



water results in the formation of 6-H-bis-[1]-benzopyrano[2,3-*b*:3',4'-*e*]pyridin-8(8*H*)-ones (**3a-h**) (Table 1). The ¹H NMR (**3a,b,d**)⁹ and mass spectra of all the compounds are consistent with the assigned structures. Reaction of **1** with 3-formylchromone (**4**) gives 3-(4-oxo-4*H*-[1]-benzopyrano-3-methynyl)-2,3-dihydro-4*H*-1-benzopyran-4-ones (**5a-j**) (Table 2). These compounds react with ammonium acetate in acetic acid or with ammonia in aqueous methanolic medium to give 3-(2'-hydroxybenzoyl)-5*H*-[1]-benzopyrano[4,3-*b*]pyridines (**6a-j**) (Table 2); the former method however gives unsatisfactory yields due to the poor solubility of the reactants (**5a-j**) in acetic acid. The *o*-hydroxybenzoyl moiety of this system may be exploited further in building up 3-phenylcoumarins and 2-benzoylbenzofuran ring systems. Thus, reaction of **6** with phenylacetyl chloride in the presence of aqueous K₂CO₃ in a two -phase system using tetrabutylammonium hydrogen sulfate (TBAHSO₄) as the phase transfer catalyst and dichloromethane (DCM) as organic solvent gives 3-(2-oxo-3-phenyl-2*H*-[1]-benzopyran-4-yl)-5*H*-[1]-benzopyrano[4,3-*b*]pyridines **7a-j** (Table 3).

Similar reaction of **6** with *p*-chlorophenacyl bromide gives 3-(2-benzoylbenzo[*b*]furan-3-yl)-5*H*-[1]-benzopyrano[4,3-*b*]pyridines, **8a-e** (Table 2) in good yields. The spectral data of all these compounds supports the assigned structures.


TABLE 1. Yields, Mps and Elemental Analyses of compounds **3a-h**

Cmpd.	Yield (%)	mp (°C)	Elemental Analyses Calcd (Found)		
			C	H	N
3a	66	218-223	75.74 (76.04)	3.68 (3.91)	4.65 (4.84)
3b	75	246-248	76.18 (76.43)	4.16 (4.34)	4.44 (4.66)
3c	60	295-297	67.97 (67.96)	3.00 (3.15)	4.17 (4.01)
3d	68	282-286	60.02 (59.72)	2.65 (2.45)	3.68 (3.58)
3e	70	230-236	67.97 (68.26)	3.00 (3.19)	4.17 (4.33)
3f	83	250-257	68.68 (68.40)	3.46 (3.23)	4.01 (3.90)
3g	65	285-290	61.64 (61.34)	2.45 (2.23)	3.78 (3.56)
3h	78	290-294	55.04 (55.35)	2.19 (2.37)	3.38 (3.60)


EXPERIMENTAL SECTION

Melting points are uncorrected. ¹H NMR spectra were determined on Varian Gemini 200 MHz spectrometer (Internal Me₄Si). Mass spectra were recorded on a VG Micromass 70-70H instrument. Thin layer chromatography (TLC) was performed on Merck silica gel 60F254 precoated plates (0.2mm).

6*H*-bis-[1]-benzopyrano[2,3-*b*:3',4'-*e*]pyridin-8(8*H*)-ones (3a-h). General Procedure.- A mixture of 4-chromanone (**1a-b**) (30 mmol) and 3-cyanochromone (**2a-d**) (25 mmol) was refluxed in ethanol (50 mL) in the presence of piperidine (1 mL) for 4 hrs. The reaction mass was cooled to room temperature and the resulting solid was collected and crystallized from ethanol to give **3a-h** in 60-83% yield (Table 1).



- a) $R_1 = R_2 = H$, b) $R_1 = CH_3$, $R_2 = H$
 c) $R_1 = Cl$, $R_2 = H$, d) $R_1 = Br$, $R_2 = H$
 e) $R_1 = Cl$, $R_2 = CH_3$



- a) $R = R_1 = H$, b) $R = CH_3$, $R_1 = H$
 c) $R = Cl$, $R_1 = H$, d) $R = Br$, $R_1 = H$,
 e) $R = Cl$, $R_1 = CH_3$

5, 6, 7	a	b	c	d	e	f	g	h	i	j
R	H	H	H	H	H	Cl	Cl	Cl	Cl	Cl
R ₁	H	CH ₃	Cl	Br	Cl	H	CH ₃	Cl	Br	Cl
R ₂	H	H	H	H	CH ₃	H	H	H	H	CH ₃

- i) EtOH, Et₃N (67-80%) ii) MeOH, aq. NH₄OH (50-63%) iii) PhCH₂COCl/TBAHSO₄, DCM, aq. K₂CO₃ (48-70%) iv) 4-Chlorophenacyl bromide, TBAHSO₄, DCM, aq. K₂CO₃ (60-84%)

TABLE 2. Yields, mps and Elemental Analyses of Compounds **5a-j**, **6a-j**, **7a-j** and **8a-e**

Cmpd	Yield ^a (%)	mp (°C)	Elemental Analyses Calcd (Found)		
			C	H	N
5a	75	180-185	74.99 (75.23)	3.98 (4.17)	
5b	80	170-174	75.46 (75.21)	4.43 (4.52)	
5c	70	198-200	67.37 (67.64)	3.28 (3.13)	
5d	68	203-205	59.55 (59.41)	2.89 (2.77)	
5e	78	205-208	68.09 (68.04)	3.71 (3.77)	
5f	77	219-221	67.37 (67.50)	3.27 (3.48)	
5g	79	225-227	68.09 (68.37)	3.71 (3.86)	
5h	67	250-253	61.15 (61.30)	2.70 (2.70)	
5i	69	258-260	54.64 (54.54)	2.41 (2.32)	
5j	72	250-252	62.03 (62.29)	3.12 (3.30)	
6a	63	108-110	75.24 (75.07)	4.32 (4.20)	4.62 (4.50)
6b	60	118-122	75.69 (75.54)	4.76 (4.64)	4.41 (4.23)
6c	51	124-126	67.56 (67.82)	3.58 (3.41)	4.15 (4.31)
6d	54	148-150	59.70 (59.60)	3.17 (3.03)	3.66 (3.26)
6e	62	151-153	68.28 (68.53)	4.01 (4.09)	3.98 (3.76)
6f	61	140-145	67.56 (67.31)	3.58 (3.39)	4.15 (4.05)
6g	59	146-149	68.28 (68.55)	4.01 (4.20)	3.98 (3.80)
6h	52	188-191	61.31 (61.05)	2.98 (2.89)	3.76 (3.66)
6i	50	180-183	54.77 (54.88)	2.66 (2.78)	3.36 (3.56)
6j	55	174-176	62.19 (62.27)	3.39 (3.48)	3.63 (3.65)
7a	66	198-200	80.38 (80.63)	4.25 (4.55)	3.47 (3.71)
7b	68	232-235	80.56 (80.89)	4.59 (4.89)	3.56 (3.56)
7c	60	238-240	74.06 (74.35)	3.68 (3.38)	3.20 (3.10)
7d	58	238-242	67.23 (67.62)	3.34 (3.66)	2.90 (2.74)
7e	70	240-245	74.42 (74.04)	4.01 (4.19)	3.10 (3.39)
7f	62	253-266	74.06 (74.41)	3.68 (3.99)	3.20 (3.45)
7g	61	188-190	74.42 (74.04)	4.01 (3.71)	3.10 (3.31)
7h	50	223-226	68.66 (68.51)	3.20 (3.04)	2.97 (2.84)
7i	48	239-242	62.75 (62.86)	2.93 (3.12)	2.71 (2.91)
7j	58	253-255	69.15 (69.40)	3.52 (3.74)	2.88 (3.08)
8a	60	143-145	74.06 (74.43)	3.68 (3.92)	3.20 (3.38)
8b	65	178-180	74.42 (74.16)	4.01 (4.30)	3.10 (3.25)
8c	69	166-168	68.66 (68.90)	3.20 (3.00)	2.97 (2.70)
8d	72	186-190	62.75 (62.38)	2.93 (2.61)	2.71 (2.47)
8e	80	198-202	69.15 (69.01)	3.52 (3.20)	2.88 (2.90)

a) After recrystallization

TABLE 3. Spectroscopic Data of Compounds **5a-j** and **6a-j**

Cmpd	¹ H NMR (CDCl ₃ , DMSO- <i>d</i> ₆), δ, J (Hz)
5a	5.15 (s, 2H), 6.80-7.09 (m, 2H), 7.38-7.55 (m, 4H) 7.65-7.75 (m, 1H), 7.92 (s, 1H), 7.96 (dd, 1H, J = 2.0, 8.5), 8.21 (d, 1H, J = 8.5).
5b	2.42 (s, 3H), 5.13 (s, 2H), 6.88-7.06 (m, 2H), 7.34-7.60 (m, 4H), 7.89 (s, 1H), 7.96 (dd, 1H, J = 2.0, 8.5), 8.02 (d, 1H, J = 8.5).
5c	5.18 (s, 2H), 6.94-7.12 (m, 2H), 7.44-7.70 (m, 4H), 7.92 (s, 1H), 8.02 (d, 1H, J = 8.5), 8.26 (d, 1H, J = 8.5).
5d	5.18 (s, 2H), 6.96-7.14 (m, 2H), 7.40-7.60 (m, 3H), 7.82 (dd, 1H, J = 2.0, 8.5), 7.96 (s, 1H), 8.04 (dd, 1H, J = 2.5, 8.5), 8.40 (d, 1H, J = 8.5).
5e	2.41 (s, 3H), 5.16 (s, 2H), 6.87-7.10 (m, 2H), 7.32-7.59 (m, 4H), 7.88 (s, 1H), 7.95 (dd, 1H, J = 2.0, 8.5), 8.04 (d, 1H, J = 8.5).
5f	5.15 (s, 2H), 6.98 (d, 1H, J = 8.5), 7.09 (d, H, J = 8.5), 7.43-7.70 (m, 4H), 7.93 (s, 1H), 8.04 (dd, 1H, J = 2.0, 8.5), 8.23 (d, 1H, J = 8.5).
5g	2.49 (s, 3H), 5.20 (s, 2H), 6.92 (d, 1H, J = 8.5), 7.38-7.49 (m, 2H), 7.53-7.68 (m, 2H), 7.98 (dd, 2H, J = 2.0, 8.5), 8.06 (s, 1H).
5h	5.15 (s, 2H), 6.89-7.10 (m, 2H), 7.43-7.69 (m, 3H), 7.96 (s, 1H), 8.03 (d, 1H, J = 2.5), 8.28 (d, 1H, J = 8.5).
5i	5.18 (s, 2H), 6.96-7.12 (m, 2H), 7.44-7.70 (m, 2H), 7.80 (dd, 1H, J = 2.0, 8.5), 7.98 (s, 1H), 8.06 (dd, 1H, J = 2.0, 8.5), 8.39 (d, 1H, J = 8.5).
5j	2.48 (s, 3H), 5.15 (s, 2H), 7.02-7.60 (m, 4H), 7.92 (s, 1H), 8.04 (d, 1H, J = 8.5), 8.32 (d, 1H, J = 2.5).
6a	5.25 (s, 2H), 6.88-7.64 (m, 7H), 7.80 (s, 1H), 8.28 (dd, 1H, J = 2.0, 8.5), 8.84 (d, 1H, J = 2.0), 11.82 (s, 1H).
6b	2.30 (s, 3H), 5.30 (s, 2H), 6.88 (s, 1H), 7.05 (s, 1H), 7.10-7.55 (m, 4H), 7.78 (s, 1H), 8.30 (d, 1H, J = 10.0), 8.86 (d, 1H, J = 2.0), 11.62 (s, 1H).
6c	5.23 (s, 2H), 6.92-7.59 (m, 6H), 7.76 (s, 1H), 8.35 (dd, 1H, J = 2.0, S.5), 8.82 (d, 1H, J = 2.0) 11.67 (s, 1H).
6d	5.28 (s, 2H), 6.72-7.78 (m, 7H), 8.25 (dd, 1H, J = 2.5, 8.5), 8.82 (d, 1H, J = 2.0), 11.64 (s, 1H).
6e	2.42 (s, 3H), 5.35 (s, 2H), 7.0 (s, 1H), 7.06-7.48 (m, 3H), 7.60 (s, 1H), 7.78 (s, 1H), 8.30 (d, 1H, J = 10.0), 8.82 (d, 1H, J = 2.0), 11.75 (s, 1H).
6f	5.23 (s, 2H), 6.86-7.60 (m, 6H), 7.78 (s, 1H), 8.20 (d, 1H, J = 2.0), 8.82 (d, 1H, J = 2.0), 1.72 (s, 1H).
6g	2.30 (s, 3H), 5.30 (s, 2H), 6.90-7.02 (m, 2H), 7.25-7.40 (m, 3H), 7.80 (s, 1H), 8.25 (d, 1H, J = 2.0), 8.82 (d, 1H, J = 2.0), 11.60 (s, 1H).
6h	5.28 (s, 2H), 6.98 (d, 1H, J = 8.5), 7.10 (d, 1H, J = 8.5), 7.36-7.60 (m, 3H), 7.80 (s, 1H), 8.30 (d, 1H, J = 2.0), 8.88 (d, 1H, J = 2.0), 11.70 (s, 1H).
6i	5.35 (s, 2H), 6.98 (d, 1H, J = 8.5), 7.02 (d, 1H, J = 8.5), 7.36-7.75 (m, 3H), 7.80 (s, 1H), 8.30 (d, 1H, J = 2.0), 8.90 (d, 1H, J = 2.0), 11.70 (s, 1H).
6j	2.44 (s, 3H), 5.16 (s, 2H), 6.96-7.02 (m, 2H), 7.28 (dd, 1H, J = 2.0, 8.5), 7.48 (s, 1H), 7.80 (s, 1H), 8.25 (d, 1H, J = 2.0), 8.95 (d, 1H, J = 2.0), 11.65 (s, 1H).

TABLE 4. Spectroscopic Data of Compounds **7a-j** and **8a-e**

Cmpd	¹ H NMR (CDCl ₃ , DMSO-d ₆), δ, J (Hz)
7a	5.1 (d, 2H, J = 10.0), 6.95 (d, 1H, J = 10.0), 7.05-7.65 (m, 12H), 8.18 (dd, 1H, J = 2.5, 10.0), 8.4 (d, 1H, J = 2.5).
7b	2.33 (s, 3H), 5.1 (d, 2H, J = 10.0), 6.9-7.38 (m, 12H), 8.18 (dd, 1H, J = 2.5, 10.0), 8.39 (d, 1H, J = 2.5).
7c	5.1 (d, 2H, J = 10.0), 6.96 (d, 1H, J = 10.0), 7.08-7.48 (m, 10H), 7.53 (dd, 1H, J = 2.5, 10.0), 8.18 (dd, 1H, J = 2.5, 10.0), 8.35 (d, 1H, J = 2.5).
7d	5.15 (d, 2H, J = 10.0), 6.94 (d, 1H, J = 10.0), 7.05-7.40 (m, 10H), 7.68 (dd, 1H, J = 2.5, 10.0), 8.2 (dd, 1H, J = 2.5, 10.0), 8.4 (d, 1H, J = 2.5).
7e	2.50 (s, 3H), 5.12 (d, 2H, J = 10.0), 6.91 (d, 1H, J = 10.0), 7.13-7.41 (m, 10H), 8.17 (dd, 1H, J = 2.5, 10.0), 8.42 (d, 1H, J = 2.5).
7f	5.10 (d, 2H, J = 10.0), 6.69 (d, 1H, J = 10.0), 7.1-7.32 (m, 8H), 7.45-7.65 (m, 3H), 8.14 (d, 1H, J = 4.5), 8.39 (d, 1H, J = 2.5).
7g	2.31 (s, 3H), 5.1 (d, 2H, J = 10.0), 6.89-7.40 (m, 11H), 8.16 (d, 1H, J = 4.5), 8.40 (d, 1H, J = 2.5).
7h	5.15 (d, 2H, J = 10.0), 6.93 (d, 1H, J = 10.0), 7.1-7.62 (m, 10H), 8.19 (d, 1H, J = 4.5), 8.40 (d, 1H, J = 2.5).
7i	5.14 (d, 2H, J = 10.0), 6.91 (d, 1H, J = 10.0), 7.10-7.40 (m, 9H), 7.70 (dd, 1H, J = 2.5, 10.0), 8.19 (d, 1H, J = 4.5), 8.40 (d, 1H, J = 2.5).
7j	2.5 (s, 3H), 5.12 (d, 2H, J = 10.0), 6.90 (d, 1H, J = 10.0), 7.10-7.38 (m, 9H), 8.16 (d, 1H, J = 4.5), 8.40 (d, 1H, J = 2.5).
8a	5.21 (s, 2H), 6.95 (d, 1H, J = 10.0), 7.06-7.70 (m, 9H), 7.90 (d, 1H, J = 10.0), 7.98 (d, 1H, J = 8.5), 8.23 (dd, 1H, J = 2.0, 8.5), 8.78 (d, 1H, J = 2.5).
8b	2.49 (s, 3H), 5.29 (s, 2H), 6.98 (d, 1H, J = 10.0), 7.10-7.56 (m, 7H), 7.80 (s, 1H), 7.98 (s, 1H), 8.02 (s, 1H), 8.3 (dd, 1H, J = 2.0, 8.5), 8.84 (d, 1H, J = 2.5).
8c	5.22 (s, 2H), 6.95 (d, 1H, J = 10.0), 7.05-7.58 (m, 6H), 7.65 (d, 1H, J = 10.0), 7.73 (d, 1H, J = 10.0), 7.89 (d, 1H, J = 8.5), 7.94 (d, 1H, J = 8.5), 8.24 (dd, 1H, J = 2.0, 8.5), 8.78 (d, 1H, J = 2.5).
8d	8.23 (s, 2H), 6.96 (d, 1H, J = 10.0), 7.08-7.60 (m, 6H), 7.69 (d, 1H, J = 10.0), 7.78 (d, 1H, J = 10.0), 7.91 (d, 1H, J = 8.5), 7.98 (d, 1H, J = 8.5), 8.25 (dd, 1H, J = 2.0, 8.5), 8.80 (d, 1H, J = 2.5).
8e	2.58 (s, 3H), 5.29 (s, 2H), 6.99 (d, 1H, J = 10.0), 7.1-7.74 (m, 7H), 7.95 (s, 1H), 8.05 (s, 1H), 8.26 (dd, 1H, J = 2.0, 8.5), 8.78 (d, 1H, J = 2.5).

3-(4-Oxo-4H-[1]-benzopyrano-3-methynyl)-2,3-dihydro-4H-[1]-benzopyran-4-ones (5a-j).

General Procedure.- A mixture of 3-formylchromone (**4a-e**) (25 mmol), 4-chromanone (**1a-b**) (30 mmol) and triethylamine (30 mmol) in ethanol (50 mL) was refluxed for 3 hrs. and cooled. The resulting pale yellow crystalline solid was filtered, washed with cold ethanol (20 mL) and recrystallized from ethanol to give **5a-j** in 67-80% yield (Table 2).

3-(2'-Hydroxybenzoyl)-5*H*-[1]-benzopyrano[4,3-*b*]pyridines (6*a-j*). **General Procedure.**- A mixture of **5a-j** (5g) methanol (155 mL) and aqueous ammonia (25% w/w, 25 mL) was refluxed for 2 hrs. Methanol was distilled off, and the reaction mass was diluted with cold water. It was then acidified with conc. HCl to give a yellow solid. The solid was collected, washed with cold water and crystallized from alcohol to give **6a-j** in 50-63% yield (Table 2).

3-(2-Oxo-3-phenyl-2*H*-[1]-benzopyran-4-yl)-5*H*-[1]-benzopyrano[4,3-*b*]pyridines (7*a-j*). **General Procedure.**- To a stirred mixture of the compound **6a-j** (7 mmol) in dichloromethane (30 mL) and aqueous K₂CO₃ (30% w/v, 30 mL), tetrabutylammonium hydrogen sulfate (50 mg) was added. A solution of phenylacetyl chloride (15 mL) was added dropwise to the above mixture under stirring during 15 min. The reaction mass was stirred at room temperature for 6-8 hrs. The organic layer was separated, washed with water (2 X 20 mL), dried (Na₂SO₄) and the solvent distilled off. The residue was crystallized from ethanol to give **7a-j** in 48-70% yield (Table 2).

3-(2-Benzoylbenzo[*b*]furan-3-yl)-5*H*-[1]-benzopyrano[4,3-*b*]pyridines (8*a-e*). **General Procedure.**- To a stirred mixture of **6a-e** (7 mmol) in dichloromethane (30 mL) and aqueous K₂CO₃ (30% w/v, 30 mL), tetrabutylammonium hydrogen sulfate (50 mg) was added, followed by dropwise addition of 4-chlorophenacyl bromide (8 mmol) in dichloromethane (10 mL). The reaction mixture was stirred at room temperature for 8-10 hrs. Usual work up gave **8a-e** in 60-84% yield (Table 2).

REFERENCES

- † Present address: Indian Institute of Chemical Technology, Hyderabad-500 007, India.
1. H. J. Kabbe and A. Widdig, *Angew. Chem. Int. Ed. Engl.* **21**, 247 (1982).
 2. C. K. Ghosh, *J. Indian Chem. Soc.*, **67**, 5 (1990).
 3. a) G. Doria, A. M. Isetta, M. Ferrari and D. Trizio, *Eur. Pat. Appl. EP* 433, 274 (1988); S. P. A. Farmitalia Carlo Erba; *Chem. Abstr.*, **111**, 57711 (1989); b) K. M. Sangwan and S. Rastogi, *Indian J. Chem.*, **20B**, 135 (1981); c) J. Freedman, *US Patent*, 3,707,476 (1972); Colgate-Palmolive Co., *Chem. Abstr.*, **78**, 72120 (1973); *US Patent*, 3,825,539 (1974); Colgate-Palmolive Co., *Chem. Abstr.*, **82**, 43396 (1975); *US Patent*, 3,553,238 (1971); Colgate-Palmolive Co., *Chem. Abstr.*, **75**, 5881 (1971); *US Patent*, 3,553,230 (1971); *Chem. Abstr.*, **75**, 5882 (1971); *US Patent*, 3,553,229 (1971); Colgate-Palmolive Co., *Chem. Abstr.*, **75**, 5883 (1971); d) C. G. Wermuth, P. Worms, J. J. Bourguignon and R. Brodin, *Eur. Pat. Appl. EP* 323, 325 (1989); SANOFI, *Chem. Abstr.*, **111**, 232848 (1989); e) R. Bogner, C. Jaszberenyi, S. Maakleit, A. Fodor, P. Deak and T. Timmer, *PCT Int. Appl. WO* 8800 197; Alkaloida vegyeszeti Gyar, *Chem. Abstr.*, **109**, 73451 (1988); f) W. Werner, W. Jungstand, W. Gutsche, K. Wohlrabe, W. Roemer and D. Tresselt, *Pharmazie*, **34**, 394 (1979); *Chem. Abstr.*, **92**, 41906 (1980); g) A. H. Phillipp and I. L. Jirkovsky, *US Patent*, 4,177,284 (1979); h) Ayerst, McKenna and Harrison Ltd., *Chem. Abstr.*, **92**, 163953 (1980); *Can. J. Chem.*, **57**, 3292 (1979); i) M. Sacauet, M. Bellassoued-Fargeau, B. Graffe and P. Maitte, *J. Heterocyclic Chem.*, **22**, 757 (1985).
 4. A. Nohara, T. Umetani and Y. Sanno, *Tetrahedron Lett.*, 1995 (1973); *Tetrahedron*, **30**, 3553 (1974).

5. A. Nohara, *Tetrahedron Lett.*, 1187 (1974).
6. G. J. Reddy and A. V. S. Rao, *Curr. Sci.*, **50**, 84 (1981).
7. G. J. Reddy, G. Sabitha and A. V. S. Rao, *Synth. Commun.*, **17**, 1851 (1987).
8. C. K. Ghosh, J. *Heterocyclic Chem.*, **20**, 1437 (1983).
9. ¹H NMR of **3c**, **3e**, **3f**, **3g** and **3h** could not be recorded because of their insolubility.

(Received August 24, 1995; in revised form December 21, 1995)