

C11—N1—C12	114.4 (3)	C1—C10—C5	118.6 (4)
C2—C1—C10	119.9 (3)	C1—C10—C9	124.5 (4)
C2—C1—C11	118.9 (3)	C5—C10—C9	116.9 (4)
C10—C1—C11	121.2 (3)	P1—C11—N1	112.8 (2)
O1—C2—C1	118.2 (3)	P1—C11—C1	112.5 (3)
O1—C2—C3	120.7 (4)	N1—C11—C1	112.3 (3)
C1—C2—C3	121.1 (4)	N1—C12—C13	110.7 (3)
C2—C3—C4	119.7 (4)	O2—C13—C12	107.8 (4)
C3—C4—C5	121.8 (4)	O5—C14—C15	111.1 (4)

Table 3. *Hydrogen-bonding geometry* (Å, °)

D—H...A	D—H	H...A	D...A	D—H...A
N1—H12...O3 ⁱ	0.810	1.998	2.761 (4)	157.06
O1—H1...O4 ⁱⁱ	0.819	1.716	2.529 (3)	171.63
O2—H2...O3 ⁱⁱⁱ	0.830	1.996	2.789 (5)	160.83

Symmetry codes: (i) $2-x, -y, 2-z$; (ii) $x-1, y, z$; (iii) $1-x, -y, 2-z$.

All non-H atoms were refined with anisotropic displacement parameters. The H atoms of the naphthyl group were placed geometrically 0.95 Å from their corresponding C atoms, while other H atoms were taken from a $\Delta\rho$ map. For all H atoms a riding model was used with $U_{\text{iso}}(\text{H}) = 1.3U_{\text{eq}}(\text{C}, \text{N}, \text{O})$.

Data collection: *CAD-4 Express* (Enraf–Nonius, 1993). Data reduction: *MolEN* (Fair, 1990). Program(s) used to solve structure: *SIMPEL* in *MolEN*. Program(s) used to refine structure: *LSFM* in *MolEN*. Molecular graphics: *ORTEPII* (Johnson, 1976) in *MolEN*. Software used to prepare material for publication: *MolEN*.

The authors wish to acknowledge the purchase of the CAD-4 diffractometer under Grant DPT/TBAG1 of the Scientific and Technical Research Council of Turkey, and the Research Foundation of Ankara University, Ankara, Turkey, for their support in the preparatory part of this study (Project No. 95-25-00-06).

Lists of structure factors, anisotropic displacement parameters, H-atom coordinates and complete geometry have been deposited with the IUCr (Reference: AB1384). Copies may be obtained through The Managing Editor, International Union of Crystallography, 5 Abbey Square, Chester CH1 2HU, England.

References

- Aguilar, M., Miralles, N. & Sastre, M. A. (1989). *Rev. Inorg. Chem.* **10**, 93–119.
- Assche, I. van, Soroka, M., Haemers, A., Hooper, M., Blanot, D. & van Heijenoort, J. (1991). *Eur. J. Med. Chem.* **26**, 505–515.
- Cameron, D. G., Hudson, H. R. & Pianka, M. (1993). *Phosphorus Sulfur Silicon*, **83**, 21–37.
- Choi, N., Failla, S., Finocchiaro, P., McPartlin, M. & Scowen, I. J. (1994). *J. Chem. Res. (S)*, pp. 340–341.
- Enraf–Nonius (1993). *CAD-4 Express Software*. Version 1.1. Enraf–Nonius, Delft, The Netherlands.
- Fair, C. K. (1990). *MolEN. An Interactive Intelligent System for Crystal Structure Analysis*. Enraf–Nonius, Delft, The Netherlands.
- Gibson, D. & Karaman, R. (1989). *Inorg. Chem.* **28**, 1928–1932.
- Johnson, C. K. (1976). *ORTEPII*. Report ORNL-5138. Oak Ridge National Laboratory, Tennessee, USA.
- Lavielle, G., Hautefaye, P., Schaeffer, C., Boutin, J. A., Cudennec, C. A. & Pierré, A. (1991). *J. Med. Chem.* **34**, 1998–2003.
- Toroš, Z. R., Prodić, B. K. & Šljukić, M. (1978). *Acta Cryst.* **B34**, 3110–3113.

Acta Cryst. (1996). **C52**, 3135–3138

1-Diphenylphosphino-2-phenyl-1,2-dicarba-closo-dodecaborane(12)

M. ANNA McWHANNELL,^a GEORGINA M. ROSAIR,^{a*} ALAN J. WELCH,^a FRANCESC TEIXIDOR^b AND CLARA VIÑAS^b

^aDepartment of Chemistry, Heriot-Watt University, Riccarton, Edinburgh EH14 4AS, Scotland, and ^bDepartament de Química, Universitat Autònoma de Barcelona–CSIC, Campus de Bellaterra, 08193 Bellaterra, Barcelona, Spain. E-mail: chegmr@bonaly.hw.ac.uk


(Received 10 May 1996; accepted 4 July 1996)

Abstract

The two crystallographically independent molecules of the title compound, 1-[(C₆H₅)₂P]-2-C₆H₅-1,2-C₂B₁₀H₁₀ (C₂₀H₂₅B₁₀P), in the asymmetric fraction of the unit cell are highly congruent. The conformation of the cage phenyl groups is defined by $\theta = 1.15^\circ$ and the average C1—C2 distance is 1.744 (8) Å, which is significantly longer than that usually found for 1,2-C₂B₁₀H₁₀ compounds. The PPh₂ substituent is tilted away from an idealized radial position on C1 as a result of intramolecular crowding.

Comment

1-Diphenylphosphino-2-phenyl-1,2-dicarba-closo-dodecaborane(12), (I), is an exceptionally stable compound with the potential for extensive derivatization of both the phosphine moiety and the carborane polyhedron. It was synthesized and characterized in order to extend the series of structural determinations of carboryl phosphines (Kivekäs, Sillanpää, Teixidor, Viñas & Nuñez, 1994; Kivekäs, Teixidor, Viñas & Nuñez, 1995) and also to establish a structural basis for its subsequent use as a carborane-containing ligand (Reid, 1992).



There are two crystallographically independent molecules in the asymmetric fraction of the unit cell. A perspective view of one molecule (atom labels unprimed) is shown in Fig. 1 and Table 2 lists selected geometric parameters. There is a high degree of congruence between the two independent molecules, with

no equivalent parameters differing by greater than 3σ . Consequently, average values are quoted hereafter.




Fig. 1. The molecular structure of (I) shown with 40% probability displacement ellipsoids for non-H atoms. The cage numbering scheme can be seen in this and Fig. 2. The phenyl ring C atoms are numbered in sequence.

The C_2B_{10} cage has the expected near icosahedral geometry. The conformation of the phenyl substituent at C2 is described by a θ value of 1.15° (θ is defined as the modulus of the average $C_{Ph}-C_{Ph}-C_{cage}-C_{cage}$ torsion angle; Cowie, Reid, Watmough & Welch, 1994) and thus the phenyl ring is face-on to the phosphine substituent in order to minimize interactions arising from the bulky PPh_2 substituent at C1. We have shown previously (Brain *et al.*, 1996) that the C1—C2 distance in icosahedral carboranes is sensitive to θ , with low θ values contributing to weak C1—C2 bonding (Lewis & Welch, 1993). In the present compound, C1—C2 is 1.744 (8) Å, which is 0.042 (10) Å longer than the corresponding value in 1- PPh_2 -2-Me-1,2- $C_2B_{10}H_{10}$ (Kivekäs *et al.*, 1994) and 0.078 (12) Å longer than in 1- PPh_2 -1,2- $C_2B_{10}H_{11}$ (Kivekäs *et al.*, 1995). Normally in undistorted 1,2-dicarboranes $C-C < C-B < B-B$ (Thomas, Rosair & Welch, 1996). In the title compound, the average C1—C2 bond is extended beyond the limit of C—B distances [1.692 (7)–1.739 (7) Å], while remaining below the range of B—B distances [1.753 (8)–1.801 (9) Å].

The P—C1 distance [1.876 (7) Å] compares well with that found in 1- PPh_2 -1,2- $C_2B_{10}H_{11}$ [1.871 (6) Å; Kivekäs *et al.*, 1995] and that in 1- PPh_2 -2-Me-1,2- $C_2B_{10}H_{10}$ [1.888 (4) Å; Kivekäs *et al.*, 1994]. Moreover, the similarity between these three compounds extends to a pronounced deformation of the PPh_2 substituent away from an ideal radial position on C1, resulting in a wide range of P1—C1— C_{cage} angles, from 111.6 (6) (P1—

C1—C2) to $130.0 (5)^\circ$ (P1—C1—B4). Previously, this deformation has been ascribed to intramolecular steric congestion between cage and PPh_2 fragments (Kivekäs *et al.*, 1994). The title molecule as viewed along the C1...B12 vector is shown in Fig. 2 and it is clear that the H5 and H3 atoms lie under the phosphine-bound phenyl rings. Ring plane-to-H atom distances are 2.72 and 2.79 Å, and are at the limit of the sum of the van der Waals radii for hydrogen and an aromatic ring, even using the lowest accepted value (1.0 Å) of the former.




Fig. 2. A view down the C1...B12 vector showing the distortion of the phosphine away from the radial position at C1.

Inspection of Fig. 2 also suggests that the lone pair of electrons on the P atom lies above the B6—C2 connectivity, displaced towards the latter. The stereochemical consequences of utilizing this lone pair of electrons in coordination to metal fragments will be the subject of future contributions.

Experimental

For the preparation of (I), 1-Ph-1,2-*closo*- $C_2B_{10}H_{11}$ (Brain *et al.*, 1996) (1.00 g, 4.54 mmol) was dissolved in diethyl ether (20 ml) under a dry nitrogen atmosphere and cooled to 273 K. MeLi (5 mmol, 3.6 ml of a 1.4 M solution in diethyl ether) was added dropwise and the resulting solution allowed to warm to room temperature with stirring over a period of 30 min. PPh_2Cl (1.10 g, 0.89 ml, 5 mmol) was added dropwise to the stirred mixture, resulting in the formation of a white precipitate. The mixture was stirred for a further 30 min, then filtered through silica (2 cm). Volatiles were removed *in vacuo*, affording a white adhesive solid. This was triturated with *n*-pentane and dried *in vacuo*, resulting in a dry white powder (1.54 g, 84%). Crystals were grown by slow

evaporation of an *n*-pentane solution at room temperature. Analysis calculated for C₂₀H₂₅B₁₀P: C 59.4, H 6.23%; found: C 59.4, H 6.62%. FTIR (CH₂Cl₂): $\nu = 2605\text{ cm}^{-1}$ (B—H). ¹H FTMNR (400.1 MHz, CDCl₃, 293 K, TMS): δ (p.p.m.) 7.85–7.15 (*m*, Ar—H). ¹¹B—{¹H} FTMNR (128.4 MHz, CDCl₃, 293 K, BF₃·Et₂O): δ (p.p.m.) 3.04 (1B), 0.36 (1B), –5.81 (8B). ³¹P—{¹H} FTMNR (162.0 MHz, CDCl₃, 293 K, H₃PO₄): δ (p.p.m.) 12.66 (*s*). MS (FAB, Noba) parent ion: 407. NMR spectra were recorded on a Bruker DPX400 spectrometer, the IR spectrum on a Nicolet Impact 400 spectrophotometer and the mass spectrum on a VG/MS9 spectrometer.

Crystal data

C ₂₀ H ₂₅ B ₁₀ P	Mo <i>K</i> α radiation
<i>M_r</i> = 404.47	$\lambda = 0.71073\text{ Å}$
Monoclinic	Cell parameters from 32 reflections
<i>P</i> ₂ ₁ / <i>c</i>	$\theta = 5.56\text{--}23.13^\circ$
<i>a</i> = 23.896 (2) Å	$\mu = 0.124\text{ mm}^{-1}$
<i>b</i> = 10.608 (4) Å	<i>T</i> = 293 (2) K
<i>c</i> = 20.173 (3) Å	Block
$\beta = 114.218 (6)^\circ$	0.30 × 0.25 × 0.20 mm
<i>V</i> = 4663.7 (17) Å ³	Colourless
<i>Z</i> = 8	
<i>D_x</i> = 1.152 Mg m ^{−3}	
<i>D_m</i> not measured	

Data collection

Siemens <i>P4</i> diffractometer	<i>R</i> _{int} = 0.059
ω scans	$\theta_{\text{max}} = 25^\circ$
Absorption correction:	<i>h</i> = −28 → 26
ψ scans (<i>SHELXTL</i> ;	<i>k</i> = −12 → 1
Sheldrick, 1994)	<i>l</i> = −1 → 23
<i>T</i> _{min} = 0.823, <i>T</i> _{max} =	3 standard reflections
0.872	monitored every 97
9893 measured reflections	reflections
8127 independent reflections	intensity decay: 6%
3636 observed reflections	
[<i>I</i> > 2σ(<i>I</i>)]	

Refinement

Refinement on <i>F</i> ²	$w = 1/[\sigma^2(F_o^2) + (0.0585P)^2]$
$R[F^2 > 2\sigma(F^2)] = 0.0689$	where $P = (F_o^2 + 2F_c^2)/3$
$wR(F^2) = 0.1810$	(Δ/σ) _{max} < 0.001
<i>S</i> = 0.988	$\Delta\rho_{\text{max}} = 0.278\text{ e Å}^{-3}$
8122 reflections	$\Delta\rho_{\text{min}} = -0.233\text{ e Å}^{-3}$
559 parameters	Extinction correction: none
H atoms riding with C—H	Atomic scattering factors
0.93, B—H 1.10 Å and	from <i>International Tables</i>
<i>U</i> (H) = 1.2 <i>U</i> _{eq} (C/B)	for <i>Crystallography</i> (1992,
	Vol. C, Tables 4.2.6.8 and
	6.1.1.4)

Table 1. Fractional atomic coordinates and equivalent isotropic displacement parameters (Å²)

	<i>x</i>	<i>y</i>	<i>z</i>	<i>U</i> _{eq}
P1	0.82304 (6)	0.65246 (14)	0.22525 (7)	0.0436 (3)
C1	0.8837 (2)	0.7729 (5)	0.2748 (3)	0.0404 (11)
C2	0.9351 (2)	0.7150 (5)	0.3594 (3)	0.0420 (12)
B3	0.8835 (2)	0.8331 (6)	0.3545 (3)	0.0438 (13)

$$U_{\text{eq}} = (1/3)\sum_i\sum_j U_{ij}a_i^*a_j^*\mathbf{a}_i\cdot\mathbf{a}_j.$$

B4	0.8778 (3)	0.9326 (6)	0.2811 (4)	0.052 (2)
B5	0.9233 (3)	0.8674 (6)	0.2395 (3)	0.053 (2)
B6	0.9561 (3)	0.7263 (7)	0.2869 (3)	0.0467 (14)
B7	0.9618 (3)	0.8352 (7)	0.4201 (3)	0.0531 (15)
B8	0.9269 (3)	0.9739 (7)	0.3719 (4)	0.062 (2)
B9	0.9522 (3)	0.9941 (7)	0.3005 (4)	0.066 (2)
B10	1.0004 (3)	0.8662 (8)	0.3039 (4)	0.063 (2)
B11	1.0071 (3)	0.7697 (7)	0.3781 (3)	0.054 (2)
B12	1.0032 (3)	0.9350 (8)	0.3854 (4)	0.067 (2)
C21	0.9255 (2)	0.5853 (5)	0.3828 (3)	0.0443 (12)
C22	0.9538 (3)	0.4819 (6)	0.3688 (3)	0.067 (2)
C23	0.9452 (3)	0.3632 (7)	0.3909 (4)	0.081 (2)
C24	0.9102 (3)	0.3468 (7)	0.4291 (3)	0.075 (2)
C25	0.8824 (3)	0.4474 (6)	0.4444 (3)	0.072 (2)
C26	0.8902 (3)	0.5678 (6)	0.4216 (3)	0.0586 (15)
C31	0.7567 (2)	0.7094 (5)	0.2406 (3)	0.0473 (12)
C32	0.7342 (2)	0.6259 (6)	0.2761 (3)	0.062 (2)
C33	0.6861 (3)	0.6602 (8)	0.2944 (3)	0.075 (2)
C34	0.6623 (3)	0.7769 (8)	0.2793 (3)	0.077 (2)
C35	0.6838 (2)	0.8634 (7)	0.2435 (3)	0.069 (2)
C36	0.7308 (2)	0.8294 (6)	0.2234 (3)	0.0583 (14)
C41	0.8042 (2)	0.6847 (5)	0.1291 (3)	0.0451 (12)
C42	0.7463 (2)	0.7164 (6)	0.0794 (3)	0.0560 (14)
C43	0.7329 (3)	0.7207 (6)	0.0056 (3)	0.066 (2)
C44	0.7764 (3)	0.6944 (6)	−0.0196 (3)	0.064 (2)
C45	0.8344 (3)	0.6614 (7)	0.0294 (3)	0.072 (2)
C46	0.8478 (2)	0.6537 (7)	0.1021 (3)	0.064 (2)
P1′	0.66505 (5)	0.33925 (13)	0.38240 (6)	0.0393 (3)
C1′	0.6125 (2)	0.2099 (4)	0.3839 (2)	0.0352 (10)
C2′	0.5590 (2)	0.2613 (4)	0.4153 (2)	0.0358 (11)
B3′	0.6168 (2)	0.1571 (6)	0.4664 (3)	0.0402 (12)
B4′	0.6275 (3)	0.0529 (6)	0.4043 (3)	0.0450 (14)
B5′	0.5781 (3)	0.1005 (6)	0.3162 (3)	0.0450 (14)
B6′	0.5375 (2)	0.2366 (6)	0.3236 (3)	0.0405 (13)
B7′	0.5403 (2)	0.1422 (6)	0.4582 (3)	0.0431 (13)
B8′	0.5816 (3)	0.0087 (6)	0.4513 (3)	0.052 (2)
B9′	0.5567 (3)	−0.0256 (6)	0.3570 (3)	0.053 (2)
B10′	0.5011 (3)	0.0863 (6)	0.3064 (3)	0.052 (2)
B11′	0.4901 (2)	0.1911 (6)	0.3689 (3)	0.0452 (14)
B12′	0.5031 (3)	0.0287 (6)	0.3896 (4)	0.055 (2)
C21′	0.5625 (2)	0.3949 (5)	0.4412 (3)	0.0400 (11)
C22′	0.5983 (3)	0.4271 (6)	0.5123 (3)	0.063 (2)
C23′	0.5988 (3)	0.5509 (7)	0.5352 (4)	0.084 (2)
C24′	0.5663 (4)	0.6414 (7)	0.4886 (5)	0.085 (2)
C25′	0.5295 (4)	0.6105 (7)	0.4174 (5)	0.090 (2)
C26′	0.5277 (3)	0.4876 (6)	0.3944 (3)	0.068 (2)
C31′	0.7327 (2)	0.3216 (5)	0.4682 (3)	0.0449 (12)
C32′	0.7680 (2)	0.2141 (6)	0.4961 (3)	0.0602 (15)
C33′	0.8173 (3)	0.2198 (8)	0.5640 (3)	0.074 (2)
C34′	0.8316 (3)	0.3302 (9)	0.6030 (3)	0.083 (2)
C35′	0.7975 (3)	0.4360 (8)	0.5756 (3)	0.078 (2)
C36′	0.7486 (3)	0.4320 (6)	0.5091 (3)	0.060 (2)
C41′	0.6882 (2)	0.2889 (5)	0.3101 (3)	0.0431 (11)
C42′	0.7443 (3)	0.2397 (7)	0.3212 (3)	0.070 (2)
C43′	0.7594 (3)	0.2247 (8)	0.2610 (4)	0.092 (2)
C44′	0.7196 (3)	0.2539 (8)	0.1933 (3)	0.087 (2)
C45′	0.6638 (3)	0.3034 (8)	0.1823 (3)	0.083 (2)
C46′	0.6485 (2)	0.3231 (6)	0.2407 (3)	0.063 (2)

Table 2. Selected geometric parameters (Å, °)

P1—C1	1.883 (5)	P1′—C1′	1.869 (5)
P1—C31	1.837 (5)	P1′—C31′	1.829 (5)
P1—C41	1.835 (5)	P1′—C41′	1.838 (5)
C1—B3	1.733 (7)	C1′—B3′	1.719 (7)
C1—B4	1.709 (8)	C1′—B5′	1.722 (7)
C1—B5	1.720 (7)	C1′—B4′	1.717 (7)
C1—B6	1.717 (7)	C1′—B6′	1.727 (7)
C1—C2	1.755 (6)	C1′—C2′	1.732 (6)
C2—C21	1.503 (7)	C2′—C21′	1.500 (7)
C2—B3	1.732 (7)	C2′—B3′	1.739 (7)
C2—B6	1.735 (7)	C2′—B6′	1.725 (6)
C2—B7	1.700 (8)	C2′—B7′	1.692 (7)
C2—B11	1.705 (7)	C2′—B11′	1.698 (7)
B3—B4	1.777 (8)	B3′—B4′	1.766 (8)
B3—B7	1.794 (8)	B3′—B7′	1.775 (7)

B3—B8	1.768 (9)	B3'—B8'	1.753 (8)
B4—B5	1.765 (9)	B4'—B5'	1.755 (8)
B4—B8	1.778 (10)	B4'—B8'	1.782 (8)
B4—B9	1.779 (9)	B4'—B9'	1.772 (8)
B5—B6	1.776 (9)	B5'—B6'	1.779 (8)
B5—B9	1.761 (10)	B5'—B9'	1.753 (9)
B5—B10	1.767 (8)	B5'—B10'	1.774 (8)
B6—B10	1.772 (10)	B6'—B10'	1.781 (9)
B6—B11	1.801 (9)	B6'—B11'	1.789 (8)
B7—B8	1.772 (10)	B7'—B8'	1.764 (9)
B7—B11	1.768 (9)	B7'—B11'	1.783 (7)
B7—B12	1.777 (10)	B7'—B12'	1.773 (9)
B8—B9	1.789 (10)	B8'—B9'	1.782 (9)
B8—B12	1.779 (10)	B8'—B12'	1.787 (9)
B9—B10	1.764 (11)	B9'—B10'	1.761 (9)
B9—B12	1.760 (10)	B9'—B12'	1.760 (9)
B10—B11	1.765 (9)	B10'—B11'	1.780 (8)
B10—B12	1.775 (11)	B10'—B12'	1.769 (9)
B11—B12	1.766 (10)	B11'—B12'	1.771 (9)
C41—P1—C31	105.2 (2)	C41'—P1'—C31'	106.5 (2)
C31—P1—C1	102.6 (2)	C31'—P1'—C1'	104.4 (2)
C41—P1—C1	103.6 (2)	C41'—P1'—C1'	102.8 (2)
B4—C1—P1	130.0 (4)	B4'—C1'—P1'	129.9 (3)
B6—C1—P1	113.1 (4)	B3'—C1'—P1'	118.7 (3)
B5—C1—P1	127.6 (3)	B5'—C1'—P1'	126.0 (3)
B3—C1—P1	116.6 (3)	B6'—C1'—P1'	112.2 (3)
C2—C1—P1	111.0 (3)	C2'—C1'—P1'	112.1 (3)

Data collection: *XSCANS* (Siemens, 1994). Cell refinement: *XSCANS*. Data reduction: *XSCANS*. Program(s) used to solve structure: *SHELXTL* (Sheldrick, 1994). Program(s) used to refine structure: *SHELXTL*. Molecular graphics: *SHELXTL*. Software used to prepare material for publication: *SHELXTL*.

The authors thank Heriot-Watt University for the provision of a Postdoctoral Fellowship (to GMR) and a University Studentship (to MAM), the British Council for an Acciones Integradas award and the Callery Chemical Company for a generous gift of B₁₀H₁₄.

Lists of structure factors, anisotropic displacement parameters, H-atom coordinates and complete geometry have been deposited with the IUCr (Reference: CF1123). Copies may be obtained through The Managing Editor, International Union of Crystallography, 5 Abbey Square, Chester CH1 2HU, England.

References

- Brain, P. T., Cowie, J., Donohoe, D. J., Hnyk, D., Rankin, D. W. H., Reed, D., Reid, B. D., Robertson, H. E., Welch, A. J., Hofmann, M. & von R. Schleyer, P. (1996). *Inorg. Chem.* **35**, 1701–1708.
- Cowie, J., Reid, B. D., Watmough, J. M. S. & Welch, A. J. (1994). *J. Organomet. Chem.* **481**, 283–293.
- Kivekäs, R., Sillanpää, R., Teixidor, F., Viñas, C. & Nuñez, R. (1994). *Acta Cryst.* **C50**, 2027–2030.
- Kivekäs, R., Teixidor, F., Viñas, C. & Nuñez, R. (1995). *Acta Cryst.* **C51**, 1868–1870.
- Lewis, Z. G. & Welch, A. J. (1993). *Acta Cryst.* **C49**, 705–708.
- Reid, B. D. (1992). PhD thesis, University of Edinburgh, Scotland.
- Sheldrick, G. M. (1994). *SHELXTL*. Siemens Analytical X-ray Instruments Inc., Madison, Wisconsin, USA.
- Siemens (1994). *XSCANS*. X-ray Single Crystal Analysis System. Version 2.1. Siemens Analytical X-ray Instruments Inc., Madison, Wisconsin, USA.
- Thomas, R. L., Rosair, G. M. & Welch, A. J. (1996). *Acta Cryst.* **C52**, 1024–1026, and references therein.

Acta Cryst. (1996). **C52**, 3138–3140

Tetraethyl (1Z,3Z)-Buta-1,3-diene-1,2,3,4-tetracarboxylate

KAZUHIDE TANI,* TSUNEAKI YAMAGATA AND YASUTAKA KATAOKA

Department of Chemistry, Faculty of Engineering Science, Osaka University, Toyonaka, Osaka 560, Japan. E-mail: yamagata@pxews1.protein.osaka-u.ac.jp



(Received 20 May 1996; accepted 8 July 1996)

Abstract

The title compound, C₁₆H₂₂O₈, was obtained as the product of a cationic Rh^I-catalyzed hydridodimerization of diethyl acetylenedicarboxylate in methanol. The molecule lies on a crystallographic inversion centre. The diene moiety of the molecule possesses *s-trans* and *Z,Z* geometry.

Comment

The title compound, (1), has been synthesized by non-catalytic reactions (Bruce, Koutsantonis, Tiekink & Nicholson, 1991; Ryu, Kusumoto, Ogawa, Kambe & Sonoda, 1989; Neunhoeffer, Lehmann & Ewald, 1977); the *Z,Z* geometry, however, has been reported without X-ray analysis. We found an excellent cationic rhodium(I) catalyst containing the (*S*)-(–)-2,2'-bis-(diphenylphosphino)-1,1'-binaphthyl [(*S*)-binap] ligand which exhibited high catalytic activity for the hydridodimerization of acetylenic compounds. The synthetic details will be published elsewhere. The title compound was prepared according to the scheme shown below.



The reaction product was purified by distillation using a Kuhrgelrohr apparatus and subsequent recrystalliza-