

View Article Online View Journal

# ChemComm

# Accepted Manuscript

This article can be cited before page numbers have been issued, to do this please use: S. Bakthavatsalam, A. Sarkar, A. Rakshit, S. Jain, A. Kumar and A. Datta, *Chem. Commun.*, 2014, DOI:



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about *Accepted Manuscripts* in the **Information for Authors**.

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard <u>Terms & Conditions</u> and the <u>Ethical guidelines</u> still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.



www.rsc.org/chemcomm

### ChemComm

# **RSCPublishing**

## COMMUNICATION

Journal Name

# **Tuning Macrocycles to Design 'Turn-on' Fluorescence Probes for Manganese (II) Sensing in Live Cells**

Cite this: DOI: 10.1039/x0xx00000x

Subha Bakthavatsalam†, Anindita Sarkar†, Ananya Rakshit, Shubhi Jain, Amit Kumar and Ankona Datta $^{\ast}$ 

Received 00th January 2014, Accepted 00th January 2014

DOI: 10.1039/x0xx00000x

www.rsc.org/

Published on 23 December 2014. Downloaded by Old Dominion University on 23/12/2014 16:09:42.

We tune the coordination environment of macrocyclic ligands to design two novel fluorescence sensors for  $Mn^{2+}$ . The BODIPY-based  $Mn^{2+}$  sensor M1 affords an excellent, 52 fold, fluorescence 'turn-on' response despite the paramagnetic nature of  $Mn^{2+}$ . The lipophilic probe is cell-permeable and confocal imaging demonstrates that the sensor distinctly detects  $Mn^{2+}$  within live cells.

Mn<sup>2+</sup> is essential for biological systems as a cofactor for enzymes in its bound form and as an antioxidant when labile.<sup>1-3</sup> Mn-containing compounds also find extensive applications in industrial processes.<sup>4</sup> Pesticides, anti-knock agents, and infant formula contain Mn in different forms.<sup>4</sup> Exposure through these sources however, can lead to Mn accumulation and neuronal defects in biological systems.<sup>4-6</sup> The subtle balance between essentiality and toxicity is extremely intriguing, and there are many open questions related to the cellular effects of Mn<sup>2+</sup> and factors maintaining homeostasis.<sup>3, 7</sup> A key requirement for attempts towards elucidating cellular mechanisms is obtaining information on the *in cellulo* localization of Mn<sup>2+</sup> both under physiological and overexposure conditions.

Optical sensing using fluorescence is ideal for detecting trace metals because of its high sensitivity.<sup>8</sup> Justifiably multiple groups have reported various elegant fluorescent sensors for transition metal ions.<sup>8-24</sup> Mn<sup>2+</sup>, however remains elusive, due to the lack of Mn-selective ligands and the paramagnetic nature of Mn<sup>2+</sup> which leads to fluorescence quenching. Hence, there are only few reports on fluorescence sensors for Mn<sup>2+.25-31</sup> We

Department of Chemical Sciences, Tata Institute of Fundamental Research, 1 Homi Bhabha Road, Colaba, Mumbai- 400005, India.

† Equal Contribution

Electronic Supplementary Information (ESI) available: Synthetic procedures, characterization, and spectra. See DOI: 10.1039/c000000x/ have therefore critically assessed the coordination preferences of  $Mn^{2+}$  to design  $Mn^{2+}$  specific ligands and report herein the synthesis and properties of  $Mn^{2+}$  selective 'turn-on' fluorescent sensors, manganese sensor 1 (M1) and manganese sensor 2 (M2). The BODIPY-based lipophilic sensors have visible excitation and emission profiles and display 52-fold and 28-fold fluorescence 'turn-on' response upon  $Mn^{2+}$  binding, for M1 and M2 respectively. M1 displayed higher selectivity towards  $Mn^{2+}$  sensing and was successfully applied for *in cellulo* imaging of  $Mn^{2+}$  using confocal microscopy.

Mn<sup>2+</sup> lies in the bottom of the Irving-Williams series that predicts the relative stabilities of complexes formed by divalent transition metal ions.<sup>32</sup> Hence, designing Mn<sup>2+</sup> specific ligands for sensor development proves to be extremely challenging. Canary and co-workers have reported that ligands containing mixed oxygen and nitrogen donor atoms might be suitable for Mn<sup>2+</sup> binding.<sup>29</sup> A comprehensive body of work on Mn<sup>2+</sup> complexes of penta-aza macrocycles acting as superoxide dismutase mimetics reported by Riley and co-workers and other groups indicate that Mn<sup>2+</sup> ions prefer pentagonal bi-pyramidal coordination geometry.<sup>33, 34</sup> We therefore hypothesize that penta-aza macrocyclic motifs with oxygen containing arms can be optimized for selective Mn<sup>2+</sup> sensing over other divalent metal ions (Scheme1).



Scheme 1 Design strategy for 'turn-on' Mn<sup>2+</sup> sensing.

ChemComm

Page 2 of 4 View Article Online DOI: 10.1039/C4CC09542A



In order to achieve efficient 'turn-on'  $Mn^{2+}$  sensing a dimethylaniline moiety is strategically fused with the ligand. The dimethylaniline moiety and the additional N-donors in the ligand will quench the fluorescence of the covalently attached BODIPY via photo-induced electron transfer (PET) mechanism in the absence of  $Mn^{2+}$ . Upon  $Mn^{2+}$  binding to the macrocycle, the fluorophore will afford an increase in fluorescence intensity (Scheme 1).

An important requirement for  $Mn^{2+}$  sensors is the selectivity for  $Mn^{2+}$  over  $Ca^{2+}$  due to the high abundance of  $Ca^{2+}$  *in cellulo*.<sup>35</sup> Specifically, anionic oxygen donors will increase  $Ca^{2+}$ binding. Hence, we include methyl ester groups as oxygen donors in the ligand design. The ratio of nitrogen to oxygen donor atoms is varied to design M1 and M2 with the aim of tuning  $Mn^{2+}$  affinity and selectivity.

The synthesis of M1 and M2 is outlined in Scheme 2. Richman-Atkins cyclization is used to synthesize the two novel differentially substituted N-tosyl-protected macrocycles 3a and 3b in 29 % and 47 % yields respectively, starting from compounds 1 and 2. The synthesis of the macrocycle precursors, compounds 1 and 2 are described in the supplementary information. The macrocycles are deprotected in the presence of sodium naphthalenide following which methyl acetate arms are attached on the free secondary amines with yields of 45 % and 21 % for compounds 4a and 4b, respectively. Vilsmeier-Haack reaction is used to generate aldehydes 5a in 33 % and 5b in 31 % yields. Condensation reactions of the aldehyde-containing macrocycles 5a and 5b with 2,4-dimethylpyrrole, followed by treatment with BF<sub>3</sub> Et<sub>2</sub>O affords M1 in 29 % yield and M2 in 14 % yield following purification.

M1 and M2 have ester arms which increase their lipophilicity. Hence, *in vitro* absorbance and fluorescence experiments to examine the 'turn-on' response of the probes were conducted in acetonitrile. Metal-unbound aka 'apo' M1 and M2 both display a single absorption maxima at 496 nm ( $\epsilon = 6.9 \times 10^4 \text{ M}^{-1} \text{ cm}^{-1}$  for M1 and M2, Figure S3). The emission maxima for apo M1 and M2 were at 508 and 507 nm ( $\lambda_{ex} = 480$  nm), respectively (Figures 1 and S2). Upon Mn<sup>2+</sup> addition neither M1 nor M2 showed any significant shifts in either the absorption ( $\lambda_{abs}=497$  nm,  $\epsilon = 7.1 \times 10^4 \text{ M}^{-1} \text{ cm}^{-1}$  for M1 and M2, Figure S3) or emission maxima ( $\lambda_{em} = 508$  nm for M1 and M2, Figures 1 and S2).



**Figure 1** Fluorescence response of M1 (2.5  $\mu$ M) to Mn<sup>2+</sup> (0 - 48  $\mu$ M) in acetonitrile.  $\lambda_{ex}$  = 480 nm. (inset) Image depicting quartz cuvettes containing M1 (10  $\mu$ M, left) and M1 (10  $\mu$ M) in presence of Mn<sup>2+</sup> (20  $\mu$ M, right). Both cuvettes were illuminated equally under long-wave UV radiation ( $\lambda_{ex}$  = 365 nm).

Both M1 and M2 displayed excellent fluorescence 'turn-on' response in the presence of  $Mn^{2+}$ . The apo forms of M1 and M2 had very low quantum yields ( $\phi = 0.0038$  for M1 and 0.0055

Published on 23 December 2014. Downloaded by Old Dominion University on 23/12/2014 16:09:42.

for M2) due to effective quenching of the BODIPY fluorophore via the dimethylaniline moiety fused to the penta-aza macrocycle (Figures 1 and S2). Upon addition of  $Mn^{2+}$ , M1 afforded a 52 fold emission enhancement ( $\phi = 0.197$ , Figure 1 and 1 inset), while M2 afforded a 28 fold emission enhancement (( $\phi = 0.156$ , Figure S2).



**Figure 2** Observed fluorescence intensity (F) over initial fluorescence intensity in the absence of any metal ion ( $F_0$ ) for (a) M1 (2.5  $\mu$ M) and (b) M2 (2.5  $\mu$ M) in the presence of various biologically relevant metal ions ([ $M^{n+}$ ]= 20  $\mu$ M) at 508 nm, in acetonitrile.  $\lambda_{ex}$  = 480 nm. Insets: F/F<sub>0</sub> at 508 nm for (a) M1 and (b) M2 in the presence of Mn<sup>2+</sup> (blue squares, fitted data red line), Ca<sup>2+</sup> (orange circles), and Fe<sup>2+</sup> (green triangles) over a [ $M^{n+}$ ] range (0-48  $\mu$ M).

The fluorescence binding plots (Figures 2a inset and 2b inset) were fitted to obtain apparent dissociation constants for  $Mn^{2+}$  binding to M1 and M2 in acetonitrile, which were  $4.0 \pm 0.9 \ \mu$ M and  $8.3 \pm 0.4 \ \mu$ M respectively. The plot for M1 could be best fitted to a 1:1 binding model (Equation S1) with no cooperativity as indicated by the corresponding Hill plot (Figure S4a). On the other hand, the binding plot for M2 was best fitted to a 1:2 (M2:Mn<sup>2+</sup>) binding model (Equation S2). A Hill plot in this case displayed cooperative binding (Figure S4b), hinting at a possible second Mn<sup>2+</sup> binding site due to the presence of additional methyl acetate arms.

We probed the selectivity of M1 and M2 towards other biologically relevant metal ions by performing fluorescence titration experiments.  $F/F_0$  values for the probes at the emission maxima, 508 nm, have been plotted for various metal ions in Figures 2a and 2b. Amongst the ions tested, both M1 and M2 afforded maximum fluorescence enhancement in the presence of Mn<sup>2+</sup> and the fluorescence response saturated above 20  $\mu$ M Mn<sup>2+</sup> concentration (Figures 2a inset and 2b inset). Importantly, Ca<sup>2+</sup>, a major competitor for Mn<sup>2+</sup> binding ligands, did not display any fluorescence enhancement with M1(Figures 2a and 2a inset) and only a modest 2.6 times increase with M2 (Figures 2b and 2b inset). Similarly, another paramagnetic metal ion  $Fe^{2+}$  did not afford any increase in emission in the presence of either sensor (Figures 2a and 2b).

M1 afforded better selectivity towards Mn<sup>2+</sup> when compared to M2. The only other biologically relevant metal ions that afforded a response with M1 were  $\mathrm{Zn}^{2+}$  and  $\mathrm{Cu}^{2+}$ (Figure 2a). Since, intracellular Cu is mostly present as Cu<sup>+</sup> in the reducing intracellular environment, interference from Cu<sup>2+</sup> during in cellulo Mn<sup>2+</sup> imaging is not expected.<sup>21</sup> Also, intracellular labile  $Zn^{2+}$  is in the pM level, while the putative labile  $Mn^{2+}$  is predicted to be in the sub  $\mu M$  level.<sup>35-37</sup> Competition experiments performed by adding Mn<sup>2+</sup> following additions of Zn<sup>2+</sup> or Cu<sup>2+</sup> to either probe indicated further increase in fluorescence emission (Figure S5). Hence, M1 and M2 respond to Mn<sup>2+</sup> even in the presence of these interfering ions, proving higher Mn<sup>2+</sup> specificity. Another metal ion that responded to the sensors was Cd<sup>2+</sup> (Figure S6). However Cd<sup>2+</sup> is highly toxic to cells and is not generally present in labile form.<sup>35</sup> Hence, we do not expect any interference due to this ion in the cellular milieu.



**Figure 3** Live-cell imaging of intracellular Mn<sup>2+</sup> by confocal microscopy in HEK 293T cells. Overlay of confocal and transmission images of cells treated with (a) M1 (b) MnCl<sub>2</sub> (25  $\mu$ M) in growth media followed by M1 (c) M1 followed by TPEN (2mM) (d) MnCl<sub>2</sub> followed by M1 and TPEN.  $\lambda_{ex}$  488 nm (e-g) Confocal images depicting co-localization of M1 (green channel,  $\lambda_{ex}$  488 nm) with Nile Red (100 nM, red channel,  $\lambda_{ex}$  543 nm) in Mn-treated cells. M1 concentration 5  $\mu$ M. Scale: 25  $\mu$ m (a-d), 10  $\mu$ m (e-g).

The probes are lipophilic and hence would be applicable for *in cellulo* imaging of  $Mn^{2+}$  in lipid-rich bodies and organelles.

ChemComm

**ChemComm Accepted Manuscript** 

In order to test the in cellulo response we chose M1 since it had higher selectivity for Mn<sup>2+</sup>. M1 was dissolved in DMSO and diluted in Thomson's buffer (< 1 % DMSO in buffer, pH 7.4) for the cell experiments. All experiments and imaging were performed on live HEK 293T cells. The probe readily entered cells within 15 min of incubation, as shown in the confocal images in figure 3. Cells treated with MnCl<sub>2</sub> for 1h showed significant fluorescence enhancement when compared to control untreated cells (Figures 3 a, b). To test whether the fluorescence increase was due to Mn<sup>2+</sup>, tetrakis-(2pyridylmethyl)ethylene-diamine (TPEN) a cell-permeable chelator of divalent metal ions was added to the Mn-treated cells. The emission intensity clearly decreased (Figure 3d) proving that the increase in emission intensity was a Mn<sup>2+</sup> selective response. The fluorescence intensity in control cells also decreased upon TPEN treatment indicating that the sensor would also be able to detect labile Mn<sup>2+</sup> under physiological conditions (Figures 3c).

Finally, to check the *in cellulo* localization of the sensors we performed co-localization experiments with Nile Red, a cell-permeable lipophilic dye known to bind to lipid membranes.<sup>38</sup> The confocal images indicate that M1 co-localizes with Nile Red (Figure 3e-g), implying that the sensor targets lipid vesicles and senses  $Mn^{2+}$  in cellular bodies with high lipid-content.

In this work we have successfully shown that penta-aza macrocycles can be optimally decorated with oxygen containing arms to develop selective 'turn-on' fluorescent sensors for  $Mn^{2+}$ . The sensor M1 shows significant increase in fluorescence intensity in  $Mn^{2+}$ -loaded cells and can detect  $Mn^{2+}$  within lipid rich entities. Selective detection of  $Mn^{2+}$  is distinctly confirmed by reduction in intensity upon TPEN treatment in  $Mn^{2+}$ -treated cells. Recent micro-SXRF and XANES studies on  $Mn^{2+}$  localization within the cell indicate  $Mn^{2+}$  accumulation in the Golgi apparatus with possible removal of excess  $Mn^{2+}$  via exocytosis through lipid vesicles.<sup>39</sup> In the future we plan to use our modular synthetic strategy of attaching arms to penta-aza macrocycles to modify and target our sensors to the Golgi, a major site implicated in  $Mn^{2+}$  homeostasis.<sup>40</sup>

A.D. acknowledges support from Department of Atomic Energy, India, The authors acknowledge Prof. Sudipta Maiti and the Biophotonics group, TIFR, for help with cell culture and confocal imaging; NMR Facility, TIFR, India; and Mass laboratory, Chemistry Department, I.I.T. Bombay, India.

### Notes and references

- K. Barnese, E. B. Gralla, J. S. Valentine and D. E. Cabelli, *Proc. Natl. Acad. Sci. U.S.A.*, 2012, **109**, 6892-6897.
- 2 R. L. McNaughton, A. R. Reddi, M. H. S. Clement, A. Sharma, K. Barnese, L. Rosenfeld, E. B. Gralla, J. S. Valentine, V. C. Culotta and B. M. Hoffman, *Proc. Natl. Acad. Sci. U.S.A.*, 2010, 107, 15335-15339.
- 3 A. R. Reddi, L. T. Jensen and V. C. Culotta, *Chem. Rev.*, 2009, 109, 4722-4732.

- 4 A. B. Bowman, G. F. Kwakye, E. H. Hernandez and M. Aschner, J. Trace Elem. Med. Biol., 2011, 25, 191-203.
- 5 J. Kaiser, Science, 2003, 300, 926-928.
- 6 S. Bakthavatsalam, S. Das Sharma, M. Sonawane, V. Thirumalai and A. Datta, *Dis Model Mech.*, 2014, 7, 1239-1251.
- 7 C. E. Blaby-Haas and S. S. Merchant, J. Biol. Chem., 2014, 289, 28129-28136.
- 8 K. P. Carter, A. M. Young and A. E. Palmer, *Chem. Rev.*, 2014, **114**, 4564-4601.
- 9 E. M. Nolan and S. J. Lippard, Acc. Chem. Res., 2009, 42, 193-203.
- 10 Y. Xu, L. Xiao, S. Sun, Z. Pei, Y. Pei and Y. Pang, Chem. Commun., 2014, 50, 7514-7516.
- 11 S. Pal, N. Chatterjee and P. K. Bharadwaj, *RSC Adv.*, 2014, **4**, 26585-26620.
- 12 M. Li, H. Ge, R. L. Arrowsmith, V. Mirabello, S. W. Botchway, W. Zhu, S. I. Pascu and T. D. James, *Chem. Commun*, 2014, **50**, 11806-11809.
- 13 Y. H. Lee, N. Park, Y. B. Park, Y. J. Hwang, C. Kang and J. S. Kim, *Chem. Commun.*, 2014, **50**, 3197-3200.
- 14 M. L. Giuffrida, E. Rizzarelli, G. A. Tomaselli, C. Satriano and G. Trusso Sfrazzetto, *Chem. Commun.*, 2014, 50, 9835-9838.
- 15 A. Balamurugan, V. Kumar and M. Jayakannan, *Chem. Commun.*, 2014, 50, 842-845.
- 16 C. J. Fahrni, Curr. Opin. Chem. Biol., 2013, 17, 656-662.
- 17 Y. Chen, C. Zhu, J. Cen, J. Li, W. He, Y. Jiao and Z. Guo, *Chem. Commun.*, 2013, 49, 7632-7634.
- 18 Z. Xu, X. Liu, J. Pan and D. R. Spring, Chem. Commun., 2012, 48, 4764-4766.
- 19 A. M. Kim, S. Vogt, T. V. O'Halloran and T. K. Woodruff, Nat. Chem. Biol., 2010, 6, 674-681.
- 20 Y. You, E. Tomat, K. Hwang, T. Atanasijevic, W. Nam, A. P. Jasanoff and S. J. Lippard, *Chem. Commun.*, 2010, 46, 4139-4141.
- 21 E. L. Que, D. W. Domaille and C. J. Chang, Chem. Rev., 2008, 108, 1517-1549.
- 22 J. Liu and Y. Lu, J. Am. Chem. Soc., 2007, 129, 9838-9839.
- 23 J. L. Bricks, A. Kovalchuk, C. Trieflinger, M. Nofz, M. Buschel, A. I. Tolmachev, J. Daub and K. Rurack, J. Am. Chem. Soc., 2005, 127, 13522-13529.
- 24 M. D. Shults, D. A. Pearce and B. Imperiali, J. Am. Chem. Soc., 2003, 125, 10591-10597.
- 25 X. Mao, H. Su, D. Tian, H. Li and R. Yang, ACS Appl. Mater. Interfaces, 2013, 5, 592-597.
- 26 K. Dutta, R. C. Deka and D. K. Das, J. Fluorescence, 2013, 23, 1173-1178.
- 27 F. Gruppi, J. Liang, B. B. Bartelle, M. Royzen, D. H. Turnbull and J. W. Canary, *Chem. Commun.*, 2012, 48, 10778-10780.
- 28 I. Kaya, M. Yildirim and M. Kamaci, Synth. Met., 2011, 161, 2036-2040.
- 29 J. Liang and J. W. Canary, Angew. Chem. Int. Ed., 2010, 49, 7710-7713.
- 30 Y. J. Lee, C. Lim, H. Suh, E. J. Song and C. Kim, Sens. Actuators, B, 2014, 201, 535-544.
- 31 B. R. Jali, K. Masud and J. B. Baruah, Polyhedron, 2013, 51, 75-81.
- 32 H. Irving and R. J. P. Williams, J. Am. Chem. Soc., 1953, 3192-3210.
- 33 D. Lieb, I. Kenkell, J. L. Miljkovic, D. Moldenhauer, N. Weber, M. R. Filipovic, F. Groehn and I. Ivanovic-Burmazovic, *Inorg. Chem.*, 2014, 53, 1009-1020.
- 34 D. Salvemini, D. P. Riley and S. Cuzzocrea, Nat. Rev. Drug Discov., 2002, 1, 367-374.
- 35 L. A. Ba, M. Doering, T. Burkholz and C. Jacob, *Metallomics*, 2009, 1, 292-311.
- 36 R. A. Bozym, R. B. Thompson, A. K. Stoddard and C. A. Fierke, ACS Chem. Biol., 2006, 1, 103-111.
- 37 C. E. Outten and T. V. O'Halloran, Science, 2001, 292, 2488-2492.
- 38 P. Greenspan, E. P. Mayer and S. D. Fowler, J. Cell Biol., 1985, 100, 965-973.
- 39 A. Carmona, S. Roudeau, L. Perrin, G. Veronesi and R. Ortega, *Metallomics*, 2014, 6, 822-832.
- 40 S. Mukhopadhyay and A. D. Linstedt, Proc. Natl. Acad. Sci. U.S.A., 2011, 108, 858-863.