Photochemistry of heteroaromatics-a novel photocycloaddition of 2-alkoxy-3-cyanopyridines with methacrylonitrile

Masami Sakamoto,*a Takeru Sano,a Masaki Takahashi,b Kentaro Yamaguchi,c Tsutomu Fujitaa and Shoji Watanabea

^a Department of Applied Chemistry, Faculty of Engineering, Chiba University, Yayoi-cho, Inage-ku, Chiba, Japan 263

^b Graduate School of Science and Technology, Chiba University, Yayoi-cho, Inage-ku, Chiba, Japan 263

^c Analytical Center, Chiba University, Yayoi-cho, Inage-ku, Chiba, Japan 263

The photochemical cycloaddition of 2-alkoxy-3-cyano-4,6dimethylpyridine with methacrylonitrile gives a bicyclic azetine, 6-alkoxy-3,5-dicyano-2,5,8-trimethyl-7-azabicyclo-[4.2.0]octa-2,7-diene (45-55% yield) and 3-acetyl-4amino-1,5-dicyano-2,5-dimethylcyclohexa-1,3-diene (15-17% yield).

The photochemical cycloaddition of aromatic compounds has received much attention from both mechanistic and synthetic perspectives.^{1,2} In particular, the photochemical [2 + 2] and [3+ 2] addition of benzene and its derivatives to various kinds of alkenes has been investigated extensively.^{3,4} Relatively little is known about the photochemical reaction of heteroaromatics in the presence of alkenes.^{5,6} Here we report that the photolysis of 2-alkoxy-3-cyano-4,6-dimethylpyridine in the presence of methacrylonitrile promotes [2 + 2] cycloaddition from the singlet excited state.

We previously reported that the photolysis of a benzene solution containing 3-cyano-2-methoxy-4,6-dimethylpyridine 1a resulted in a 92% yield of the transpositional isomer 2a (Table 1, entry 1).^{7,8} However, irradiation in the presence of alkenes resulted in considerably different photochemical behaviour. When an argon-purged benzene solution of 1a and methacrylonitrile (1.0 mol dm^{-3}) in a Pyrex vessel was irradiated using a high pressure mercury lamp and the photolysate was subjected to chromatography in silica gel, a bicyclic azetine, 3,5-dicyano-6-methoxy-2,5,8-trimethyl-7-azabicyclo[4.2.0]octa-2,7-diene 3a (45% yield) and 3-acetyl-4-amino-1,5-dicyano-2,5-dimethylcyclohexa-1,3-diene 4a (17% yield) were obtained and the yield of 2a decreased (Table 1, entry 2).

The chemical structure of 3a was determined by spectroscopy. The mass spectrum (FAB) showed a molecular ion peak

Table 1 Photolysis of 2-alkoxy-3-cyano-4,6-dimethylpyridine 1 in the presence of methacrylonitrile

Me 1a R b R	$ \underbrace{\bigvee_{N=0}^{CN}}_{OR}^{CN} + \underbrace{\bigvee_{0}^{N}}_{C}^{N} $ $ = Me $ $ = Et $	he hv NC benzene N Me	Me N OR Me		Me ⁺ Me CN		9
				Yield (%) ^c			
Entry	Pyridine 1 ^a	Conc. of Alkene	Conv. (%) ^b	2	3	4	
1	a	0	54	92	_	_	_
2	а	1.0	72	8	45	17	
3	b	0	58	90			
4	b	1.0	80	6	55	15	

^a Concentration of pyridines was 0.02 mol dm⁻³. ^b All samples were irradiated using a 1 kW high-pressure mercury lamp in an argon atmosphere through a Pyrex filter for 1.5 h. c Yields were determined based on consumed pyridines 1.

at 230 (MH⁺ calculated for $C_{13}H_{16}N_3O$), indicating that 3a is a 1:1 adduct of 1a with methacrylonitrile. The ¹H NMR displayed methylene protons at δ 2.22 and 2.64 (ABq, J 15.8 Hz, 2 H, 4-CH₂) and a methine proton at δ 3.88 (s, 1 H, 1-CH) in addition to four singlet peaks derived from methyl protons at δ 1.57 (s, 3 H, 5-Me), 2.10 (s, 3 H, 8-Me), 2.19 (s, 3 H, 2-Me) and 3.43 (s, 3 H, OMe). The ¹³C NMR exhibited sp² carbon peaks at δ 188.0 (s, 8-C=N), 149.0 (s, 2-C) and 106.9 (s, 3-C) in addition to the peaks derived from methylene carbon at δ 35.2 (4-C), the methine group at δ 53.7 (d, 1-C), and singlet peaks at δ 120.5 (s, 3-CN), 116.9 (s, 5-CN), 91.9 (s, 6-C) and 39.7 (s, 5-C).

The structure of cyclohexadiene derivative 4a was unequivocally established by X-ray crystallography (Fig. 1).† In



Fig. 1 ORTEP drawing of 4a. Selected bond lengths (Å) and angles (°); O(1)-C(7) 1.237 (7), N(2)-C(12) 1.120(8), N(3)-C(11) 1.148(9), N(4)-C(1) 1.334(8), C(1)-C(2) 1.386(8), C(1)-C(6) 1.537(8), C(2)-C(3) 1.463(9), C(2)-C(7) 1.457(8), C(3)-C(4) 1.337(9), C(3)-C(10) 1.52(1), C(4)–C(5) 1.490(9), C(4)–C(11) 1.431(9), C(5)–C(6) 1.537(9), C(6)–C(9) 1.503(9), C(6)–C(12) 1.461(9), C(7)–C(8) 1.499(10); N(4)–C(1)–C(2) 123.8(6), N(4)-C(1)-C(6) 116.5(6), C(2)-C(1)-C(6) 116.5(6), C(2)-C(1)-C(6) 119.7(6), C(1)-C(2)-C(3) 118.4(6), C(1)-C(2)-C(7) 120.0(6), C(3)-C(2)-C(7) 121.5(6), C(2)-C(3)-C(4) 119.5(6), C(2)-C(3)-C(10) 119.4(7), C(4)-C(3)-C(10) 120.7(7), C(3)-C(4)-C(5) 119.3(6), C(3)-C(4)-C(11) 121.6(7), C(5)–C(4)–C(11) 119.1(7), C(4)–C(5)–C(6) 110.8(6), C(1)– C(6)-C(5) 107.3(5), C(1)-C(6)-C(9) 113.4(6), C(1)-C(6)-C(12) 106.8(6), C(5)-C(6)-C(9) 110.5(6), C(5)-C(6)-C(12) 108.9(6), C(9)-C(6)-C(12) 109.8(6), O(1)-C(7)-C(2) 121.8(6), O(1)-C(7)-C(8) 117.2(7), C(2)-C(7)-C(8) 120.6(7), N(3)-C(11)-C(4) 178.1(10), N(2)-C(12)-C(6) 177.8(8).

addition, the hypothetical structure was supported by the fact that 3a was easily hydrolysed to 4a.

Table 2 shows the quantum yields for the cycloaddition of 1a in the presence of methacrylonitrile under various conditions. Cycloaddition was not observed at methacrylonitrile concentrations lower than 0.1 mol dm^{-3} (Table 2, entry 1), and the quantum efficiency increased with increasing alkene concentration (Table 2, entries 1-3). The quantum yield was found to be independent of the solvent polarity (Table 2, entries 3-6), and quenching by 2,5-dimethylhexa-2,4-diene 7 was found to be inefficient (Table 2, entry 7). The addition of alkenes did not affect the absorption spectrum, which showed no interaction between pyridine and alkenes in the ground state. The fluorescence spectrum exhibited emission at 322 nm in cyclohexane. The intensity of this fluorescence decreased after the addition of alkenes; however, new fluorescence emission due to the formation of pyridine-alkene exciplex was not observed. These results indicate that photocycloaddition proceeded from the pyridine singlet excited state.

A mechanism has been postulated for the formation of adducts **3** and **4** that involves [2 + 2] photocycloaddition between the C(2)–C(3) of pyridine ring and the alkenyl double bonds leading to cyclobutane **5**, which easily transforms to azacyclooctatriene **6** (Scheme 1).⁹ Upon excitation with a quantum of light, the azacyclooctatriene ring cyclizes into 8-azabicyclo[4.2.0]octa-4,7-diene **3**, which further hydrolyses into cyclohexadiene **4** during the isolation procedure.

Frontier-MO calculations by the AM1 method available in the MOPAC program help to explain the highly regioselective cycloaddition.‡ Using this method, the orbital energies and coefficients of the singlet excited state of **1a** (LSOMO: orbital energy, -6.56318 eV; coefficients, N-1, 0.04716; C-2, 0.37764; C-3, 0.57750; C-4, 0.00499; C-5, -0.45440; C-6, -0.37039, and HSOMO: orbital energy, -3.55472; coefficients, N-1, 0.14025; C-2, 0.43035; C-3, -0.50828; C-4,

 Table 2 Quantum yield of photochemical reaction of 1a

Entry	Solventa	Conc. of alkene/	$10^{-2} \Phi \cdot b$
Linu y	Sorvent		10 - ¥add
1	C ₆ H ₆	0.1	0 (1.1) ^c
2	C ₆ H ₆	0.2	0.6
3	C ₆ H ₆	1.0	2.7
4	Et ₂ O	1.0	1.9
5	MeCN	1.0	2.6
6	MeOH	1.0	2.7
7	C ₆ H ₆	$1.0 + 7^{d}$	2.6

^{*a*} Concentration of pyridine **1a** was 0.02 mol dm⁻³. ^{*b*} A varelophenone actinometer was used in reference to the 313 nm line. ^{*c*} Photoaddition was not observed. Value in parentheses shows quantum yield for the *trans*-isomerization from **1a** to **2a**. ^{*d*}2,5-Dimethylhexa-2,4-diene **7** (0.05 mol dm⁻³) was added.



Footnotes

† *Crystal data* for **4a**: C₁₂H₁₃N₃O, *M*_W 215, colourless prismatic crystal, monoclinic space group *P*2₁/c, *a* = 9.368(3), *b* = 12.566(2), *c* = 10.203(2) Å, β = 97.13(2)°, *V* = 1191.7(5) Å³, *Z* = 4, ρ = 1.20 g cm⁻³, μ(Cu-Kα) = 6.42 cm⁻¹. Final *R* and *R*_w were 0.075 and 0.081 for 1294 reflections. Atomic coordinates, bond lengths and angles, and thermal parameters have been deposited at the Cambridge Crystallographic Data Centre (CCDC). See Information for Authors, Issue No. 1. Any request to the CCDC for this material should quote the full literature citation and the reference number 182/80.

 \ddagger These results are calculated by 'PASOCON MOPA C/386', which is based on the MOPAC (V6.0 QCPE No. 455), of Toray System Center.

§ The HSOMO coefficient on C-6 position is larger than the coefficient on C-3 position ($-0.54475 v_s -0.50828$). For a more accurate approach, the predicted $\Delta\Delta E$ of the reaction was calculated using the Salem–Klopman equation (see for example I. Fleming, *Frontier Orbital and Organic Chemical Reactions*, Wiley-Interscience New York, 1976). The $\Delta\Delta E$ is proportional to (ca1cb1)² + (ca2cb2)² where cX is the orbital coefficient in the reactive site for the addition reaction of the a1–a2 and the b1–b2 bonds. The value of (cC3cC2')² + (cC2cC1')² is 1.7535×10^{-1} and 1.4922×10^{-1} respectively to that of (cC6cC2')² + (cC5cC1')² for the alternative pathway. These results support that the initial bond formation occurs between C(2)–C(3) bond of 1 and C(1')–C(2') bond of methacrylonitrile leading to the experimentally observed product.

References

- 1 D. Bryce-Smith and A. Gilbert, Tetrahedron, 1976, 32, 1309.
- 2 A. Gilbert, Synthetic Organic Photochemistry, ed. W. H. Horspool, Plenum: New York, 1984.
- 3 P. A. Wender, L. Siggel and J.M. Nuss, Organic Photochemistry, ed. A. Padwa, Marcel Dekker, New York and Basel, 1991, vol. 10, pp. 357.
- 4 J. Cornelisse, Chem. Rev., 1993, 93, 615.
- 5 For the photocycloaddition of pyridines with alkenes, only one example has been reported. Pentafluoropyridine reacts with ethylene at the C(3)–C(4) bond leading to cyclobutane; see M. G. Barlow, D. E. Brown and R. N. Haszeldine, *J. Chem. Soc., Chem. Commun.*, 1977, 669.
- 6 I. Saito, K. Kanehira, K. Shimozono and T. Matsuura, *Tetrahedron Lett.*, 1980, 21, 2737; R. Bernardi, T. Caronna, S. Morrocchi and P. Traldi, *Tetrahedron Lett.*, 1981, 22, 155.
- 7 M. Sakamoto, M. Kimura, T. Fujita, T. Nishio, I. Iida and S. Watanabe, J. Am. Chem. Soc., 1991, 113, 5859; M. Sakamoto, T. Takahashi, M. Kimura, M. Fujihira, T. Fujita, I. Iida, T. Nishio and S. Watanabe, J. Org. Chem., 1994, 59, 5117.
- 8 The formation of the transpositional isomer is reasonably explained in terms of a 1,3-shift via Dewar pyridines or azaprizmanes; see S. Captlain and A. Lablache-Combier, J. Chem. Soc., Chem. Commun., 1970, 1247 and R. D. Chambers and R. Middleton, J. Chem. Soc., Perkin Trans. 1, 1977, 1500.
- 9 Similar transformations are documented for photocycloadducts of benzenes with alkenes, leading to cyclooctatrienes and bicyclo[4.2.0]octadienes; see P. J. Wagner and K. Nahm, J. Am. Chem. Soc., 1987, 109, 4404.

Received, 5th February 1996; Com. 6/00810K