

## Organic Preparations and Procedures International: The New Journal for Organic Synthesis

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/uopp20>

### UNEXPECTED DIECKMANN CONDENSATION IN THE SYNTHESIS OF METHYL 6-(6-METHOXY-2-NAPHTHYL)-6-OXOHXANOATE WITH AN ARYL CADMIUM REAGENT

Carmen Balo <sup>a</sup>, Franco Fernández <sup>a</sup>, Xerardo García-Mera <sup>a</sup> & Carmen López <sup>a</sup>

<sup>a</sup> Departamento de Química Orgánica, Facultad de Farmacia, Universidad de Santiago, E-15706, Santiago de Compostela, SPAIN  
Published online: 11 Feb 2009.

To cite this article: Carmen Balo, Franco Fernández, Xerardo García-Mera & Carmen López (2009) UNEXPECTED DIECKMANN CONDENSATION IN THE SYNTHESIS OF METHYL 6-(6-METHOXY-2-NAPHTHYL)-6-OXOHXANOATE WITH AN ARYL CADMIUM REAGENT, *Organic Preparations and Procedures International: The New Journal for Organic Synthesis*, 32:6, 563-567, DOI: [10.1080/00304940009355951](https://doi.org/10.1080/00304940009355951)

To link to this article: <http://dx.doi.org/10.1080/00304940009355951>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing,

systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

## OPPI BRIEFS

**UNEXPECTED DIECKMANN CONDENSATION IN THE SYNTHESIS  
OF METHYL 6-(6-METHOXY-2-NAPHTHYL)-6-OXOHXANOATE  
WITH AN ARYLCADMIUM REAGENT**


*Submitted by* Carmen Balo, Franco Fernández, Xerardo García-Mera and Carmen López\*  
(04/21/00)

*Departamento de Química Orgánica, Facultad de Farmacia,  
Universidade de Santiago, E-15706-Santiago de Compostela, SPAIN*

Methyl 6-(6-methoxy-2-naphthyl)-6-oxohexanoate (**3**) exhibits anti-inflammatory properties,<sup>1</sup> and has been used both as a fluorescent probe<sup>2</sup> and as an intermediate in the synthesis of other fluorescent molecules used for the study of the polarity of lipid bilayers in biological membranes.<sup>3</sup> The synthesis of this useful molecule in moderate yields (40%) by a Friedel-Crafts reaction has been reported recently.<sup>4</sup> The present work was undertaken in an attempt to improve the synthesis of **3** by reaction of a weakly ionic naphthalene organocadmium derivative with methyl adipoyl chloride (methyl 6-chloro-6-oxohexanoate, **1**) in the hope that addition would occur selectively at the most electrophilic carbonyl group.

The reaction of an excess of the cadmium reagent of **2** with **1** gave, in addition to 6,6'-dimethoxy-2,2'-binaphthyl (4%) and adipic acid (10%), the desired product **3** and a yellow solid **4**, whose structure was established as described below.

Compound **4** showed a highly deshielded D<sub>2</sub>O-exchangeable signal at  $\delta$  14.73 in its <sup>1</sup>H NMR spectrum and reacted with diazomethane to afford a derivative **6**, with an additional signal for a methoxy group at  $\delta$  3.60. However, neither the IR spectrum of **4** nor that of **6** were compatible with these compounds being an acid and its methyl ester respectively. In fact, an authentic sample of 6-(6-methoxy-2-naphthyl)-oxohexanoic acid (**7**), obtained by saponification of **3**, had IR and <sup>1</sup>H NMR spectra very different from those of **4**. A close inspection of their spectroscopic data led us to assign the structures of 2-[hydroxy(6-methoxy-2-naphthyl)methylene]cyclopentanone to **4** and that of its methyl ether to **6**.



Additional support for the structure proposed for compound 4 came from the interesting observation that in CDCl<sub>3</sub> solution, it gave rise to a new molecular species (5), easily detected in both the <sup>1</sup>H and <sup>13</sup>C NMR spectra. The concentration of this new species increased slowly and stabilized after 14 days at a 4/5 ratio of 56:44 (± 2; from <sup>1</sup>H NMR integrals). Although not all the spectral parameters of 5 could be determined, <sup>1</sup>H and <sup>13</sup>C NMR data obtained were sufficient for tentative characterization of this compound as 2-(6-methoxy-2-naphthoyl)cyclopentanone. Thus 4, the only species detected in the solid state (IR), is transformed into an equilibrium mixture of tautomers 4 and 5 soon after dissolved in CDCl<sub>3</sub>.

The formation of the unexpected side-product is attributable to compound 3 having undergone a Dieckmann condensation, *via* the formation of a carbanion at the methylene α to the ketone group, due to the strongly basic medium in which 3 was initially formed. The isolation of 4 in such noteworthy yield (32%, as against 40% for the target compound 3) is especially surprising given the fact that no analogous cyclized side-products have been detected in similar syntheses of its higher or lower side-chain homologues.<sup>7</sup> A plausible explanation for this particular behavior is that Dieckmann condensation of 3 leads to an easy to generate, stable five-membered ring.

Thus, the yield of the coupling reaction between the ester-chloride 1 and the organocadmium derivative of 2 should have been at least 72%. However, attempts made to improve the yield of 3 were unsuccessful. Decreasing the time or the temperature of reaction led to an increase in the isolated amounts of 6-methoxynaphthalene and adipic acid; on the other hand, extending reaction time led to an increase in the Dieckmann condensation side product. Production of 4 in 80% yield was achieved when 3 was refluxed for 4 h in methanolic sodium methoxide.

## EXPERIMENTAL SECTION

Melting points (uncorrected): Reichert Kofler Thermopan. IR (in KBr discs, for solids, or films between NaCl plates, for oils): Perkin Elmer FT-IR 1640. NMR (in CDCl<sub>3</sub> with TMS as internal standard): Bruker AMX-300 (300 MHz for <sup>1</sup>H and 75 MHz for <sup>13</sup>C NMR). EIMS (at 70 eV): Kratos MS-50. Microanalyses: Perkin-Elmer 240B Elemental Analyser. Silica gel 60 (70-230 mesh) for chromatography was from Merck. Reagents and solvents were of commercial grade and were used as received.

**Methyl 6-Chloro-6-oxohexanoate (1).** Thionyl chloride (27 mL, 370 mmol) was slowly added to monomethyl adipate (10.0 g, 60 mmol) at 0° and the mixture was heated at 70° for 1 h. The excess thionyl chloride was removed by warming the mixture *in vacuo* until the oily liquid residue had constant mass (12 g). IR: 1799 (acid chloride C=O), 1732 (ester C=O) cm<sup>-1</sup>.

**Reaction of bis(6-Methoxy-2-naphthyl)cadmium with 1.** Magnesium (3.75 g, 154 mmol) was placed in a flask and the apparatus was flame dried under argon. A solution of **2** (10.0 g, 42.2 mmol) in dry THF (35 mL) was added, followed by an iodine crystal and 3 drops of CH<sub>3</sub>I, and the stirred mixture spontaneously began to reflux. While stirring the mixture, further **2** (21.0 g, 88.6 mmol) in dry THF (75 mL) was added at a rate to maintain the reflux. The mixture was vigorously stirred and heated at reflux for an additional 35 min, and the formation of the Grignard reagent was confirmed by the Gilman test.<sup>8</sup> The reaction mixture was then cooled to 0° in an ice-bath, and finely powdered, oven-dried (110°) CdCl<sub>2</sub> (16.0 g, 87.3 mmol) was added at once and the mixture was vigorously stirred until it gave a negative result in the Gilman test (*ca.* 30 min). The THF was distilled off and residual solvent was removed by co-distillation with dry toluene (50 mL). Additional dry toluene (50 mL) was added to the crude arylcadmium reagent, and ester-chloride **1** (11.5 g, 64.4 mmol) was slowly dripped into the mixture. The reaction was successively heated at reflux for 1 h, cooled to 0° and acidified with 2 N H<sub>2</sub>SO<sub>4</sub>. The organic layer was separated, washed with water and dried (Na<sub>2</sub>SO<sub>4</sub>). The solvent was eliminated to afford a yellow paste which was chromatographed on silica gel, with 8:2 toluene/EtOAc as eluent, to afford 2-methoxynaphthalene (7.8 g) and a complex mixture of other products (16.5 g). Further chromatography of the latter mixture, with DCM as eluent, gave a spectroscopically pure (<sup>1</sup>H NMR) yellow solid which was identified as **4** (5.46 g, 32%). An analytical sample of **4** was obtained by recrystallization from EtOAc. Subsequent elution of the chromatographic column with 8:2 DCM/EtOAc gave a white solid identified as **3** (7.67g, 40%). An analytical sample of **3** was obtained by triple recrystallization from toluene. Further elution of the column with EtOAc left a solid residue from which alkaline extraction allowed the isolation of adipic acid (0.90 g), while recrystallization of the neutral fraction from DCM led to isolation of 6,6'-dimethoxy-2,2'-binaphthyl (0.81 g).

**Methyl 6-(6-Methoxy-2-naphthyl)-6-oxohexanoate (3):** white solid, mp 81-82° (toluene), (*lit.*<sup>4</sup> 81-83°). Spectral data as previously described.<sup>4</sup>

*Anal.* Calcd for C<sub>18</sub>H<sub>20</sub>O<sub>4</sub> (300.3): C, 71.98; H, 6.71. Found: C, 72.06; H, 6.92

**6,6'-Dimethoxy-2,2'-binaphthyl:** white solid, mp 180-181° (DCM). IR: 1624 and 1591 (aromatic C=C) cm<sup>-1</sup>. <sup>1</sup>H NMR: δ 8.26 (1 H, d, *J* = 1.6 Hz, 1-H); 7.83 (1 H, d, *J* = 9.0 Hz, 8-H); 7.74 (1 H, d, *J* = 8.7 Hz, 4-H); 7.47 (1 H, dd, *J* = 8.7, 1.6 Hz, 3-H); 7.23 (1 H, dd, *J* = 9.0, 2.5 Hz, 7-H); 7.12 (1 H, d, *J* = 2.5 Hz, 5-H); 3.92 (3 H, s, OCH<sub>3</sub>). EIMS: *m/z* (%) 102 (13), 114 (27), 145 (32), 173 (43), 189 (25), 271 (14), 299 (25), 314 (100, M).

*Anal.* Calcd for C<sub>22</sub>H<sub>18</sub>O<sub>2</sub> (314.4): C, 84.05; H, 5.77. Found: C, 83.90; H, 5.88

**2-[Hydroxy(6-methoxy-2-naphthyl)methylene]cyclopentanone (4):** yellow solid, mp 123-124° (EtOAc). IR: 1626 (C=O), 1585 (C=C) cm<sup>-1</sup>. <sup>1</sup>H NMR: δ 14.73 (1 H, br s, OH); 8.21 (1 H, s, 1'-H);

7.84 (1 H, dd,  $J = 8.7, 1.7$  Hz, 3'-H); 7.80 (1 H, d,  $J = 9.0$  Hz, 8'-H); 7.77 (1 H, d,  $J = 8.7$  Hz, 4'-H); 7.19 (1 H, dd,  $J = 9.0, 2.3$  Hz, 7'-H); 7.14 (1 H, d,  $J = 2.3$  Hz, 5'-H); 3.94 (3 H, s, 6'-OCH<sub>3</sub>); 2.98 (2 H, t,  $J = 7.1$  Hz, 3-H<sub>2</sub>); 2.52 (2 H, t,  $J = 7.9$  Hz, 5-H<sub>2</sub>); 2.00 (2 H, quintet,  $J = 7.5$  Hz, 4-H<sub>2</sub>). <sup>13</sup>C NMR:  $\delta$  210.35, 169.31, 159.48, 136.15, 130.87, 129.91, 129.14, 128.27, 127.07, 125.28, 119.87, 109.31, 105.86, 55.69, 37.79, 28.90, 21.58.

*Anal.* Calcd for C<sub>17</sub>H<sub>16</sub>O<sub>3</sub> (268.3): C, 76.10; H, 6.01. Found: C, 75.89; H, 5.86

**2-(6-Methoxy-2-naphthoyl)cyclopentanone (5):** tautomeric form of **4**, detected by <sup>1</sup>H and <sup>13</sup>C NMR. Equilibrium between **5** and **4**, at a 44:56 ratio, is reached after 14 days in CDCl<sub>3</sub> solution at rt. <sup>1</sup>H NMR (only signals clearly distinguishable from those corresponding to **4**):  $\delta$  8.45 (1 H, s, 1'-H); 8.02 (1 H, dd,  $J = 8.7, 1.8$  Hz, 3'-H); 7.87 (1 H, d,  $J = 8.9$  Hz, 8'-H); 7.20 (1 H, dd,  $J = 8.9, 2.4$  Hz, 7'-H); 4.39 (1 H, t,  $J = 7.9$  Hz, 2-H); 2.70-2.18 (4 H, m, 3-H<sub>2</sub> + 5-H<sub>2</sub>). <sup>13</sup>C NMR:  $\delta$  214.08, 195.49, 160.24, 137.74, 132.18, 132.04, 131.70, 128.05, 127.36, 125.51, 119.97, 105.94, 57.29, 55.69, 39.34, 27.63, 21.44.

**2-[Methoxy(6-methoxy-2-naphthyl)methylene]cyclopentanone (6):** An ethereal 0.5 M solution of CH<sub>2</sub>N<sub>2</sub><sup>9</sup> (5 mL) was slowly added to a solution of **4** (0.49 g, 1.8 mmol) in THF, and the reaction was left for 2 h at rt. The solvent was evaporated *in vacuo*, and the residue was identified as the dimethoxy compound **6** (0.51 g, 99%), white solid, mp 107-108° (EtOAc). IR: 1650 (C=O), 1625 (exocyclic or aromatic C=C) cm<sup>-1</sup>. <sup>1</sup>H NMR:  $\delta$  8.17 (1 H, s, 1'-H); 7.81 (1 H, d,  $J = 9.8$  Hz, 8'-H); 7.79 (1 H, dd,  $J = 8.6, 1.6$  Hz, 3'-H); 7.71 (1 H, d,  $J = 8.6$  Hz, 4'-H); 7.19 (1 H, dd,  $J = 9.8, 2.3$  Hz, 7'-H); 7.14 (1 H, d,  $J = 2.3$  Hz, 5'-H); 3.94 (3 H, s, 6'-OCH<sub>3</sub>), 3.60 (3 H, s,  $\beta$ -OCH<sub>3</sub>); 2.81-2.72 (4 H, m, 3-H<sub>2</sub> + 5-H<sub>2</sub>); 1.97 (2 H, quintet,  $J = 7.5$  Hz, 4-H<sub>2</sub>). <sup>13</sup>C NMR:  $\delta$  193.72, 166.96, 159.24, 136.72, 135.59, 131.08, 130.00, 128.14, 126.41, 126.29, 119.32, 114.04, 105.91, 57.78, 55.63, 32.07, 31.53, 19.84.

*Anal.* Calcd for C<sub>18</sub>H<sub>18</sub>O<sub>3</sub> (282.3): C, 76.57; H, 6.43. Found: C, 76.85; H, 6.65

**6-(6-Methoxy-2-naphthyl)-6-oxohexanoic acid (7):** A solution of **3** (1.00 g, 3.33 mmol) in MeOH (50 mL) was treated with 5 N NaOH (5 mL) and the mixture was refluxed for 7 h. The solvent was evaporated at reduced pressure, the solid residue was dissolved in water, and the resulting solution was extracted once with Et<sub>2</sub>O. The aqueous layer was acidified with 5 N HCl, then extracted with Et<sub>2</sub>O. The ethereal extract was dried (Na<sub>2</sub>SO<sub>4</sub>) and concentrated to afford **7** (0.82 g, 86%) as a white solid, mp 134-135° (AcOH). <sup>1</sup>H NMR:  $\delta$  12.43 (1 H, br s, CO<sub>2</sub>H); 8.40 (1 H, s, 1'-H); 8.00 (1 H, dd,  $J = 8.7, 1.7$  Hz, 3'-H); 7.86 (1 H, d,  $J = 9.0$  Hz, 8'-H); 7.77 (1 H, d,  $J = 8.7$  Hz, 4'-H); 7.20 (1 H, dd,  $J = 9.0, 2.4$  Hz, 7'-H); 7.15 (1 H, d,  $J = 2.4$  Hz, 5'-H); 3.95 (3 H, s, 6'-OCH<sub>3</sub>); 3.12 (2 H, t,  $J = 6.9$  Hz, 5-H<sub>2</sub>); 2.45 (2 H, t,  $J = 7.0$  Hz, 2-H<sub>2</sub>); 1.91-1.74 (4 H, m, 3-H<sub>2</sub> + 4-H<sub>2</sub>). <sup>13</sup>C NMR:  $\delta$  199.90, 178.83, 160.14, 137.67, 132.75, 131.53, 129.92, 128.26, 127.53, 125.01, 120.11, 106.16, 55.82, 38.36, 34.09, 24.80, 24.19.

*Anal.* Calcd for C<sub>17</sub>H<sub>18</sub>O<sub>4</sub> (286.3): C, 71.31; H, 6.34. Found: C, 71.58; H, 6.39

**Acknowledgments.**— The authors thank the *Xunta de Galicia* (XUGA 20309B98) for financially supporting this work.

## REFERENCES

1. W. V. Murray, M. P. Wachter, A. M. Kasper, D. C. Argentieri, R. J. Capetola and D. M. Ritchie, *Eur. J. Med. Chem.*, **26**, 159 (1991).
2. J. Catalán, P. Pérez, J. Laynez and F. García Blanco, *J. Fluoresc.*, **1**, 215 (1991).
3. M. I. Sáñez, A. Suárez, M. A. Ríos, M. C. Balo, F. Fernández and C. López, *Photochem. and Photobiol.*, **64**, 486 (1996).
4. C. Balo, F. Fernández, E. Lens and C. López, *Org. Prep. Proced. Int.*, **29**, 201 (1997).
5. a) J. Vene, *Bull. Soc. Chim. Fr.*, D 163 (1950). b) J. Cason, *J. Am. Chem. Soc.*, **68**, 2078 (1946).
6. O. Caamaño, A. Eirín, F. Fernández and E. Uriarte, *An. Quím.*, **83 C**, 81 (1987).
7. C. Balo, Ph. D. Dissertation, University of Santiago de Compostela, Spain (1995).
8. H. Gilman and J. F. Nelson, *Rec. Trav. Chim. Pays-Bas*, **55**, 518 (1936).
9. T. H. Black, *Aldrichimica Acta*, **16**, 3 (1983).

\*\*\*\*\*

# REDUCTIVE COUPLING OF KETONES WITH IMINES PROMOTED BY THE TITANIUM TETRACHLORIDE/SAMARIUM SYSTEM

Submitted by Yongmin Ma and Yongmin Zhang\*  
(04/18/00)

Department of Chemistry, Zhejiang University Xixi Campus  
Hangzhou, 310028, P.R. CHINA  
and  
Laboratory of Organometallic Chemistry  
Shanghai Institute of Organic Chemistry  
Chinese Academy of Science, Shanghai, 200032, P. R. CHINA

In the early 1970's, three groups discovered that low-valent titanium abstracted oxygen from aldehydes and ketones leading to the formation of olefins.<sup>1</sup> Since then, a large number of coupling reactions induced by low-valent titanium reagent have been reported.<sup>2</sup> Recently, our group reported the intermolecular and intramolecular reductive coupling of ketone-nitrile promoted by TiCl<sub>4</sub>/Sm system.<sup>3</sup> We now describe our preliminary results on the reductive couplings of aromatic aldimines