
This article was downloaded by: [University of Sydney] On: 03 September 2014, At: 12:01 Publisher: Taylor & Francis Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Phosphorus, Sulfur, and Silicon and the Related Elements

Publication details, including instructions for authors and subscription information: <u>http://www.tandfonline.com/loi/gpss20</u>

Synthesis of Organophosphorus Compounds in Terms of Elemental Phosphorus, Sulfur and their Derivatives

Elvira S. Batyeva , Il'yas S. Nizamov , Lidiya I. Kursheva & Liliya V. Frolova

^a A. E. Arbuzov Institute of Organic and Physical Chemistry, Arbuzov str. 8, Kazan 420088, RUSSIA

^b A. E. Arbuzov Institute of Organic and Physical Chemistry, Arbuzov str. 8, Kazan 420088, RUSSIA

^c A. E. Arbuzov Institute of Organic and Physical Chemistry , Arbuzov str. 8, Kazan 420088, RUSSIA

^d A. E. Arbuzov Institute of Organic and Physical Chemistry, Arbuzov str. 8, Kazan 420088, RUSSIA Published online: 17 Mar 2008. To cite this article: Elvira S. Batyeva , Il'yas S. Nizamov , Lidiya I. Kursheva & Liliya V. Frolova (1999) Synthesis of Organophosphorus Compounds in Terms of Elemental Phosphorus, Sulfur and their Derivatives, Phosphorus, Sulfur, and Silicon and the Related Elements, 144:1, 77-80, DOI: <u>10.1080/10426509908546186</u>

To link to this article: http://dx.doi.org/10.1080/10426509908546186

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at http://www.tandfonline.com/page/terms-and-conditions

Synthesis of Organophosphorus Compounds in Terms of Elemental Phosphorus, Sulfur and their Derivatives

ELVIRA S. BATYEVA, IL'YAS S. NIZAMOV, LIDIYA I. KURSHEVA and LILIYA V. FROLOVA

A. E. Arbuzov Institute of Organic and Physical Chemistry, Arbuzov str. 8, Kazan 420088, RUSSIA

We have developed new facile methods for synthesizing various organophosphorus compouds on the basis of elemental phosphorus and sulfur, phosphorus sulfides and esters of thiophosphorous acids.

Keywords: elemental phosphorus; sulfur; organophosphorus compouds

INTRODUCTION

One of the timely problems of phosphorus chemistry is the development of ecologically pure methods for their preparation. The traditional techniques for their synthesis are based on phosphorus chlorides. We have developed new facile methods for synthesizing various organophosphorus compouds on the basis of elemental phosphorus and sulfur, phosphorus sulfides and esters of thiophosphorous acids.

RESULTS AND DISCUSSION

Most of the reactions of elemental phosphorus (white or red) with organic substances studied have been carried out under basic and phase transfer catalyst conditions^[1-2]. Proton donating substances such as alcohols were used in the reactions of a mixture of elemental phosphorus and sulfur. However, these reactions are accompanied by the evolution of hydrogen sulfide and result in the mixture of products which hamper their isolation and purification^[3]. In this connection we investigated the reaction of elemental phosphorus and sulfur with non-containing proton reagents such as dialkyl disulfides, thioacetals, disilylated diols and stannylated alcohols^[4, 5].

We have observed that the reaction of red phosphorus and elemental sulfur with dialkyl disulfides leads to trialkyl pentathiophosphates with the S-S bond at $130-140^{\circ}C^{[4]}$. These pentathiophosphates decompose during distillation to form trialkyl tetrathiophosphates with elimination of sulfur.

$$4 P_{red} + 10/8 S_8 + 6 RS-SR \xrightarrow{130.140 °C} 4 (RS)_2 P-S-SR \xrightarrow{S}$$

-1/4 S_8
-1/4 S_8
-1/4 S_8
-1/4 S_8

The formation of tetrathiophosphates and trithiophosphites occurs already at room temperature in the presence of elemental iodine in dry benzene. Similar results were obtained when white phosphorus was used. We have also found that the interaction of red phosphorus with thioacetals in the presence of elemental sulfur and iodine results in products of a (1-alkylthio)alkyl tetrathiophosphate structure.


Cyclic silyldithiophosphates were synthesized by treating disilylated diols with a mixture of red phosphorus and elemental sulfur.

$$4 P_{red} + 10/8 S_8 + 4 He - 0-SIMe_3 - 160 °C + Me - 0-SIMe_3 + 2 (Me_3SI)_2S$$


Similarly stannyldithiophosphates were formed in the reaction of mixture of red phosphorus and sulfur with stannylated alcohols.

We consider that the reactions of elemental phosphorus and sulfur with organic substances may proceed through the intermediate formation of phosphorus sulfides. We decided to confirm this assumption and studied the reactions of phosphorus sulfides with dialkyl disulfides and thioacetals. These reactions resulted in the same final products that were found in the case of the reactions of mixtures of elemental phosphorus and sulfur as well. In the case of the reactions of dialkyl disulfides we observed the formation of the same intermediate pentathiophosphates with the S-S bond. Use of nucleophilic reagents such as potassium carbonate, sodium sulfide and sodium alkylthiolates results in the reactivity enhancement of phosphorus sulfides with dialkyl disulfides. Iodine leads to an increased reaction rate and yields of products formed in both types of reactions.

It was known that the reaction of the mixture of red phosphorus and sulfur with anisole results in the formation of Lawesson's reagent ^[3]. Lawesson's reagent and its analogues have proved to be effective initial substances for the synthesis of new types of cyclic organothiophosphorus compounds through their silyl derivatives. For this purpose we have obtained a series of disilyl esters of dithiophosphonic and trithiophosphonic acids on the basis of the reactions of 1,3,2,4dithiadiphosphetane-2,4-disulfides with disilylated sulfides and diols . We have also shown the prospects of these disilyldithiophosphonates and trithiophosphonates in the reactions with dimethylsilane and diphenylstannane dichlorides which resulted in new eight membered silicon and germanium-containing heterocycles .



We have managed to synthesize a new type of seven membered phosphocyclanes on the basis of disilyldithiophosphonates in their reactions with diphenylgermanium and tin dichlorides.



When the reaction of disulfides and white phosphorus and later on with thiols involving the formation of trialkyltrithiophosphite in good yields was discovered it has become possible to use the latter as initial material for synthesizing various organophosphorus compounds^[1]. We used the reaction based on the trialkylphosphites for obtaining organophosphorus compounds without any sulfur atoms.

$$(RS)_{3}P + 3 R'OH \xrightarrow{MX} (R'O)_{2}PH + 2 RSH + RSR'$$
$$M = Cu, Zn, Cd, Hq; X = Cl, Br, I$$

We have studied the reactions of triethyl and triphenyl trithiophosphites with excess alcohols (EtOH, MeOH) in the presence of metal halides.

Thus, novel methods for synthesizing organophosphorus compounds were developed in terms of elemental phosphorus and sufur and their derivatives.

References

- [1] C. Brown, R. F. Hudson, G. A. Wartew, Phosphorus and Sulfur, 5, 67 (1978).
- [2] B. A. Trofimov, T. N. Rakhmatulina, N. K. Gusarova, S. F. Malysheva, Uspekhi Khimii (Russian Chemical Review) (Russ.). 60, 2619 (1991).
- [3] F. N. Mazitova, V. K. Khairullin. Dokl. Akad. Nauk SSSR (Russ.), 250, 871 (1980); Zh. Obshch. Khim. (Russ.), 51, 958 (1981).
- [4] I. S. Nizamov, L. A. Al'metkina, G. G. Garifzyanova, E. S. Batyeva, V. A. Al'fonsov, Izv. Akad. Nauk. Ser. Khim. (Russ.), 1310 (1993).
- [5] E. S. Batyeva, Russ. Chem. Bull., 42, 1627 (1993).