

Copper-Catalyzed “Click” Reaction/ Direct Arylation Sequence: Modular Syntheses of 1,2,3-Triazoles


Lutz Ackermann,* Harish K. Potukuchi, Dirk Landsberg, and Rubén Vicente

Institut für Organische and Biomolekulare Chemie, Georg-August-Universität,
Tammannstrasse 2, 37077 Göttingen, Germany

lutz.ackermann@chemie.uni-goettingen.de

Received May 9, 2008

ABSTRACT



Inexpensive copper catalysts enabled modular one-pot multicomponent syntheses of fully decorated triazoles through a sustainable “click” reaction/direct arylation sequence.

The remarkable efficiency and selectivity accomplished with copper catalysts^{1,2} in Huisgen cycloadditions of organic azides with alkynes³ resulted in its application in various research areas, ranging from bioorganic and medicinal chemistry to material sciences.⁴ This “click” reaction proceeded highly regioselectively when using terminal alkynes, thereby providing 1,4-disubstituted 1,2,3-triazoles **2** as the sole products. On the contrary, [3 + 2] cycloadditions of the corresponding internal alkynes for the synthesis of fully decorated 1,2,3-triazoles **4** were found either not generally applicable or gave rise to mixtures of regioisomers.⁵

Catalytic C–H-bond functionalizations are attractive because of their ecologically and economically benign nature. Among these, regioselective direct arylations of (hetero)arenes represent a valuable alternative to traditional cross-couplings with stoichiometric amounts of organometallic reagents.⁶ The overall efficiency of this strategy was significantly improved by its combination with mechanistically distinct processes within sustainable one-pot reactions. Unfortunately, this approach proved thus far restricted to the use of catalysts based on relatively expensive transition metals, such as palladium⁷ or ruthenium.⁸

As part of our program on the development of sustainable C–H-bond functionalizations,⁹ we wished to explore the

(1) Tornøe, C. W.; Christensen, C.; Meldal, M. *J. Org. Chem.* **2002**, *67*, 3057–3064.

(2) Rostovtsev, V. V.; Green, L. G.; Fokin, V. V.; Sharpless, K. B. *Angew. Chem., Int. Ed.* **2002**, *41*, 2596–2599.

(3) Huisgen, R. *Angew. Chem.* **1963**, *75*, 604–637.

(4) Selected recent reviews: (a) Nandivada, H.; Jiang, X.; Lahann, J. *Adv. Mater.* **2007**, *19*, 2197–2208. (b) Angell, Y. L.; Burgess, K. *Chem. Soc. Rev.* **2007**, *36*, 1674–1689. (c) Fournier, D.; Hoogenboom, R.; Schubert, U. S. *Chem. Soc. Rev.* **2007**, *36*, 1369–1380. (d) Moses, J. E.; Moorhouse, A. D. *Chem. Soc. Rev.* **2007**, *36*, 1249–1262. (e) Lutz, J.-F. *Angew. Chem., Int. Ed.* **2007**, *46*, 1018–1025. (f) Dondoni, A. *Chem. Asian J.* **2007**, *2*, 700–708.

(5) (a) Zhang, L.; Chen, X.; Xue, P.; Sun, H. H. Y.; Williams, I. D.; Sharpless, K. B.; Fokin, V. V.; Jia, G. *J. Am. Chem. Soc.* **2005**, *127*, 15998–15999. (b) Díez-González, S.; Correa, A.; Cavallo, L.; Nolan, S. P. *Chem. Eur. J.* **2006**, *12*, 7558–7564. (c) Díez-González, S.; Nolan, S. P. *Synlett* **2007**, 2158–2167. (d) Majireck, M. M.; Weinreb, S. M. *J. Org. Chem.* **2006**, *71*, 8680–8683. (e) See also: Li, L.; Zhang, G.; Zhu, A.; Zhang, L. *J. Org. Chem.* **2008**, *73*, 3630–3633.


(6) For recent reviews, see: (a) Li, B.-J.; Yang, S.-D.; Shi, Z.-J. *Synlett* **2008**, 949–957. (b) Ackermann, L. *Top. Organomet. Chem.* **2007**, *24*, 35–60. (c) Satoh, T.; Miura, M. *Top. Organomet. Chem.* **2007**, *24*, 61–84. (d) Alberico, D.; Scott, M. E.; Lautens, M. *Chem. Rev.* **2007**, *107*, 174–238. (e) Bergman, R. G. *Nature* **2007**, *446*, 391–393. (f) Campeau, L.-C.; Stuart, D. R.; Fagnou, K. *Aldrich. Acta* **2007**, *40*, 35–41. (g) Seregin, I. V.; Gevorgyan, V. *Chem. Soc. Rev.* **2007**, *36*, 1173–1193. (h) Daugulis, O.; Zaitsev, V. G.; Shabashov, D.; Pham, Q. N.; Lazareva, A. *Synlett* **2006**, 3382–3388. (i) Yu, J.-Q.; Giri, R.; Chen, X. *Org. Biomol. Chem.* **2006**, *4*, 4041–4047.

(7) Selected recent examples for sequential palladium-catalyzed direct arylation-based processes: (a) Li, B.-J.; Tian, S.-L.; Fang, Z.; Shi, Z.-J. *Angew. Chem., Int. Ed.* **2008**, *47*, 1115–1117. (b) Ackermann, L.; Althammer, A. *Angew. Chem., Int. Ed.* **2007**, *46*, 1627–1629. (c) Dong, C.-G.; Hu, Q.-S. *Angew. Chem., Int. Ed.* **2006**, *45*, 2289–2292.

(8) Ackermann, L.; Born, R.; Álvarez-Becedo, P. *Angew. Chem., Int. Ed.* **2007**, *46*, 6364–6367.

application of less expensive copper complexes¹⁰ to direct arylation-based¹¹ sequential catalyzes. During these studies, we observed that a single copper precursor sequentially enabled two sustainable catalytic transformations, namely atom-economical “click” reactions and novel C–H bond functionalizations of 1,2,3-triazoles.^{12,13} Herein, we report on this approach, which allowed for modular syntheses of fully substituted 1,2,3-triazoles **4** through a chemo- and regioselective one-pot, four-component coupling (Scheme 1).

Scheme 1. Copper-Catalyzed Sequential Synthesis of Triazoles



To probe the viability of the envisioned sequential catalytic protocol, we first tested copper-catalyzed direct arylations of 1,4-disubstituted 1,2,3-triazoles **2** (Table 1). Triazoles **2** displaying electron-donating alkyl-substituents on the heteroarenes were efficiently converted with a variety of substituted aryl iodides **3** (entries 1–9). Pronucleophiles with aryl groups at either position of the heterocycle were regioselectively functionalized at the electron-rich triazole moieties (entries 4–8). Notably, the copper-catalyzed direct arylation of a monosubstituted 1,2,3-triazole proceeded also exclusively at position C-5, yielding triazole **4i** exclusively (entry 9).

With an effective protocol for copper-catalyzed direct arylations of triazoles **2** in hand, we tested its application to the one-pot sequential synthesis of fully¹⁴ substituted triazoles **4**. At the outset, we chose preformed alkyl-substituted azides as starting materials for the initial catalytic “click” reaction (Table 2).

(9) Ackermann, L. *Synlett* **2007**, 507–526.

(10) For selected economical copper-catalyzed C–H bond functionalizations, see: (a) Tsang, W. C. P.; Zheng, N.; Buchwald, S. L. *J. Am. Chem. Soc.* **2005**, *127*, 14560–14561. (b) Chen, X.; Hao, X.-S.; Goodhue, C. E.; Yu, J.-Q. *J. Am. Chem. Soc.* **2006**, *128*, 6790–6791. (c) Li, Z.; Bohle, D. S.; Li, C.-J. *Proc. Natl. Acad. Sci. U.S.A.* **2006**, *103*, 8928–8933. (d) Brasche, G.; Buchwald, S. L. *Angew. Chem., Int. Ed.* **2008**, *47*, 1932–1934. (e) For the use of stoichiometric amounts of copper, see: Uemura, T.; Imoto, S.; Chatani, N. *Chem. Lett.* **2006**, *35*, 842–843.

(11) For the elegant development of copper-catalyzed direct arylations, see: (a) Do, H.-Q.; Daugulis, O. *J. Am. Chem. Soc.* **2007**, *129*, 12404–12405. (b) Do, H.-Q.; Daugulis, O. *J. Am. Chem. Soc.* **2008**, *130*, 1128–1129. (c) For the use of stoichiometric amounts of copper in direct arylations, see: Yoshizumi, T.; Tsurugi, H.; Satoh, T.; Miura, M. *Tetrahedron Lett.* **2008**, *49*, 1598–1600.

(12) For palladium-catalyzed direct arylations of 1,2,3-triazoles **2**, see: (a) Chuprakov, S.; Chernyak, N.; Dudnik, A. S.; Gevorgyan, V. *Org. Lett.* **2007**, *9*, 2333–2336. (b) Iwasaki, M.; Yorimitsu, H.; Oshima, K. *Chem. Asian J.* **2007**, *2*, 1430–1435. (c) Ackermann, L.; Vicente, R.; Born, R. *Adv. Synth. Catal.* **2008**, *350*, 741–748.

(13) Ruthenium-catalyzed direct arylations of 1,2,3-triazoles **2**: Ackermann, L.; Vicente, R.; Althammer, A. *Org. Lett.* **2008**, *10*, 2299–2305.

(14) One-pot syntheses of 1,4-disubstituted 1,2,3-triazoles **2** from in situ generated azides were reported: (a) Feldman, A. K.; Colasson, B.; Fokin, V. V. *Org. Lett.* **2004**, *6*, 3897–3899. (b) Kacprzak, K. *Synlett* **2005**, 943–946. (c) Andersen, J.; Bolvig, S.; Liang, X. *Synlett* **2005**, 2941–2947.

Table 1. Copper-Catalyzed Direct Arylations of Triazoles **2**^a










entry	R ¹	R ²	4		yield
1	Oct	Hex		4a	74%
2	Oct	Hex		4b	72%
3	Bn	Hex		4c	86%
4	Oct	Ph		4d	98%
5	Oct	Ph		4e	73%
6	Ph	Bu		4f	71%
7	3-MeC ₆ H ₄	Bu		4g	92%
8	2-MeC ₆ H ₄	Bu		4h	92%
9	Bn	H		4i	93%

^a Reaction conditions: **2** (1.00 mmol), **3** (3.00 mmol), CuI (10 mol %), LiOt-Bu (2.00 mmol), DMF (3.0 mL), 140 °C, 24 h; yields of isolated product.

Importantly, these studies revealed that a single inexpensive copper catalyst could be employed for two mechanistically distinct reactions, providing triazoles **4** in high yields. This protocol proved broadly applicable, allowing for the use of alkyl- (entry 1) as well as aryl-substituted alkynes (entries 2–9). Further, electrophiles with either electron-donating (entries 1–5) or electron-withdrawing (entries 6–9) substituents were successfully converted, even when more sterically demanding (entries 1–4, and 9).

Subsequently, we probed a modular synthesis of *N*-aryltriazoles **4** through a chemoselective coupling of four

Table 2. Sequential Copper-Catalyzed Synthesis of Triazoles **4**^a







$\text{H}-\text{C}\equiv\text{C}-\text{R} \xrightarrow[\text{b) ArI (3), LiOt-Bu, DMF, 140 }^\circ\text{C}]{\text{a) AlkN}_3, \text{CuI (10 mol \%), DMF, 60 }^\circ\text{C}} \text{Alk}-\text{N}_2-\text{C}(\text{Ar})=\text{N}-\text{R}$				
entry	R	Alk	4	yield
1	Bu	Bn		4j 67%
2	Ph	Oct		4d 69%
3	Ph	Bn		4k 70%
4 ^b	Ph	Bn		4l 84%
5	Ph	Bn		4m 61%
6 ^b	Ph	Bn		4n 69%
7 ^b	Ph	Bn		4o 80%
8 ^b	Ph	Bn		4p 79%
9 ^b	Ph	Oct		4q 63%

^a Reaction conditions: (a) **1** (1.00 mmol), AlkN₃ (1.00 mmol), CuI (10 mol %), DMF (3 mL), 60 °C, 4 h; (b) **3** (3.00 mmol), LiO-*t*-Bu (2.00 mmol), DMF (2.0 mL), 140 °C, 20 h; yields of isolated product. ^b **3** (3.00 mmol), CuI (10–30 mol %), LiO-*t*-Bu (2.00 mmol), DMF (2.0 mL), 140 °C, 20 h.

components. A copper catalyst modified with *N,N'*-dimethylethylenediamine (DMEDA) as stabilizing ligand yielded disubstituted triazoles **2** efficiently. Importantly, their regioselective direct arylations were also achieved with the same copper catalyst (Table 3).¹⁵ Therefore, a variety of triazoles **4** were obtained, displaying both electron-rich (entries 1–4)

(15) **Representative Procedure (Table 3, Entry 2).** A suspension of CuI (19 mg, 0.10 mmol, 10 mol %) and NaN₃ (69 mg, 1.05 mmol) in DMF (3 mL) was treated with 1-hexyne (82 mg, 1.00 mmol), 3-iodobenzene (204 mg, 1.00 mmol), and *N,N'*-dimethylethylenediamine (13 mg, 0.15 mmol, 15 mol %) and was stirred under N₂ at ambient temperature for 2 h. Then, LiO-*t*-Bu (160 mg, 2.00 mmol), 2-iodoanisole (702 mg, 3.00 mmol), and DMF (2.0 mL) were added, and the resulting suspension was stirred under N₂ at 140 °C for 20 h. At ambient temperature, Et₂O (50 mL) and H₂O (50

Table 3. Sequential Copper-Catalyzed Four-Component Synthesis^a

$\text{Bu}-\text{C}\equiv\text{C}-\text{H} + \text{NaN}_3 + \text{I}-\text{C}_6\text{H}_4-\text{R}^1 \xrightarrow[\text{b) I}-\text{C}_6\text{H}_4-\text{R}^2, \text{LiO}-t\text{-Bu, 140 }^\circ\text{C, 20 h}]{\text{a) CuI (10 mol \%), DMEDA (15 mol \%), DMF, 22 }^\circ\text{C, 2 h}} \text{Bu}-\text{N}_2-\text{C}(\text{R}^1)=\text{N}-\text{C}(\text{R}^2)-\text{C}_6\text{H}_4-\text{R}^2$				
entry	R ¹	R ²	4	yield
1	3-Me	4-MeO		4r 70%
2	3-Me	2-MeO		4g 81%
3	H	4-MeO		4f 73%
4	H	2-MeO		4s 74%
5	4-F	2-MeO		4t 70%
6	4-Cl	2-MeO		4u 67%

^a Reaction conditions: (a) **1** (1.00 mmol), **3** (1.00 mmol), NaN₃ (1.05 mmol), CuI (10 mol %), DMEDA (15 mol %), DMF (3.0 mL), 22 °C, 2 h; (b) **3** (3.00 mmol), LiO-*t*-Bu (2.00 mmol), DMF (2.0 mL), 140 °C, 20 h; yields of isolated product. DMEDA = *N,N'*-dimethylethylenediamine.

as well as functionalized electron-deficient (entries 5 and 6) aryl-substituents. It is noteworthy that this four-component one-pot synthesis involved the formation of one C–C- and three C–N-bonds in a highly selective fashion.

In conclusion, a novel copper-catalyzed direct arylation of 1,4-disubstituted triazoles **2** set the stage for the development of a modular one-pot approach to diversely substituted 1,2,3-triazoles **4**. The unprecedented direct arylation-based

mL) were added, and the separated aqueous layer was extracted with Et₂O (2 × 75 mL). The combined organic layers were washed with satd aq NH₄Cl (50 mL), H₂O (50 mL), and brine (50 mL), dried over Na₂SO₄, and concentrated in vacuum. The remaining residue was purified by column chromatography on silica gel (*n*-hexane/EtOAc 12/1 → 10/1) to yield **4g** as an off-white solid (260 mg, 81%).

sequential copper-catalysis combined atom-economical “click” reactions with sustainable C–H-bond functionalizations, enabling the chemoselective coupling of up to four components through the formation of one C–C- and three C–N-bonds.

Acknowledgment. Support by the DFG, the DAAD (fellowship to H.K.), the Alexander-von-Humboldt founda-

tion (fellowship to R.V.), and the Fonds der Chemischen Industrie is gratefully acknowledged.

Supporting Information Available: Experimental procedures, characterization data, and ^1H NMR and ^{13}C NMR spectra for all compounds. This material is available free of charge via the Internet at <http://pubs.acs.org>.

OL801078R