


This article was downloaded by: [University of Borås]

On: 07 October 2014, At: 12:12

Publisher: Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Organic Preparations and Procedures International: The New Journal for Organic Synthesis

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/uopp20>

STEREOSELECTIVE SYNTHESIS OF (E)-VINYL ARYL SULFIDES via HYDROZIRCONATION OF TERMINAL ALKYNES

Xian Huang^a, Ping Zhong^a & Wei-Rong Guo^a

^a Department of Chemistry, Xi-Xi Campus Zhejiang University, Hangzhou, 310028, P. R. CHINA

Published online: 11 Feb 2009.

To cite this article: Xian Huang, Ping Zhong & Wei-Rong Guo (1999) STEREOSELECTIVE SYNTHESIS OF (E)-VINYL ARYL SULFIDES via HYDROZIRCONATION OF TERMINAL ALKYNES, Organic Preparations and Procedures International: The New Journal for Organic Synthesis, 31:2, 201-203, DOI:

[10.1080/00304949909355711](https://doi.org/10.1080/00304949909355711)

To link to this article: <http://dx.doi.org/10.1080/00304949909355711>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms &

OPPI BRIEFS

STEREOSELECTIVE SYNTHESIS OF (E)-VINYL ARYL SULFIDES

via HYDROZIRCONATION OF TERMINAL ALKYNES


Submitted by
(07/24/98)

Xian Huang*, Ping Zhong and Wei-rong Guo

Department of Chemistry, Xi-Xi Campus
Zhejiang University, Hangzhou, 310028, P. R. CHINA

Vinyl sulfides have attracted special attention as important intermediates in various synthetic transformations,^{1,2} and their stereospecific synthesis has been attempted.^{3,4} However, no convenient synthesis of (E)-vinyl sulfides from aryl or alkyl substituted alkynes has been reported. For example, the addition of thiophenol to monosubstituted ethynes is very slow at room temperature and the products exhibit predominantly the Z configuration, or a mixture of the Z and E configurations.⁵⁻⁷ It was reported that the palladium-catalyzed cross-coupling of 9-RS-9-BBN with (E)-1-iodo-1-alkenes afforded (E)-vinyl sulfides⁸, but the synthesis of 9-RS-9-BBN and (E)-1-iodo-1-alkenes are not easy.

Vinylzirconium complexes reacted with phenyltellurenyl iodide or arylselenenyl bromides readily to afford (E)-vinyl tellurides⁹ or (E)-vinyl selenides.¹⁰ Considering the high electrophilicity of arylsulfenyl chloride, we decided to react them with the vinylzirconium complexes **2** produced by hydrozirconation of terminal alkyne **1**. The experimental results showed that the reaction proceeded easily to afford (E)-vinyl sulfides **3** whose configuration was confirmed by the coupling constants (*J* = 15-16 Hz) of the vinyl protons and the absorption bands at 945-960 cm⁻¹ in IR spectrum.



The present method has advantages of high stereoselectivity, readily available starting materials, simple procedures and mild reaction conditions as well as high yields.

EXPERIMENTAL SECTION

¹H NMR spectra were recorded on PMX-60 spectrometer in CDCl₃ as the solvent with TMS as an internal standard. *J* values are given in Hz. IR spectra were determined as films on PE-683 spectrophotometer. Silica gel 60 GF254 was used for analytical and preparative TLC. All reactions were

carried out in pre-dried glassware (140°, 4h) and cooled under a stream of dry nitrogen. All solvents were dried, deoxygenated and redistilled before use. The arylsulenyl chlorides were prepared from aryl thiols and sulfuryl chloride.¹¹

TABLE. Synthesis of (E)-Vinyl Sulfides **3a-g**

Product	R	Ar	Yield(%)
3a	Ph	Ph	85
3b	Ph	4-MeC ₆ H ₄	80
3c	Ph	4-ClC ₆ H ₄	90
3d	<i>n</i> -Bu	Ph	86
3e	<i>n</i> -Bu	4-MeC ₆ H ₄	74
3f	AcOCH ₂	Ph	75
3g	CH ₃ OCH ₂	Ph	88

General Procedure for the Synthesis of (E)-Vinyl Sulfides 3a-g. A mixture of Cp₂Zr(H)Cl¹² (1mmol) and terminal alkyne **1** (1mmol) in THF (5mL) was stirred at room temperature for 20 min. The resulting solution was cooled to 0° and into it was injected ArSCl¹¹ (1mmol). Then the mixture was stirred at room temperature for 30 min. The solvent was removed on a rotary evaporator under reduced pressure. The residue was extracted with light petroleum (3 x 6 mL) and filtered through a short plug of silica gel. After evaporation of the filtrate, the residue was purified by preparative TLC on silica gel eluting with light petroleum.

Compound 3a: Colorless oil, lit.¹³, bp. 150-151°/1.5mmHg, IR (film): 3040, 1612, 1594, 1470, 950, 688 cm⁻¹; ¹H NMR: δ 6.90-7.40 (m, 10H), 6.9 (d, 1H, J = 15), 6.65 (d, 1H, J = 15).

Anal. Calcd for C₁₄H₁₂S: C, 79.20; H, 5.70. Found: C, 78.98; H, 5.83

Compound 3b: Colorless oil, IR (film): 3070, 1610, 1580, 1505, 948, 775, 689 cm⁻¹; ¹H NMR: δ 7.05-7.40 (m, 9H), 6.80 (d, 1H, J = 15), 6.60 (d, 1H, J = 15), 2.26 (s, 3H).

Anal. Calcd for C₁₅H₁₄S: C, 79.60; H, 6.23. Found: C, 79.48; H, 6.35

Compound 3c: Colorless oil, lit.¹³, bp. 164-165°/1.0mmHg, IR (film): 3050, 1608, 1580, 1480, 945, 810 cm⁻¹; ¹H NMR: δ 7.15-7.55 (m, 9H), 6.85 (d, 1H, J = 15), 6.56 (m, 1H).

Anal. Calcd for C₁₄H₁₁ClS: C, 68.15; H, 4.49. Found: C, 67.96; H, 4.56

Compound 3d: Colorless oil, IR (film): 3040, 1595, 1450, 955, 785 cm⁻¹; ¹H NMR: δ 7.00-7.50 (m, 5H), 6.05 (d, 1H, J = 15), 5.70 (m, 1H), 2.10 (m, 2H), 0.88-1.60 (m, 7H).

Anal. Calcd for C₁₂H₁₆S: C, 74.94; H, 8.39. Found: C, 74.76; H, 8.51

Compound 3e: Colorless oil, IR (film): 3040, 1500, 950, 785 cm⁻¹; ¹H NMR: δ 6.90-7.40 (m, 4H), 6.05 (d, 1H, J = 15), 5.80 (m, 1H), 2.50 (s, 3H), 2.10 (m, 2H), 0.70-1.60 (m, 7H).

Anal. Calcd for C₁₃H₁₈S: C, 75.67; H, 8.79. Found: C, 75.51; H, 8.93

Compound 3f: Colorless oil, IR (film): 3070, 1740, 1600, 1495, 955 cm⁻¹; ¹H NMR: δ 7.10-7.50 (m, 5H), 6.30 (d, 1H, J = 15), 5.70 (m, 1H), 4.45 (d, 2H, J = 6), 2.00 (s, 3H).

Anal. Calcd for C₁₁H₁₂O₂S: C, 63.43; H, 5.81. Found: C, 63.28; H, 5.97

Compound 3g: Colorless oil, IR (film): 3080, 1595, 1488, 950 cm^{-1} ; ^1H NMR: δ 7.00-7.50 (m, 5H), 6.30 (d, 1H, $J = 15$), 5.70 (m, 1H), 3.82 (d, 2H, $J = 6$), 3.20 (s, 3H).

Anal. Calcd for $\text{C}_{10}\text{H}_{12}\text{OS}$: C, 66.63; H, 6.71. Found: C, 66.40; H, 6.82

The ^1H NMR spectra of **3a**,⁷ **3b**,⁷ **3c**,⁷ of **3d**,⁶ **3e**⁸ and of **3f**¹⁴ and **3g**¹⁴ were identical to those reported in the references cited.

Acknowledgement.- Project 29772007 was supported by the National Nature Science Foundation of China and this work was also supported by The Laboratory of Organometallic Chemistry, Chinese Academy of Science.

REFERENCES

1. T. Takeda, F. Kanamori, H. Matsusita and T. Fujiwara, *Tetrahedron Lett.*, **32**, 6563 (1991).
2. D. F. Andres, E. G. Laurent, B. S. Marquet, H. Benotmane and A. Bensadat, *Tetrahedron* **51**, 2605 (1995).
3. Y. Takikawa, K. Shimada, H. Matsumoto, H. Tanabe and S. Takizawa, *Chemistry Lett.*, 1351 (1983).
4. M. Hoshi, Y. Masuda and A. Arase, *Chem. Commun.*, 1068 (1985).
5. L. Benati, L. Capella, P. C. Montecocchi and P. Spagnolo, *J. Chem. Soc. Perkin Trans. 1*, 1035 (1995).
6. L. Benati, P. C. Montecocchi and P. Spagnolo, *ibid.*, 2103 (1991).
7. A. A. Oswald, K. Griesbaum and B. E. Hudson, Jr. and J. M. Bregman, *J. Am. Chem. Soc.*, **86**, 2877 (1964).
8. T. Ishiyama, M. Mori, A. Suzuki and N. Miyaara, *J. Organomet. Chem.*, **525**, 225 (1996).
9. J. W. Sung, C.-W. Lee and D. Y. Oh, *Tetrahedron Lett.*, **36**, 1503 (1995).
10. X. Huang and L.-S. Zhu, *J. Chem. Soc. Perkin Trans. 1*, 767 (1996).
11. D. N. Harpp, B. T. Friedlander and R. A. Smith, *Synthesis*, 181 (1979).
12. S. L. Buchwald, S. J. LaMaire, R. B. Nielsen, B. T. Watson and S. M. King, *Tetrahedron Lett.*, **28**, 3895 (1987).
13. R. F. Brookes, J. E. Cranham, D. Greenwood and H. A. Stevenson, *J. Sci. Food Agr.*, **8**, 561 (1957).
14. A. Behzadi and L. N. Owen, *J. Chem. Soc. Perkin Trans 1*, 25 (1974).
