An efficient method for the preparation of peptide alcohols†

Alan Roy Katritzky,*^a Nader Elmaghwry Abo-Dya,^{a,b} Srinivasa Rao Tala,^a Kapil Gyanda^a and Zakaria Kamel Abdel-Samii^b

Received 23rd March 2009, Accepted 13th July 2009 First published as an Advance Article on the web 26th August 2009 DOI: 10.1039/b905730g

N-Protected LL-dipeptide alcohols **3a–p**, diastereomeric mixture (**3d** + **3d**') and tripeptide alcohols **6a–c** were synthesized by treatment of various amino alcohols with *N*-protected(α -aminoacyl)benzotriazoles **1a–c**, **1f–m**, (**1a** + **1a**') and *N*-protected(α -dipeptidoyl)benzotriazoles **5a**, **5b** respectively in good yields with complete retention of chirality.

Introduction

Peptide alcohols are synthetic intermediates for peptide aldehydes which are inhibitors of m-calpain,^{1a} Subtilisin 72,^{1b} α -Chymotrypsin,^{1b} HIV-protease,^{1c,d} multicatalytic proteinase complex (MPC)^{1e} and other biologically and pharmaceutically active molecules.^{1f,g} Peptide alcohols serve as reversible activators of catalytic activity of the 20S proteasome.² Peptide alcohols are also used as ligands or precursors for ligands in asymmetric catalysis,^{3a-g} and in the preparation of a prototype calix[4]arenebased receptor for carbohydrate recognition.⁴

Reported preparations of peptide alcohols have utilized: (i) coupling reactions with DCC/HOBT,^{1c,3d} EDC/HOBT,^{3g} HATU, DIPEA,^{1a} NMM, BuⁱOCOCl,^{3b} DEPBT,⁵ (ii) aminoacyl chlorides,⁶ (iii) activated esters,^{3b,e} (iv) reduction of esters,^{4,7} ketones,⁶ (v) enzymatic synthesis,⁸ and (vi) solid phase synthesis.⁹

Utilization of these methods involves: (i) cumbersome reaction conditions,^{3b,3e,6} (ii) producing mixtures of diastereomeric alcohols on reduction,⁶ (iii) long reaction times (up to 53 hours),⁸ (iv) use of expensive reagents and (v) low yields.⁶ Thus a mild and efficient approach to peptide alcohols would be of utility.

N-Acylbenzotriazoles are advantageous for N-, O-, C-, S-acylation,^{10a-j} especially where the corresponding acid chlorides are unstable or difficult to prepare.^{10k,l} N-Fmoc-(α-aminoacyl)benzotriazoles and their Boc- and Cbz- analogs enabled the preparation of chiral di-, tri- and tetra-peptides in good yields from natural amino acids in the solution phase.^{10b,11a-b} Recently, we have also prepared tri-, tetra-, penta-, hexa-, and heptapeptides in 71% average crude yields by microwave-assisted solid phase peptide synthesis utilizing N-Fmoc-(α-aminoacyl)benzotriazoles and N-Fmoc-protected(α -dipeptidoyl)benzotriazoles.^{12a,b} We now report the preparation of di- and tri-peptide alcohols in solution phase by reaction of Nprotected(α-aminoacyl)benzotriazoles N-protected(α and dipeptidoyl)benzotriazoles with amino alcohols.

Results and discussion

1. Preparation of dipeptide alcohols 3a-p and diastereomeric mixture (3d + 3d')

The starting *N*-protected(α -aminoacyl)benzotriazoles **1a–e**, **1g–m** and (**1a + 1a'**) (*N*-Cbz-L-Ala-Bt **1a**, *N*-Cbz-DL-Ala-Bt (**1a + 1a'**), *N*-Fmoc-L-Val-Bt **1b**, *N*-Fmoc-L-Met-Bt **1c**, *N*-Fmoc-L-Phe-Bt **1d**, *N*-Fmoc-Gly-Bt **1e**, *N*-Cbz-L-Phe-Bt **1g**, *N*-Fmoc-L-Glu(O'Bu)-Bt **1h**, *N*-Fmoc-L-Asp(O'Bu)-Bt **1i**, *N*-Fmoc-L-Lys(Boc)-Bt **1j**, *N*-Fmoc-L-Trp-Bt **1k**, *N*-Fmoc-L-His(Trt)-Bt **1l** and *N*-Fmoc-L-Cys(Trt)-Bt **1m**) were prepared in 75–90% yield from corresponding *N*-protected amino acids following our previously published one-step procedure.¹⁰⁶ *N*-Boc-L-Val-Bt **1f** was prepared as described previously.^{10e}

Reaction of *N*-protected(α -aminoacyl)benzotriazoles **1a–c**, **1f–m** and racemic mixture (**1a** + **1a**') with various amino alcohols **2** in THF at room temperature for 6 h gave *N*-protected dipeptide alcohols **3a–p** and diastereomeric mixture (**3d** + **3d**') in 46–89% yield, all isolated without column chromatography (Scheme 1, Table 1). The dipeptide alcohols **3a–p** were characterized by ¹H-NMR, ¹³C-NMR, HRMS and elemental analysis. Diastereomeric mixtures were prepared to confirm that the original chirality of the amino acid used was maintained during the formation of peptide alcohol by means of chiral HPLC analysis. Chiral HPLC (detection at 254 nm, flow rate 1 mL/min, hexanes– isopropanol (9:1) as solvent using Whelk-O1 chiral column) showed a single peak for **3d** at 21.12 min. By contrast, two peaks were observed for the corresponding racemic mixtures (**3d** + **3d**') at 19.25 and 21.12 min, confirming the enantiopurity of **3d**.



Scheme 1 Preparation of dipeptide alcohols.

2. Preparation of *N*-Fmoc-α-dipeptides 4a, 4b from *N*-Fmoc-(α-aminoacyl)benzotriazoles 1d, 1e

N-Fmoc-protected dipeptides **4a** and **4b** were prepared by reaction of *N*-Fmoc-protected(α -aminoacyl)benzotriazoles **1d** and **1e**, respectively, with the corresponding amino acids in

^aCenter for Heterocyclic Compounds, Department of Chemistry, University of Florida, Gainesville, FL 32611-7200, USA. E-mail: katritzky@ chem.ufl.edu; Fax: +1-352-392-9199; Tel: +1-352-392- 0554

^bDepartment of Pharmaceutical Organic Chemistry, Faculty of Pharmacy, Zagazig University, Zagazig-44519, Egypt

[†] Electronic supplementary information (ESI) available: General experimental procedures with spectral data for compounds **3a–p**, **4a**, **4b**, **5a**, **5b** and **6a–c**. See DOI: 10.1039/b905730g

Table 1Preparation of dipeptide alcohols 3a-p and (3d + 3d')

Entry	1	Amino alcohol 2	Product 3 , yield $(\%)^a$
1	1f	L-Phenylalaninol	<i>N</i> -Boc-L-Val-L-Phenylalaninol 3a , 86
2	1f	L-Leucinol	N-Boc-L-Val-L-Leucinol 3b, 77
3	1f	L-t-Leucinol	N-Boc-L-Val-L-t-Leucinol 3c, 81
4	1a	L-Phenylalaninol	N-Cbz-L-Ala-L-Phenylalaninol 3d, 88
5	(1a + 1a')	L-Phenylalaninol	N-Cbz-DL-Ala-L-Phenylalaninol (3d + 3d'), 89
6	la	L-Leucinol	N-Cbz-L-Ala-L-Leucinol 3e ^b , 82
7	1c	L-Phenylalaninol	N-Fmoc-L-Met-L-Phenylalaninol 3f, 60
8	1b	L-Phenylglycinol	N-Fmoc-L-Val-L-Phenylglycinol 3g, 54
9	1a	L-Phenylglycinol	N-Cbz-L-Ala-L-Phenylglycinol 3h , 85
10	1g	L-Phenylglycinol	N-Cbz-L-Phe-L-Phenylglycinol 3i, 60
11	lg	L-Leucinol	N-Cbz-L-Phe-L-Luccinol 3i, 46
12	1ที่	L-Phenylalaninol	N-Fmoc-L-Glu(O'Bu)-L-Phenylalaninol 3k, 80
13	1i	L-Phenylglycinol	N-Fmoc-L-Asp(O'Bu)-L-Phenylglycinol 31, 77
14	1j	L-Leucinol	N-Fmoc-L-Lys(Boc)-L-Leucinol 3m, 85
15	1 k	L-Phenylalaninol	N-Fmoc-L-Trp-L-Phenylalaninol 3n , 83
16	11	L-t-Leucinol	N-Fmoc-L-His(Trt)-L-t-Leucinol 30, 60
17	1m	L-Leucinol	N-Fmoc-L-Cys(Trt)-L-Leucinol 3p, 70

Table 2 Preparation of N-Fmoc- α -dipeptides 4a, 4b from N-Fmoc-protected(α -aminoacyl)benzotriazoles 1d, 1e

Entry	1	Amino acid	Product 4, yield $(\%)^a$	4 , mp (°C)	
1 2	1d 1e	L-Ala-OH L-Leu-OH	<i>N</i> -Fmoc-L-Phe-L-Ala-OH 4a , 87 <i>N</i> -Fmoc-Gly-L-Leu-OH 4b , 85	208–210 ^b 133–134 ^c	
^a Isolated yield. ^b Lit., ^{12b} mp 208.7–210.6 °C. ^c Lit., ¹⁴ mp 133–135 °C.					

acetonitrile–water (3:1) in the presence of triethylamine at room temperature in 85-87% yield as described previously (Scheme 2, Table 2).¹⁰⁶



Scheme 2 Preparation of *N*-Fmoc- α -dipeptides 4a, 4b from *N*-Fmoc-protected(α -aminoacyl)benzotriazoles 1d, 1e.

3. Preparation of N-Fmoc-(a-dipeptidoyl)benzotriazoles 5a, 5b

N-Fmoc-(α -dipeptidoyl)benzotriazoles **5a** and **5b** were prepared by treatment of *N*-Fmoc- α -dipeptides **4a** and **4b**, respectively, with benzotriazole and thionyl chloride in CH₂Cl₂ at –15 °C for 3 h in 78–82% yield as described previously (Scheme 3, Table 3).^{12b} Compounds **5a**, **5b** were characterized by ¹*H*-NMR, ¹³*C*-NMR and elemental analysis. The *N*-Fmoc-(α -dipeptidoyl)benzotriazoles **5a**, **5b** were utilized for the synthesis of *N*-Fmoc-protected tripeptide alcohols **6a–c**.



Scheme 3 Preparation of N-Fmoc-(α -dipeptidoyl)benzotriazoles 5a, 5b from N-Fmoc- α -dipeptides 4a, 4b.

Table 3 Preparation of N-Fmoc-(α-dipeptidoyl)benzotriazoles 5a, 5b

Entry	4	Product 5	5, yield (%) ^{<i>a</i>}	5, mp (°C)
1 2	4a 4b	N-Fmoc-L-Phe-L-Ala-Bt, 5a N-Fmoc-Gly-L-Leu-Bt, 5b	5a , 82 5b , 78	155–156 165–167
" Isolat	ed yie	ld. ^{<i>b</i>} Lit., ¹² mp 155.2–156.9 °C.		

4. Preparation of tripeptide alcohols 6a-c

N-Fmoc-protected tripeptide alcohols **6a–c** were prepared in solution phase by treatment of *N*-Fmoc-(α -dipeptidoyl)benzotriazoles **5a** and **5b** with amino alcohols **2** in THF for 4 h at 0 °C followed by 3 h at room temperature in good yields, all isolated without column chromatography (Scheme 4, Table 4). These compounds were characterized by ¹*H*-NMR, ¹³*C*-NMR, and elemental analysis. No detectable racemization of the tripeptide alcohols was observed in chiral HPLC analysis.



Scheme 4 Preparation of tripeptide alcohols 6a-c from *N*-Fmoc- $(\alpha$ -dipeptidoyl)benzotriazoles 5a, 5b.

In comparison our methodology offers: (1) mild and simple reaction and work-up conditions: we isolated all products under milder reaction conditions and by simply washing or recrystallization; (2) high yields: most of our products were obtained in
 Table 4
 Preparation of tripeptide alcohols 6a-c from N-Fmoc-(α-dipeptidoyl)benzotriazoles 5a, 5b

Entry	5	Amino alcohol 2	Product 6 , yield (%) ^{<i>a</i>}
1	5a	L-Phenylalaninol	N-Fmoc-L-Phe-L-Ala-L-Phenylalaninol ^b 6a, 69
2	5a	L-Phenylglycinol	N-Fmoc-L-Phe-L-Ala-L-Phenylglycinol ^b 6b, 65
3	5b	L-Phenylalaninol	N-Fmoc-Gly-L-Leu-L-Phenylalaninol ^b 6c, 69
^{<i>a</i>} Isolated vield. ^{<i>b</i>} R	etention times for 6a . 28.7: 6	b , 13.65; and 6c , 26.67 min.	

high yield (46–89%); (3) no racemization occurred; (4) less time to complete; and (5) uses inexpensive reagents.

Conclusions

In conclusion, a convenient and efficient method for the preparation of di-, and tri-peptide alcohols under mild and simple reaction conditions has been developed by reacting *N*-protected(α -aminoacyl)benzotriazoles and *N*-protected(α -dipeptidoyl)benzotriazoles with amino alcohols at room temperature. All the peptide alcohols were obtained in moderate to good yields with no detectable racemization for chiral compounds.

Experimental

Starting materials and solvents were purchased from commercial sources and used without further purification. Melting points were determined on a Fisher melting apparatus and are uncorrected. Column chromatography was carried out using silica gel 200–425 mesh. HPLC analyses were performed on Shimadzu SPD-20-A using a Whelk-O1 chiral column with detection at 254 nm, a flow rate of 1.0 mL/min and hexanes–isopropanol (9:1) as the eluting solvent. ¹H NMR (300 MHz) and ¹³C NMR (75 MHz) spectra were recorded on a 300 MHz NMR spectrometer with CDCl₃ or DMSO-d₆ as solvents. J values are given in Hz. [α]_D values are given in 10⁻¹ deg cm² g⁻¹. Elemental analyses were performed on a Carlo Erba-1106 instrument.

Preparation of dipeptide alcohols 3a-p and (3d + 3d'); general procedure

To a mixture of *N*-protected(α -aminoacyl)benzotriazoles **1** (1 mmol) in dry THF (10 mL) was added the α -amino alcohol **2** (1 mmol) and the mixture was stirred for 6 h at room temperature. The solvent was evaporated under vacuum and the residue was dissolved in ethyl acetate (15 mL), washed with Na₂CO₃ (3 × 5 mL), brine (1 × 5 mL), water (1 × 5 mL) and dried over Na₂SO₄. The solvent was evaporated and the solid residue was crystallized from ethyl acetate–hexanes to afford the corresponding dipeptide alcohols **3**.

N-Boc-L-Val-L-Phenylalaninol 3a^{1*a*}. (Yield 0.19 g, 86%.) White microcrystals; mp 145.0–146.0 °C (EtOAc–hexanes). $[\alpha]_D^{24}$ –49.4 (*c* 1.0 in MeOH). (Found: C, 65.26; H, 8.94; N, 7.99. Calc. for C₁₉H₃₀N₂O₄: C, 65.12; H, 8.63; N, 7.99%.) $\delta_{\rm H}$ (300 MHz; CDCl₃; Me₄Si): 7.37–7.26 (5 H, m, Ar*H*), 6.32 (1 H, d, *J* 7.2, N*H*), 4.98 (1 H, br s, N*H*), 4.24 (1 H, br s, O*H*), 3.89 (1 H, t, *J* 6.0, CH₂C*H*), 3.77–3.59 (2 H, m, PhCH₂), 3.00–2.86 (2 H, m, OCH₂), 2.22–2.11 (1 H, m, (CH₃)₂C*H*), 1.82 (1 H, br s, O*H*), 1.50 (9 H, s, 3 × CCH₃), 0.96 (3 H, d, *J* 6.9, CH(CH₃)) and 0.87 (3 H, d, *J* 5.7, CH(CH₃));

 $\delta_{\rm C}$ (75 MHz; CDCl₃; Me₄Si): 172.0, 156.2, 137.8, 129.3, 128.6, 126.6, 80.2, 63.4, 60.6, 52.9, 37.0, 30.6, 28.4, 19.3 and 17.9.

Preparation of tripeptide alcohols 6a-c; general procedure

To a solution of *N*-Fmoc-(α -dipeptidoyl)benzotriazole **5** (0.29 mmol) in THF (5 mL), amino alcohol (0.29 mmol) was added at 0 °C. The reaction mixture was stirred at 0 °C for 4 h followed by 3 h at room temperature. Hexanes (5 mL) were added and the solid separated was washed with diethyl ether (2 × 10 mL) and dried to give the corresponding tripeptide alcohol **6**.

Fmoc-L-Phe-L-Ala-L-Phenylalaninol 6a. (Yield 0.11 g, 69%.) White microcrystals; mp 185.0–186.0 °C (from tetrahydrofuranhexanes). [α]_D²⁴ –67.0 (*c* 1.0 in MeOH). (Found: C, 71.29; H, 6.25; N, 6.83. Calc. for C₃₆H₃₇N₃O₅H₂O: C, 70.92; H, 6.12; N, 6.89%.) $\delta_{\rm H}$ (300 MHz; DMSO-d₆; Me₄Si): 8.12 (1 H, d, *J* 7.2, N*H*), 7.87 (2 H, d, *J* 7.5, Ar*H*), 7.73 (1 H, d, *J* 8.1, N*H*), 7.61–7.66 (2 H, m, Ar*H*), 7.41(2 H, t, *J* 8.1, Ar*H*), 7.15–7.34 (13 H, m, Ar*H*, NH), 4.81 (1 H, br s, C*H*), 4.25–4.29 (2 H, m, C*H*₂), 4.13–4.17 (4 H, m, 2 × C*H*₂), 3.82–3.89 (1 H, m, C*H*), 2.62–3.01 (5 H, m, C*H*₂, 2 × C*H*, O*H*) and 1.18 (3 H, d, *J* 6.9, C*H*₃); $\delta_{\rm C}$ (75 MHz: DMSO-d₆; Me₄Si): 171.8, 171.2, 155.9, 143.8, 140.7, 138.3, 129.3, 129.2, 128.2, 128.1, 127.7, 127.2, 126.3, 126.0, 125.3, 120.2, 65.7, 62.1, 56.1, 52.4, 48.41, 46.6, 37.5, 36.4 and 18.6.

Acknowledgements

We thank Dr C. D. Hall for helpful discussions and Dr Q.-Y. Chen for some preliminary experiments.

Notes and references

- (a) A. D. Abell, M. A. Jones, A. T. Neffe, S. G. Aitken, T. P. Cain, R. J. Payne, S. B. McNabb, J. M. Coxon, B. G. Stuart, D. Pearson, H. Y.-Y. Lee and J. D. Morton, J. Med. Chem., 2007, 50, 2916; (b) T. L. Voyushina, J. V. Potetinova, E. I. Milgotina and V. M. Stepanov, Bioorg. Med. Chem., 1999, 7, 2953; (c) X. Zhang, J. Rodrigues, L. Evans, B. Hinkle, L. Ballantyne and M. Peña, J. Org. Chem., 1997, 62, 6420; (d) B. Kammermeier, G. Beck, W. Holla, D. Jacobi, B. Napierski and H. Jendralla, Chem.-Eur. J., 1996, 2, 307; (e) A. Vinitsky, C. Cardozo, L. Sepp-Lorenzino, C. Michaudt and M. Orlowski, J. Biol. Chem., 1994, 269, 29860; (f) J. A. Read and C. T. Walsh, J. Am. Chem. Soc., 2007, 129, 15762; (g) V. Schanen, C. Riche, A. Chiaroni, J.-C. Quirion and H.-P. Husson, Tetrahedron Lett., 1994, 35, 2533.
- 2 S. Wilk and W.-E. Chen, Mol. Biol. Rep., 1997, 24, 119.
- 3 (a) A. S. Y. Yim and M. Wills, *Tetrahedron*, 2005, **61**, 7994; (b) I. M. Pastor, P. Västilä and H. Adolfsson, *Chem.-Eur. J.*, 2003, **9**, 4031; (c) I. M. Pastor, P. Västilä and H. Adolfsson, *Chem. Commun.*, 2002, 2046; (d) H.-Q. Cai, C. Chen, L. Liu, J.-M. Ni and R. Wang, *J. Mol. Catal. A: Chem.*, 2006, **253**, 86; (e) P. Västilä, I. M. Pastor and H. Adolfsson, *J. Org. Chem.*, 2005, **70**, 2921; (f) I. M. Pastor and H. Adolfsson, *Tetrahedron Lett.*, 2002, **43**, 1743; (g) S. R. Gilbertson and P. Lan, *Tetrahedron Lett.*, 2002, **43**, 6961.

- 4 M. Segura, B. Bricoli, A. Casnati, E. M. Muñoz, F. Sansone, R. Ungaro and C. Vicent, J. Org. Chem., 2003, 68, 6296.
- 5 P. Liu, B. Sun, X. Chen, G. Tian and Y. Ye, *Synth. Commun.*, 2002, **32**, 473.
- 6 M. L. Di Gioia, A. Leggio, A. L. Pera, A. Liguori and C. Siciliano, *Eur. J. Org. Chem.*, 2004, 463.
- 7 K. Soai, H. Oyamada and M. Takase, M., *Bull. Chem. Soc. Jpn.*, 1984, 57, 2327.
- 8 J. V. Potetinova, T. L. Voyushina and V. M. Stepanov, *Bioorg. Med. Chem. Lett.*, 1997, 7, 705.
- 9 J. Swistok, J. W. Tilley, W. Danho, R. Wagner and K. Mulkerins, *Tetrahedron Lett.*, 1989, **30**, 5045.
- 10 (a) A. R. Katritzky, P. Angrish, D. Hur and K. Suzuki, Synthesis, 2005, 397; (b) A. R. Katritzky, P. Angrish and K. Suzuki, Synthesis, 2006, 411; (c) A. R. Katritzky, K. Suzuki, S. K. Singh and H. Y. He, J. Org. Chem., 2003, 68, 5720; (d) A. R. Katritzky, H. Y. He and K. Suzuki, J. Org. Chem., 2000, 65, 8210; (e) A. R. Katritzky, M. Y. Wang, H. F. Yang, S. M. Zhang and N. G. Akhmedov, ARKIVOC, 2002, viii, 134; (f) A. R. Katritzky, A. A. Shestopalov and K. Suzuki, ARKIVOC, 2005, vii, 36; (g) A. R. Katritzky, S. R. Tala and S. K.

Singh, *Synthesis*, 2006, 3231; (*h*) A. R. Katritzky, R. Chen and Q-. Tala S. R., *Org. Biomol. Chem.*, 2008, **6**, 2400; (*i*) A. R. Katritzky, A. Singh, D. N. Haase and M. Yoshioka, *Arkivoc*, 2009, **viii**, 47; (*j*) A. R. Katritzky, S. R. Tala, N. Abo-Dya and Z. K. Abdel-Samii, *Synthesis*, 2009, 1708; (*k*) A. R. Katritzky and S. A. Belyakov, *Aldrichimica Acta*, 1998, **31**, 35; (*l*) A. R. Katritzky, J. Li and J. Xie, *Tetrahedron*, 1999, **55**, 8263.

- 11 (a) A. R. Katritzky, S. Kazuyuki and S. K. Singh, *Synthesis*, 2004, 2645; (b) A. R. Katritzky, Q. Chen and S. R. Tala, *Chem. Biol. Drug Des.*, 2009, **73**, 611.
- 12 (a) A. R. Katritzky, N. M. Khashab, M. Yoshioka, D. N. Haase, K. R. Wilson, J. V. Johnson, A. Chung and C. Haskell-Luevano, *Chem. Biol. Drug Des.*, 2007, **70**, 465; (b) A. R. Katritzky, M. Yoshioka, T. Narindoshvili, A. Chung and N. M. Khashab, *Chem. Biol. Drug Des.*, 2008, **72**, 182.
- 13 M. Rainaldi, A. Moretto, C. Peggion, F. Formaggio, S. Mammi, E. Peggion, J. A. Galvez, M. D. Diaz-de-Villegas, C. Cativiela and C. Toniolo, *Chem.-Eur. J.*, 2003, 9, 3567.
- 14 T. C. Barden, B. L. Buckwatter, R. T. Testa, P. J. Peterson and V. J. Lee, J. Med. Chem., 1994, 37, 3205.