

Synthesis and Transformations of Methyl (*E*)-2-(Acetylamino)-3-cyanoprop-2-enoate und Methyl (*E*)-2-(Benzoylamino)-3-cyanoprop-2-enoate, Versatile Reagents for the Preparation of Polyfunctional Heterocyclic Systems

by Lucija Pizzioli, Brina Ornik, Jurij Svete, and Branko Stanovnik*


Faculty of Chemistry and Chemical Technology, University of Ljubljana, Aškerčeva 5, POB 537,
1000 Ljubljana, Slovenia

Methyl (*E*)-2-(acetylamino)-3-cyanoprop-2-enoate (**2a**) and its 2-benzoyl analog **2b** were prepared from the corresponding methyl (*Z*)-2-(acylamino)-3-(dimethylamino)propenoates **1**. Multifunctional compounds **2** are versatile synthons for preparation of polysubstituted heterocyclic systems such as pyrroles **4**, pyrimidines **5** and **6**, pyridazines **7**, pyrazoles **8**, **9**, and **11**, and isoxazoles **10**.

Introduction. – Derivatives of 2-amino-3-cyanoprop-2-enoates have been prepared from dichloromaleic anhydride [1] and by condensation of alkyl cyanofornates and analogous compounds with active methylene compounds [2–4]. Inspite of the fact that these compounds have been described as precursors for the synthesis of α,β -didehydro- α -amino acids, reactions involving either CN or ester group have not been reported.

Results and Discussion. – In our studies on the syntheses and transformations of alkyl 2-(acylamino)-3-(dimethylamino)prop-2-enoates as simple and versatile reagents for the preparation of various cyclic systems [5][6], we observed that the Me₂N group in methyl (*Z*)-2-(acetylamino)-3-(dimethylamino)prop-2-enoate (**1a**) and methyl (*Z*)-2-(benzoylamino)-3-(dimethylamino)prop-2-enoate (**1b**), obtained from *N*-acetylglutamine [6] and *N*-benzoylglutamine (hippuric acid) [7][8], respectively, can be easily exchanged by a CN group to give methyl (*E*)-2-(acetylamino)-3-cyanoprop-2-enoate (**2a**) and methyl (*E*)-2-(benzoylamino)-3-cyanoprop-2-enoate (**2b**) in 70 and 52 % yield, respectively (Scheme 1). The configuration around the C=C bond in compounds **2** was established by X-ray analysis¹⁾.


Since the compounds **2** are multifunctional, they turned out to be useful synthons for the preparation of several polyfunctional substituted five- and six-membered heterocyclic



¹⁾ X-Ray data of structure analyses of the compounds **2a** and **2b** will be published elsewhere.

systems. There are several types of synthetic methods for preparation of pyrroles [9], pyrimidines [10–11], pyridazines [12–14], pyrazoles [15][16], and isoxazoles [17][18]. Especially, systems with an amino, cyano, carboxylic, and/or ester group(s) require tedious, multistep procedures, resulting usually in low overall yields.

In the reaction of **2** with NH_3 , CN , and COOMe groups are involved in cyclization to give pyrrole derivative **4**, while in the reaction with primary amines CN and AcNH groups react to give substituted pyrimidine derivatives **5** and **6**. The reaction of hydrazine



with CN and COOMe groups produced 4,6-diamino-substituted pyridazin-3(2*H*)-ones **7**.

1,3-Dipolar cycloaddition of CH_2N_2 to the activated $\text{C}=\text{C}$ bond takes place under mild conditions to form methyl 3-(acetylamino)-4-cyano-4,5-dihydro-3*H*-pyrazole-3-carboxylate (**8a**) and its 3-benzoylamino analog **8b**, as the primary cycloadducts (Scheme 2). Thermal elimination of acetamide from **8a** leads to the 3,4-disubstituted pyrazole derivative **9**. On the other hand, the 1,3-dipolar cycloaddition of nitrile oxides **10** requires more drastic conditions, under which acetamide or benzamide is eliminated spontaneously to give the corresponding substituted isoxazole derivative **11**. Similarly, 1,3-dipolar cycloaddition of diphenylnitrile imine, generated *in situ* from *N*-phenylbenzohydrazonoyl chloride in the presence of Et_3N , yields methyl 4-cyano-2,5-diphenylpyrazole-3-carboxylate (**12**) by elimination of benzamide from the primary cycloadduct, which was not isolated. The structures of compounds were determined by NMR spectra, elemental analyses, and, in some instances, by X-ray structure analyses.

Experimental Part

General. M.p.: Kofler micro hot stage. The ^1H -NMR spectra: Varian EM 360 L and Bruker Avance 300 DPX spectrometers. MS: Autospeck Q spectrometer. Elemental analyses: Perkin-Elmer Analyser 2400. Methyl (*Z*)-2-(acetylamino)-3-(dimethylamino)prop-2-enoate (**1a**) [6] and methyl (*Z*)-2-(benzoylamino)-3-(dimethylamino)-prop-2-enoate (**1b**) [7] were prepared according to literature procedures.

Methyl (E)-2-(Acetylamino)-3-cyanoprop-2-enoate (2a). To a mixture of **1a** (0.930 g, 5 mmol), KCN (0.358 g, 5.5 mmol), and toluene (15 ml), glacial AcOH (15 ml) was added dropwise. Then, the mixture was heated under reflux for 1.5 h, cooled to r.t., and the solvent was evaporated *in vacuo*. To the oily residue, H_2O (30 ml) was added and the precipitate collected by filtration to give **2a** in 71% yield. M.p. 115–116°. ^1H -NMR (300 MHz, CDCl_3): 2.23 (s, COMe); 4.03 (s, MeO); 7.37 (s, CH); 8.27 (br. s, NH). MS: 168 (29, M^+). Anal. calc. for $\text{C}_7\text{H}_{10}\text{N}_2\text{O}_3$ (168.15): C 50.00, H 4.80, N 16.66; found: C 50.20, H 4.61, N 16.55. X-Ray analysis confirms the (*E*)/configuration.)

Methyl (E)-2-(Benzoylamino)-3-cyanoprop-2-enoate (2b). Similarly, **2b** was prepared from **1b** in 52% yield. M.p. 96–98° (ProH). ^1H -NMR (60 MHz, $(\text{D}_6)\text{DMSO}$): 3.87 (s, MeO); 6.10 (s, CH); 7.50–7.80 (*m*, H–C(3'), H–C(4'), H–C(5') of Ph); 7.93–8.10 (*m*, H–C(2'), H–C(6') of Ph); *NH*COPh exchanged. Anal. calc. for $\text{C}_{12}\text{H}_{10}\text{N}_2\text{O}_3$ (230.22): C 62.61, H 4.38, N 12.17; found: C 62.30, H 4.35, N 12.11.

Methyl (RS)-2-(Acetylamino)-3-cyanopropanoate (3a). To a soln. of **2a** (0.168 g, 1 mmol) in EtOH (10 ml), Pd/C (10%, 0.020 g) was added and the mixture was hydrogenated at normal pressure for 5 h. The filtrate was evaporated *in vacuo*. Et_2O (5 ml) was added to the oily residue, and the precipitate was collected by filtration to give **3a** in 98% yield. M.p. 72–75°. ^1H -NMR (300 MHz, CDCl_3): 2.09 (s, COMe); 2.95 (*dd*, $J = 4.3$, 17, H–C(3)); 3.09 (*dd*, $J = 5.3$, 17, H–C(3)); 3.87 (s, MeO); 4.73–4.78 (br. *ddd*, $J = 4.3$, 5.3, 17, H–C(2)); 6.43 (br. s, NH). Anal. calc. for $\text{C}_7\text{H}_{10}\text{N}_2\text{O}_3$ (170.17): C 49.41, H 5.92, N 16.46; found: C 49.16, H 5.71, N 16.42.

Methyl (RS)-2-(Benzoylamino)-3-cyanopropanoate (3b). Similarly, **3b** was prepared from **2b** in 99% yield. M.p. 135–138° (from AcOEt/MeOH). ^1H -NMR (300 MHz, CDCl_3): 3.06 (*dd*, $J = 4.4$, 17, H–C(3)); 3.23 (*dd*, $J = 5.5$, 17, H–C(3)); 3.80 (s, MeO); 4.93–4.98 (*m*, $J = 4.4$, 5.5, 5.6, 17, H–C(2)); 7.17 (*d*, $J = 5.6$, NH); 7.45–7.56 (*m*, H–C(3'), H–C(4'), H–C(5') of Ph); 7.82–8.85 (*m*, H–C(2'), H–C(6') of Ph). Anal. calc. for $\text{C}_{12}\text{H}_{12}\text{N}_2\text{O}_3$ (232.24): C 62.06, H 5.21, N 12.06; found: C 61.74, H 5.13, N 12.03.

*3-(Acetylamino)-5-imino-1,5-dihydro-2*H*-pyrrol-2-one (4a).* A soln. of **2a** (0.168 g, 1 mmol) in EtOH (5 ml) was treated with NH_3 for 10 min. The soln. was cooled on ice, and the precipitate was collected by filtration to give **4a** in 53% yield. M.p. > 350°. ^1H -NMR (300 MHz, $(\text{D}_6)\text{DMSO}$): 2.12 (s, COMe); 6.60 (s, CH); 9.11 (br. s, *NH*COMe); 9.87 (br. s, NH); 10.22 (br. s, NH). Anal. calc. for $\text{C}_6\text{H}_7\text{N}_3\text{O}_2$ (153.14): C 47.06, H 4.61, N 27.44; found: C 47.10, H 4.55, N 27.45.

*3-(Benzoylamino)-5-imino-1,5-dihydro-2*H*-pyrrol-2-one (4b).* Similarly, **4b** was prepared from **2b** in 45% yield. M.p. > 230° (dec.). ^1H -NMR (60 MHz, $(\text{D}_6)\text{DMSO}$): 6.83 (s, H–C(4)); 7.47–7.67 (*m*, H–C(3'), H–C(4'), H–C(5') of Ph); 7.90–8.10 (*m*, H–C(2'), H–C(6') of Ph); 8.70–9.80 (br. s, *NH*COPh, NH); 10.23 (br. s, NH). Anal. calc. for $\text{C}_{11}\text{H}_9\text{N}_3\text{O}_2$ (215.21): C 61.39, H 4.22, N 19.53; found: C 61.18, H 4.20, N 19.82.

Methyl 2-Methyl-6-(methylamino)pyrimidine-4-carboxylate (5). A soln. of **2a** (0.168 g, 1 mmol) and MeNH₂ (40%, 0.0775 g) in MeOH (3 ml) was heated under reflux for 3 h. The solvent was evaporated *in vacuo*, and the solid residue was purified by radial chromatography using *Chromatotron* (CHCl₃/AcOEt 3:2) to give **5** in 19% yield. M.p. 195–198°. ¹H-NMR (300 MHz, CDCl₃): 2.24 (s, Me); 3.11 (s, NHMe); 3.41 (s, MeO); 7.04 (s, CH); 7.92 (br. s, NH). MS: 181 (54, M⁺). Anal. calc. for C₈H₁₁N₃O₂ (181.19): C 53.03, H 6.12, N 23.19; found: C 53.33, H 6.18, N 23.17.

N-Benzyl-6-(benzylamino)-2-methylpyrimidine-4-carboxamide (6a). A soln. of **2a** (0.168 g, 1 mmol) and PhCH₂NH₂ (0.5 ml, 5 mmol) in EtOH (3 ml) was heated under reflux for 4 h. The soln. was left overnight at r.t., and the precipitate was collected by filtration to give **6a** in 38% yield. M.p. 155–156°. ¹H-NMR (300 MHz, CDCl₃): 2.48 (s, Me); 4.57, 4.59 (2s, 2 CH₂); 5.63 (br. t, NH); 7.11 (s, H-C(5)); 7.25–7.36 (m, 2 Ph); 8.36 (br. t, CONH). Anal. calc. for C₂₀H₂₀N₄O (332.40): C 72.27, H 6.06, N 16.85; found: C 72.10, H 6.12, N 16.79.

N-Benzyl-6-(benzylamino)-2-phenylpyrimidine-4-carboxamide (6b). Similarly, **6b** was prepared from **2b** and PhCH₂NH₂ in 47% yield. M.p. 208–210° (EtOH). ¹H-NMR (60 MHz, (D₆)DMSO): 4.63 (d, J = 7, CH₂NH); 4.80 (d, J = 6, CH₂NHCO); 7.17 (s, H-C(5)); 7.30–7.57 (m, H-C(3'), H-C(4'), H-C(5') of CONHCH₂Ph, 2 PhCH₂); 8.40 (t, J = 7, NHCH₂); 8.47–8.67 (m, H-C(2'), H-C(6') of Ph-C(2)); 9.47 (t, J = 6, CONHCH₂). Anal. calc. for C₂₅H₂₂N₄O (394.48): C 76.12, H 5.62, N 14.20; found: C 75.83, H 5.56, N 14.03.

4-(Acetylamino)-6-amino-2H-pyridazin-3-one (7a). A mixture of **2a** (0.168 g, 1 mmol) and hydrazine hydrate (0.1 ml, 2 mmol) in EtOH (3 ml) was stirred at r.t. for 2 h. The precipitate was collected by filtration and recrystallized from EtOH to give **7a** in 62% yield. M.p. 235°. ¹H-NMR (300 MHz, (D₆)DMSO): 2.17 (s, COMe); 5.54 (s, NH₂); 7.69 (s, CH); 9.55 (s, NHCOMe); 11.88 (br. s, NH). MS: 168 (49, M⁺). Anal. calc. for C₆H₈N₄O₂ (168.16): C 42.86, H 4.80, N 33.32; found: C 42.71, H 4.85, N 32.95.

6-Amino-4-(benzoylamino)-2H-pyridazin-3-one (7b). Similarly, **7b** was prepared from **2b** in 81% yield. M.p. 284–286° (MeOH). ¹H-NMR (60 MHz, (D₆)DMSO): 5.60 (br. s, NH₂); 7.57–7.77 (m, H-C(3'), H-C(4'), H-C(5') of Ph); 7.90 (s, CH); 7.90–8.07 (m, H-C(2'), H-C(6') of Ph); 9.57 (s, NHCOPh); 12.20 (br. s, NH). Anal. calc. for C₁₁H₁₀N₄O₂ (230.23): C 57.39, H 4.38, N 24.34; found: C 57.35, H 4.41, N 24.44.

Methyl 3-(Acetylamino)-4-cyano-4,5-dihydro-3H-pyrazole-3-carboxylate (8a). To a soln. of **2a** (0.168 g, 1 mmol) in CHCl₃ (3 ml), excess of CH₃N₂ (in Et₂O, 10 mmol) was added and left overnight at –20°. The mixture was left for a few hours in the open air. The precipitate was collected by filtration to give **8a** in 58% yield. M.p. 128–132°. ¹H-NMR (300 MHz, (D₆)DMSO): 1.98 (s, COMe); 3.64 (dd, J = 7.4, 9.6, H-C(4)); 3.77 (s, MeO); 4.83 (dd, J = 7.4, 18, H-C(5)); 5.32 (dd, J = 9.6, 18, H-C(5)); 9.38 (s, NH). MS: 211 (43, M⁺). Anal. calc. for C₈H₁₀N₄O₃ (210.19): C 45.71, H 4.80, N 26.66; found: C 45.93, H 4.84, N 26.54.

Methyl 3-(Benzoylamino)-4-cyano-4,5-dihydro-3H-pyrazole-3-carboxylate (8b). Similarly, **8b** was prepared from **2b** in 86% yield. M.p. 98° (CH₂Cl₂/Et₂O). ¹H-NMR (60 MHz, (D₆)DMSO, D₂O): 3.80 (deg. dd and s, J = 7, 10, H-C(4), MeO); 4.83 (dd, J = 7, 18, H-C(5)); 5.45 (dd, J = 10, 18, H-C(5)); 7.50–7.70 (m, H-C(3'), H-C(4'), H-C(5') of Ph); 7.87–8.07 (m, H-C(2'), H-C(6') of Ph); 9.83 (s, NHCOPh, exchanged). Anal. calc. for C₁₃H₁₂N₄O₃ (272.26): C 57.35, H 4.44, N 20.58; found: C 57.30, H 4.52, N 20.35.

Methyl 4-Cyanopyrazole-3-carboxylate (9). The compound **8a** (0.210 g, 1 mmol) in xylene (3 ml) was heated under reflux for 3 h. The precipitate was collected by filtration to give **9** in 79% yield. M.p. 221–227° ([19]: m.p. 225–228°). ¹H-NMR (300 MHz, (D₆)DMSO): 3.88 (s, MeO); 8.69 (s, CH); 14.40 (s, NH). MS: 151 (72, M⁺). Anal. calc. for C₆H₅N₃O₂ (151.12): C 47.69, H 3.33, N 27.80; found: C 47.33, H 3.15, N 27.89.

Methyl 4-Cyano-3-(2,4,6-trimethoxyphenyl)isoxazole-5-carboxylate (11). A mixture of **2a** (0.168 g, 1 mmol) and 2,4,6-trimethoxybenzenecarbonitrile oxide (**10**; 0.209 g, 1 mmol) in CHCl₃ (3 ml) was heated under reflux for 6 h. The solvent was evaporated *in vacuo*, and MeOH (1 ml) was added to the oily residue. The precipitate was collected by filtration to give **11** in 72% yield. M.p. 129–131°. ¹H-NMR (300 MHz, CDCl₃): 3.82 (s, 2,6'-(OMe)₂); 3.88 (s, 4'-OMe); 4.07 (s, COOMe); 6.21 (s, H-C(3'), H-C(5') of Ar). Anal. calc. for C₁₅H₁₄N₂O₆ (318.29): C 56.60, H 4.43, N 8.80; found: C 56.71, H 4.44, N 8.45.

The compound **11** was obtained also from **4b** in 71% yield.

Methyl 4-Cyano-1,3-diphenylpyrazole-5-carboxylate (12). To a mixture of **2b** (0.230 g, 1 mmol) and *N*-phenylbenzohydrazonyl chloride (0.230 g, 1 mmol) in CH₂Cl₂ (3 ml), Et₃N (0.2 ml) was added, and the mixture was heated under reflux for 0.5 h. The solvent was evaporated *in vacuo*. The oily residue was dissolved in EtOH (3 ml), H₂O (5 ml) was added and the precipitate was collected by filtration to give **11** in 81% yield. M.p. 172–174°. ¹H-NMR (60 MHz, (D₆)DMSO): 3.87 (s, MeO); 7.50–7.67 (m, Ph-N(1), H-C(3'), H-C(4'), H-C(5') of Ph-C(3)); 7.93–8.13 (m, H-C(2'), H-C(6') of Ph-C(3)). Anal. calc. for C₁₈H₁₃N₃O₂ (303.32): C 71.28, H 4.32, N 13.85; found: C 70.80, H 4.29, N 14.03.

The authors wish to express their gratitude to the *Ministry of Science and Technology*, Slovenia, for their financial support.

REFERENCES

- [1] G. Bonse, H. U. Blank, *Liebigs Ann. Chem.* **1981**, 1658.
- [2] T. Iimori, Y. Nii, T. Izawa, S. Kobayashi, M. Ohno, *Tetrahedron Lett.* **1979**, 27, 2525–2528.
- [3] Jpn Kokai Tokkyo Koho JP 82 67.545; *Chem. Abstr.* **1982**, 97, 198561.
- [4] Jpn Kokai Tokkyo Koho JP 81 87.541; *Chem. Abstr.* **1981**, 95, 204437.
- [5] a) B. Stanovnik, 'Methyl 2-Benzoylamino-3-dimethylaminopropenoate in the Synthesis of Heterocyclic Systems', in 'Progress in Heterocyclic Chemistry', Eds. H. Suschitzky and E. F. V. Scriven, Pergamon Press, Oxford, 1993, Vol. 5, pp. 34–53; b) B. Stanovnik, *Molecules* **1996**, 1, 123.
- [6] L. Kralj, A. Hvala, J. Svete, L. Golič, B. Stanovnik, *J. Heterocycl. Chem.* **1997**, 34, 247.
- [7] B. Stanovnik, J. Svete, M. Tišler, L. Žorž, A. Hvala, I. Simonič, *Heterocycles* **1988**, 27, 903.
- [8] Japan Kokai, 75 58063; *Chem. Abstr.* **1975**, 83, 193075.
- [9] a) R. J. Sundberg, 'Pyrroles and their Benzo Derivatives: Synthesis and Applications, in Comprehensive Heterocyclic Chemistry', Eds. A. R. Katritzky and C. W. Rees, Vol. 4, ed. by C. W. Bird and G. W. H. Cheeseman, Pergamon, Oxford–New York, 1984, pp. 313–376; b) 'Pyrroles and their Benzo Derivatives: Synthesis', in Comprehensive Heterocyclic Chemistry II', Eds. A. R. Katritzky, C. W. Rees, and E. F. V. Scriven, Vol. 2, ed. by C. W. Bird, Pergamon, Oxford, 1996, pp. 119–206.
- [10] D. J. Brown, 'Pyrimidines and their Benzo Derivatives, in Comprehensive Heterocyclic Chemistry', Eds. A. R. Katritzky and C. W. Rees, Vol. 3, ed. by A. J. Boulton and A. McKillop, Pergamon, Oxford–New York, 1984, pp. 57–155.
- [11] D. J. Brown, R. F. Evans, W. B. Cowden, M. D. Fenn, 'The Pyrimidines', John Wiley and Sons, New York, 1994, pp. 49–239.
- [12] a) M. Tišler, B. Stanovnik, 'Pyridazines and their Benzo Derivatives', in 'Comprehensive Heterocyclic Chemistry', Eds. A. R. Katritzky and C. W. Rees, Vol. 3, ed. by A. J. Boulton and A. McKillop, Pergamon, Oxford–New York, 1984, pp. 1–56; b) W. J. Coates, 'Pyridazines and their Benzo Derivatives', in 'Comprehensive Heterocyclic Chemistry II', ed. by A. R. Katritzky, C. W. Rees and E. F. V. Scriven, Vol. 6, ed. by A. J. Boulton, Pergamon, Oxford, 1996, pp. 1–91.
- [13] 'Pyridazines', Ed. R. N. Castle, John Wiley and Sons, New York, 1973.
- [14] M. Tišler, B. Stanovnik, 'Advances in Pyridazine Chemistry', *Adv. Heterocycl. Chem.* **1990**, 49, 385–474, and ref. cit. therein.
- [15] a) J. Elguero, 'Pyrazoles and their Benzo Derivatives', in 'Comprehensive Heterocyclic Chemistry', Eds. A. R. Katritzky and C. W. Rees, Vol. 5, ed. by K. T. Potts, Pergamon, Oxford, New York, 1984, pp. 167–303; b) 'Pyrazoles', in 'Comprehensive Heterocyclic Chemistry II', Eds. A. R. Katritzky, C. W. Rees, and E. F. V. Scriven, Vol. 3, ed. by I. Shinkai, Pergamon, Oxford, 1996, pp. 1–75.
- [16] R. H. Wiley, L. C. Behr, R. Fusco, C. H. Jarbe, 'Pyrazoles, Pyrazolines, Pyrazolidines, Indazoles and Condensed Rings', John Wiley and Sons, New York, 1967.
- [17] S. A. Lang, Jr., Y.-i Lin, 'Isoxazoles and their Benzo Derivatives', in 'Comprehensive Heterocyclic Chemistry', Eds. A. R. Katritzky and C. W. Rees, Vol. 6, ed. by K. T. Potts, Pergamon, Oxford–New York, 1984, pp. 1–130.
- [18] a) P. Grünanger, P. Vita-Finzi, 'Isoxazoles', John Wiley and Sons, New York, 1991; b) M. Sutharchanadevi, R. Murugan, Isoxazoles, in 'Comprehensive Heterocyclic Chemistry II', Eds. A. R. Katritzky, C. W. Rees and E. F. V. Scriven, Vol. 3, ed. by I. Shinkai, Pergamon, Oxford, 1996, pp. 221–260.
- [19] R. Huisgen, A. Mitra, J. R. Moran, *Heterocycles* **1986**, 24, 2429.

Received September 30, 1997