

Tetrahedron 54 (1998) 2459-2476

SYNTHESIS OF NEW 2*H*-1,4-THIAZINES AND THEIR DERIVATIVES UTILIZING *N*,*N*-(DI-*N*-ALKYL)-*N*'-ARYLTHIOCARBAMOYLAMIDINES

Moon-Kook Jeon and Kyongtae Kim*

Department of Chemistry, Seoul National University, Seoul 151-742

Sung Hoon Kim

Korea Institute of Science and Technology, P. O. Box 131, Cheongyang, Seoul 130-650, Korea

Received 8 September 1997; accepted 26 December 1997

Abstract: The reactions of N,N-(di-n-alkyl)-N'-arylthiocarbamoylamidines (1) with 2-bromo-1-phenylethanone in the presence of K₂CO₃ in THF at reflux gave 3-(di-n-alkylamino)-2-arylimino-5-phenyl-2H-1,4-thiazines (7) in 32 to 62 % yields. Treatment of compounds 1 with bromoacetyl bromide in the presence of pyridine in CH₂Cl₂ at 0 °C afforded 5-(di-n-alkylamino)-6-arylimino-2H-1,4-thiazin-3-ones (12) in 41 to 84 % yields, whereas the same reactions of 1 with 2-bromopropionyl bromide under the same conditions gave 4-(di-n-alkylamino)-5-arylimino-2-(1-bromoethylidene)-2H-thiazolines (17) as minor compounds in addition to thiazin-3-ones 16, analogous to compounds 12. The reactions of 1 with ethyl bromoacetate in CH₂Cl₂ at room temperature, however, gave [(arylimino)(S-ethoxycarbonylmethyl)]methyl-N,N-(di-n-propyl)amidine hydrobromides (19) in 71 to 88 % yields. Compounds 7, 12, 16, 17, and 19 are all new and the mechanisms of their formations are proposed. © 1998 Elsevier Science Ltd. All rights reserved.

We have previously described the synthesis of N,N-(di-n-alkyl)-N²-arylthiocarbamoylamidines (1)¹ and demonstrated their synthetic utilities by preparing 4-dialkylamino-5-aryliminothiazoline-2-thiones (2), 4-dialkylamino-5-arylimino-5H-2-oxo-1,2,3-dithiazoles (3), 4-dialkylamino-5-arylimino-5H-2,2-dioxo-1,2,3-dithiazoles (3), 4-dialkylamino-5-arylimino-5H-2,2-dioxo-1,2,3-dithiazoles (4), 4-dialkylamino-5-arylimino-2-phenyliminothiazolines (5), and 4-dialkylamino-3-arylimino-2,5-benzothiazocine-1,6-diones (6).² All of these compounds 2-5, having a five-membered cyclic skeleton except for compound 6, were prepared from compounds 1, which have four-atom unit as nucleophiles, reacting with one-atom unit such as thiophosgene, thionyl chloride, sulfuryl chloride, and N-phenylimidoyl dichloride as electrophiles.



0040-4020/98/\$19.00 © 1998 Elsevier Science Ltd. All rights reserved. *PII:* S0040-4020(98)00010-6

In a connection with the study to explore the potential synthetic utilities of compounds 1, it was our aim to prepare 1,4-thiazine derivatives, which have attracted much attention due to the potential biological activities such as antibacterial³ and cardiotonic⁴ activities, by the cyclization of compounds 1 with two-carbon atom unit.

Of the 1,4-thiazines, 2H-1,4-thiazines have been most extensively studied.⁵ However, the methods of synthesis are rather limited. Treatment of bis(aroylmethyl)sulfide with ammonia gas gave 3,5-diaryl-2H-1,4-thiazines.⁶ 2-(Dialkylhydrazono)thioacetophenones react with acetylenic dienophiles such as methyl propiol-ate or dimethyl acetylenedicarboxylate to give 2-amino-2H-1,4-thiazines after amino group transposition.⁷ Similarly, 2-hydrazonothioacetophenones are cyclized with methyl vinyl ketones in benzene with a few crystals of hydroquinone added, to afford 3-acetyl-3,4-dihydro-2H-1,4-thiazines, which undergo thermal elimination reaction at 60°C to give the corresponding 2H-1,4-thiazines.⁸ In the presence of silica, 4-dimethylamino-3,4-dihydro-2H-1,4-thiazines afford 2H-1,4-thiazines.⁹

We have studied the reactions of compounds 1 with α -monohaloketones, α -monohaloacyl halides, and α -monobromo ester in order to prepare new 2H-1,4-thiazine derivatives. The results are described herein.

RESULTS AND DISCUSSION

Reactions with *a*-Monohaloketones

The reactions of 1 with 2-bromo-1-phenylethanone in the presence of K_2CO_3 in THF at reflux gave 3-(di*n*-alkylamino)-2-arylimino-5-phenyl-2*H*-1,4-thiazines 7 as major products. Reaction conditions, yields, and physical properties of compounds 7 are summarized in Table 1 and their analytical and spectroscopic data in Table 2. Treatment of compound 1c (X = Me, R = *n*-Pr) with pyridine (2 equiv.) in CH₂Cl₂ for 12 days at reflux gave 7c in 52 %yield, which was lower than the yield obtained under the foregoing condition.



The structures of compounds 7a-i were determined on the basis of the spectroscopic data and elemental analyses. A vinyl proton signal of componds 7 appears at δ 6.08 - 6.22 ppm, which is close to that (δ 6.45 ppm) of the analogous type of the compound, 3,5-diphenyl-2*H*-1,4-thiazine.¹⁰ The stereochemistry of the arylimino group at C-2 was determined on the basis of X-ray analysis of 7i (X = Cl, R = Et). ORTEP drawing of 7i shows clearly that the 4-chlorophenyl group is directed toward S-1 (Fig. 1).

The formation of compounds 7 can be explained by a nucleophilic displacement of the bromine atom of 2-bromo-1-phenylethanone by the thione sulfur of compouds 1 to give an intermediate 9a, followed by an intramolecular nucleophilic attack of the imino nitrogen of 9a to the carbonyl carbon atom to give a cyclized intermediate 9b, which loses a water molecule concomitant with loss of a HBr molecule to give 7 (Scheme 1).



Fig. 1. ORTEP drawing of 7i.

Selected bond lengths (Å): S-C4 1.714(4), S-Cl 1.757(4), N1-C1 1.263(5), N1-C15 1.412(5), N2-C2 1.301(5), C3-C4 1.342(5). Selected bond angles (deg): C4-S-C1 103.3(2), C1-N1-C15 120.2(4), C2-N2-C3 125.1(3), C4-C3-N2 123.5(4), C3-C4-S 124.1(3), C2-C1-S 115.6(3).

Table 1. Reaction conditions, yields, and physical properties of compounds 7

Compounds	х	R	1	PhCOCH ₂ Br	Time	Yield *	тр ^ь	Color
			mmol	mmol	h	%	°C	
7 a	NO ₂	<i>n-</i> Pr	0.535	0.819	51	37	79-80	red
7b	Cl	<i>n-</i> Pr	0.588	0.588	40	43	89-9 0	yellow
7c	Me	<i>n</i> -Pr	0.624	0.944	41	62	80-81	yellow
7 d	MeO	<i>n</i> -Pr	0.538	0.819	68	60	79-81	yellow
7e	NO ₂	<i>n</i> -Bu	0.517	0.779	18	51	55-56	red
7 f	Cl	<i>n</i> -Bu	0.491	0.754	46	51	69-71	yellow
7g	Me	<i>n-</i> Bu	0.517	0.794	20	62	64-66	yellow
7h	MeO	<i>n-</i> Bu	0.526	0.809	41	32	74-76	yellow
7i	Cl	Et	0.693	1.04	29	49	129-130	yellow

^a Isolated yields. ^b Recrystallized from EtOH except for 7i (from *n*-hexane).

Table 2. Analytical, ¹H NMR, and IR data of compounds 7

Comp-	Molecular	Analytical	IR (neat)	¹ H NMR (CDCl ₃)
ounds	Formula	Calcd / Found	cm ⁻¹	δ, ppm
7 a	$C_{22}H_{24}N_4O_2S$	C, 64.68 / 64.59	2944, 1570, 1509,	0.60-1.12 (m, 6H, 2CH ₃), 1.55-
		H, 5.92 / 5.99	1478, 1453, 1413,	1.96 (m, 4H, 2CH ₂), 3.51-3.87
		N, 13.71 / 13.63	1366, 1331, 1248,	(m, 4H, 2NCH ₂), 6.22 (s, 1H,
		S, 7.85 / 7.91	1210, 1162, 1102,	SCH=), 6.95 (d, 2H, $J = 8.0$ Hz,
			1074, 1006, 853,	ArH), 7.21-7.45 (m, 3H, ArH),
			715	7.58-7.88 (m, 2H, ArH), 8.27 (d,
				2H, J = 8.0 Hz, ArH)
7b	$C_{22}H_{24}ClN_3S$	C, 66.40 / 66.46	2944, 1571, 1520,	0.60-1.09 (m, 6H, 2CH ₃), 1.48-
		H, 6.08/ 6.15	1474, 1411, 1366,	1.94 (m, 4H, 2CH ₂), 3.55-3.82
		N, 10.56 / 10.48	1326, 1291, 1250,	(m, 4H, 2NCH ₂), 6.15 (s, 1H,
		S, 8.06 / 8.17	1202, 1157, 1085,	SCH=), 6.78 (d, 2H, J = 8.0 Hz,
			1010, 714	ArH), 7.22-7.48 (m, 3H, ArH),
				7.33 (d, 2H, $J = 8.0$ Hz, ArH),
				7.64-7.82 (m, 2H, ArH)
7c	$C_{23}H_{27}N_3S$	C, 73.17 / 73.09	2944, 1571, 1523,	0.74-1.06 (m, 6H, 2CH ₃), 1.40-
		H, 7.21 / 7.24	1493, 1451, 1411,	1.92 (m, 4H, 2CH ₂), 2.26 (s, 3H,
		N, 11.13 / 11.09	1366, 1326, 1293,	CH ₃), 3.45-3.84 (m, 4H,
		S, 8.49 / 8.58	1248, 1094, 1067,	2NCH ₂), 6.08 (s, 1H, SCH=),
			1029, 1003, 915,	6.76 (d, 2H, $J = 8.0$ Hz, ArH),
			770, 714, 688	7.03-7.35 (m, 3H, ArH), 7.18 (d,
				2H, J = 8.0 Hz, ArH), 7.53-7.85
				(m, 2H, ArH)
7d	C23H27N3OS	C, 70.20 / 70.10	2944, 1570, 1523,	0.78-1.08 (m, 6H, 2CH ₃), 1.55-
		H, 6.91 / 6.94	1491, 1451, 1410,	1.84 (m, 4H, 2CH ₂), 3.52-3.87
		N, 10.68 / 10.71	1366, 1326, 1285,	(m, 4H, 2NCH ₂), 3.80 (s, 3H,
		S, 8.15 / 8.21	1240, 1094, 1066,	CH ₃ O), 6.10 (s, 1H, SCH=),
			1027, 1006, 914,	6.89 (s, 4H, ArH), 7.17-7.46 (m,
			830, 770, 718,	3H, ArH), 7.56-7.85 (m, 2H,
			688	ArH)
7e	$C_{24}H_{28}N_4O_2S$	C, 66.03 / 66.09	2928, 1570, 1510,	0.77-1.06 (m, 6H, 2CH ₃), 1.06-
		H, 6.46 / 6.50	1478, 1448, 1413,	1.88 (m, 8H, 2CH ₂ CH ₂), 3.55-
		N, 12.83 / 12.76	1365, 1328, 1230,	3.86 (m, 4H, 2NCH ₂), 6.20 (s,
		S, 7.34 / 7.42	1160, 1101, 101 8 ,	1H, SCH=), 6.92 (d, 2H, $J = 8.0$
			998, 944, 853,	Hz, ArH), 7.18-7.44 (m, 3H,
			712, 693	ArH), 7.56-7.89 (m, 2H, ArH),
				8.29 (d, 2H, $J = 8.0$ Hz, ArH)
7f	$C_{24}H_{28}CIN_3S$	C, 67.66 / 67.73	2936, 1571, 1523,	0.79-1.09 (m, 6H, 2CH ₃), 1.09-

		H, 6.62 / 6.58	1474, 1451, 1413,	1.82 (m, 8H, 2CH ₂ CH ₂), 3.48-
		N, 9.86/9.81	1365, 1326, 1288,	3.87 (m, 4H, 2NCH ₂), 6.12 (s,
		S, 7.53 / 7.64	1227, 1083, 1018,	1H, SCH=), 6.75 (d, 2H, $J = 8.0$
			946, 853, 829,	Hz, ArH), 7.05-7.40 (m, 3H,
			768, 722, 646	ArH), 7.32 (d, 2H, $J = 8.0$ Hz,
				ArH), 7.55-7.85 (m, 2H, ArH)
7 g	$C_{25}H_{31}N_3S$	C, 74.03 / 73.96	2944, 1571, 1523,	0.70-1.09 (m, 6H, 2CH ₃), 1.09-
		H, 7.70/7.75	1494, 1450, 1413,	1.89 (m, 8H, 2CH ₂ CH ₂), 2.35 (s,
		N, 10.36 / 10.33	1365, 1328, 1288,	3H, CH ₃), 3.50-3.85 (m, 4H,
		S, 7.90 / 7.85	1277, 1101, 1016,	2NCH ₂), 6.11 (s, 1H, SCH=),
			944, 858, 819,	6.78 (d, 2H, $J = 8.0$ Hz, ArH),
			770, 714, 688	7.02-7.45 (m, 3H, ArH), 7.24 (d,
				2H, $J = 8.0$ Hz, ArH), 7.58-7.85
				(m, 2H, ArH)
7h	$C_{25}H_{31}N_3OS$	C, 71.22 / 71.13	2920, 1570, 1523,	0.78-1.06 (m, 6H, 2CH ₃), 1.06-
		H, 7.41 / 7.47	1491, 1450, 1411,	1.86 (m, 8H, 2CH ₂ CH ₂), 3.51-
		N, 9.97/9.92	1366, 1285, 1242,	3.91 (m, 4H, 2NCH ₂), 3.84 (s,
		S, 7.60 / 7.71	1178, 1160, 1099,	3H, CH ₃ O), 6.10 (s, 1H, SCH=),
			1077, 1026, 998,	6.88 (s, 4H, ArH), 7.10-7.78 (m,
			966, 944, 857,	3H, ArH), 7.61-7.85 (m, 2H,
			830, 770, 720,	ArH)
			688 , 6 30	
7i	$C_{20}H_{20}ClN_3S$	C, 64 .94 / 64.70	2952, 2912, 1563,	1.28 (t, 6H, $J = 8.0$ Hz, 2CH ₃),
		H, 5.45 / 5.52	1518, 1472, 1418,	3.75 (q, 4H, $J = 8.0$ Hz, 2NCH ₂),
		N, 11.36 / 11.39	1365, 1346, 1326,	6.14 (s, 1H, SCH=), 6.81 (d, 2H,
		S, 8.67/8.75	1274, 1080, 1014,	J = 8.0 Hz, ArH), 7.20-7.45 (m,
			834, 717, 642	3H, ArH), 7.35 (d, 2H, $J = 8.0$
				Hz, ArH), 7.64-7.83 (m, 2H,
				ArH)

For the chemical reactivities of compounds 7, compoud 7b was treated with 20 % agueous hydrochloric acid in THF for 24 h at room temperature. From the reaction mixture were isolated 2-(4-chlorophenylimino)-5-phenyl-4H-1,4-thiazin-3-one (10) and [S-(benzoylmethyl)(4-chlorophenylimino)]methyl-N,N-(di-*n*-propyl)amidine (11) in 53 % and 18 % yields, respectively. The result indicates that 7b undergoes hydrolysis via two pathways: A nucleophilic attack of water to an imino carbon bearing a di-*n*-propylamino group of 7b may result in the formation of compound 10. On the other hand, protonation of an olefinic carbon, followed by a nucleophilic attack of water would lead to compound 11, which is identical with deprotonated compound 9a (X = Cl, R = n-Pr) (Scheme 2).

Reduction of compound 7b with LiAlH₄ in THF at either room or reflux temperature did not proceed and 7b was recovered in 87 % yield. On the other hand, treatment of compound 7b with *m*-chloroperbenzoic acid (*m*-CPBA) in CH₂Cl₂ at room temperature gave a complex mixture from which 7b (27%) was isolable.



Reactions with α -monobromoacyl bromides

The reactions of compounds 1 with bromoacetyl bromide in the presence of pyridine (2 equiv.) at 0 °C gave a mixture showing two spots on TLC ($R_f = 0.90-0.95$ and 0.10-0.50, *n*-hexane : EtOAc = 1 : 1), which were separated by column chromatography. The former was unidenifiable complex mixture and the latter was identified as 5-(di-*n*-alkylamino)-6-arylimino-2*H*-1,4-thiazin-3-ones (12). Reaction conditions, yields, and melting points of compounds 12 are summarized in Table 3 and their analytical and spetroscopic data in Table 4.



The structures of compouds 12 were determined on the basis of the spectroscopic data and elemental analyses. The stereochemistry of arylimino group of 12 was determined based on X-ray analysis of 12i, which showed that the arylimino group was directed toward S-1 (Fig. 2). The possible formation of regioisomer 13 was eliminated in view of the chemical shift of the methylene protons appeared in the range of δ 3.53-3.59 ppm. The chemical shift of the corresponding proton NMR signal of the isomer 13 would be expected to

appear at further downfield because of the presence of two electron-withdrawing groups (a carbonyl group and an imino nitrogen atom) directly bonded to the methylene carbon.¹¹ However, even crude compounds isolated from chromatography did not show absorptions of methylene protons below the range foregoing up to around δ 6.5 ppm. Furthermore, according to the principle of hard and soft acids and bases (HSAB principle) concept,¹² it is reasonable to expect the bond formations between the imino nitrogen (hard base) and the acyl carbon atom (hard acid), and the thione sulfur (soft base) and the carbon atom (soft acid) of the bromomethyl group.

Compounds	Х	R	1	BrCOCH ₂ Br	Time	Yield *	mp ^b
-			mmol	mmol	min	%	°C
12a	NO ₂	<i>n-</i> Pr	0.511	0.57	95	62	129-130
12b	Cl	<i>п</i> -Рг	0.630	0.69	90	75	104-107
12c	Me	<i>n</i> -Pr	0.609	1.09	80	41	86-89
12d	MeO	<i>n</i> -Pr	0.518	0.57	105	47	109-110
12e	NO ₂	<i>n-</i> Bu	0.482	0.57	110	66	140-141
12f	Cl	<i>n</i> -Bu	0.490	0.57	105	84	109-110
12g	Me	<i>n</i> -Bu	0.540	0.57	85	59	99-100
12h	MeO	<i>n-</i> Bu	0.514	0.57	115	50	101-103
1 2 i	NO ₂	Et	0.31	0.34	130	47	180-183 (dec.)

Table 3. Reaction conditions, yields, and melting points of compounds 12

^a Isolated yields. ^b Recrystallized from a mixture of *n*-hexane and dichloromethane except for 12f (from *n*-hexane) and 12i (from ethyl acetate).

Table 4. Analytical, ¹H NMR, and IR data of compounds 12

Comp-	Molecular	Analytical	IR (KBr)	¹ H NMR (CDCl ₃)
ounds	Formula	Calcd / Found	cm ⁻¹	δ, ppm
12a	C ₁₆ H ₂₀ N ₄ O ₃ S	C, 55.16 / 55.09	2936, 1651, 1574,	0.60-1.14 (m, 6H, 2CH ₃), 1.32-2.03
		H, 5.79/ 5.84	1541, 1509, 1456,	(m, 4H, 2CH ₂), 3.13-3.90 (m, 4H,
		N, 16.08 / 16.14	1426, 1334, 1045	2NCH ₂), 3.59 (s, 2H, CH ₂), 6.99 (d,
		S, 9.20 / 9.29		2H, J = 8.0 Hz, ArH), 8.25 (d, 2H,
				J = 8.0 Hz, ArH
12b	C16H20CIN3OS	C, 56.88 / 56.94	2944, 1648, 1595,	0.60-1.14 (m, 6H, 2CH ₃), 1.37-2.03
		H, 5.97/ 6.00	1539, 1477, 1469,	(m, 4H, 2CH ₂), 3.12-3.89 (m, 4H,
		N, 12.44 / 12.41	1434, 1370, 1320,	2NCH ₂), 3.57 (s, 2H, CH ₂), 6.95 (d,
		S, 9.49/9.54	1286, 1083, 1053,	2H, J = 8.0 Hz, ArH), 7.35 (d, 2H,
			920	J = 8.0 Hz, ArH
12c	C17H23N3OS	C, 64.32 / 64.27	2944, 1662, 1589,	0.69-1.12 (m, 6H, 2CH ₃), 1.54-1.94

		H, 7.30 / 7.32	1533, 1496, 1454,	(m, 4H, 2CH ₂), 2.35 (s, 3H, CH ₃),
		N, 13.24 / 13.27	1426, 1365, 1309,	3.45-3.81 (m, 4H, 2NCH ₂), 3.56 (s,
		S, 10.10 / 10.17	1283, 1250, 1043	2H, CH ₂), 6.84 (d, 2H, $J = 8.0$ Hz,
				ArH), 7.22 (d, 2H, $J = 8.0$ Hz,
				ArH)
12d	$C_{17}H_{23}N_3O_2S$	C, 61.24 / 61.29	2944, 1653, 1595,	0.60-1.15 (m, 6H, 2CH ₃), 1.36-2.01
		H, 6.95 / 6.98	1573, 1530, 1496,	(m, 4H, 2CH ₂), 3.17-3.92 (m, 4H,
		N, 12.60 / 12.64	1458, 1371, 1320,	2NCH ₂), 3.56 (s, 2H, CH ₂), 3.80 (s,
		S, 9.61 / 9.67	1288, 1248, 1048,	3H, CH ₃ O). 6.92 (s, 4H, ArH)
			1029	
12e	$C_{18}H_{24}N_4O_3S$	C, 57.43 / 57.46	2936, 1662, 1600,	0.60-1.10 (m, 6H, 2CH ₃), 1.10-2.03
		H, 6.43 / 6.47	1586, 1552, 1504,	(m, 8H, 2CH ₂), 3.20-3.90 (m, 4H,
		N, 14.88 / 14.91	1458, 1430, 1341,	2NCH ₂), 3.57 (s, 2H, CH ₂), 6.97 (d,
		S, 8.52 / 8.58	1320, 1275, 1221,	2H, J = 8.0 Hz, ArH), 8.26 (d, 2H,
			1160, 1099, 1056,	J = 8.0 Hz, ArH)
			862, 699	
12f	C ₁₈ H ₂₄ ClN ₃ OS	C, 59.08 / 59.13	2944, 1650, 1598,	0.60-1.08 (m, 6H, 2CH ₃), 1.08-2.20
		H, 6.61 / 6.59	1542, 1472, 1430,	(m, 8H, 2CH ₂), 3.19-3.90 (m, 4H,
		N, 11.48 / 11.50	1370, 1320, 1277,	2NCH ₂), 3.56 (s, 2H, CH ₂), 6.83 (d,
		S, 8.76/ 8.80	1054	2H, $J = 8.0$ Hz, ArH), 7.35 (d, 2H,
				J = 8.0 Hz, ArH)
12g	C19H27N3OS	C, 66.05 / 66.11	2936, 1672, 1595,	0.60-1.10 (m, 6H, 2CH ₃), 1.10-2.00
		H, 7.88 / 7.90	1555, 1494, 1453,	(m, 8H, 2CH ₂), 2.34 (s, 3H, CH ₃),
		N, 12.16 / 12.19	1432, 1366, 1314,	3.14-3.93 (m, 4H, 2NCH ₂), 3.53 (s,
		S, 9.28 / 9.36	1274, 1043	2H, CH ₂), 6.80 (d, 2H, $J = 8.0$ Hz,
				ArH), 7.19 (d, 2H, $J = 8.0$ Hz,
				ArH)
1 2h	$C_{19}H_{27}N_3O_2S$	C, 63.13 / 63.19	2944, 1651, 1576,	0.60-1.07 (m, 6H, 2CH ₃), 1.07-1.98
		H, 7.53 / 7.50	1526, 1494, 1456,	(m, 8H, 2CH ₂), 3.18-3.91 (m, 4H,
		N, 11.62 / 11.60	1371, 1318, 1283,	2NCH ₂), 3.55 (s, 2H, CH ₂), 3.78 (s,
		S, 8.87/ 8.95	1250, 1050	3H, CH ₃ O), 6.90 (s, 4H, ArH)
12i	$C_{14}H_{16}N_4O_3S$	C, 52.49 / 52.56	3088, 2968, 2920,	1.31 (t, 3H, $J = 7.1$ Hz, CH ₃), 1.37
		H, 5.03 / 5.06	1651, 1598, 1574,	(t, 3H, $J = 7.1$ Hz, CH ₃), 3.62 (s,
		N, 17.49 / 17.44	1536, 1504, 1466,	2H, CH ₂), 3.72 (q, 2H, $J = 7.1$ Hz,
		S , 10.01 / 10.10	1438, 1333, 1309,	NCH ₂), 3.77 (q, $2H$, $J = 7.1$ Hz,
			1096, 1042, 981	NCH ₂), 7.03 (d, 2H, $J = 8.9$ Hz,
				ArH), 8.30 (d, 2H, $J = 8.9$ Hz,
				ArH)

For comparison, 1c (X = Me, R = *n*-Pr) was treated with K_2CO_3 (2 equiv.) in THF at 0°C for 150 min. However, 12c was obtained in 28 % yield together with the recovery of the starting material 1c (35 %). The same reaction carried out at room temperature for 24 h gave 12c (12 %) together with an unknown complex mixture.



Fig. 2. ORTEP drawing of 12i.

Selected bond lengths (Å): S-C1 1.748(3), S-C4 1.809(4), C1-C2 1.503(5), C3-C4 1.495(5), N2-C2 1.303(4), N2-C3 1.366(5), N1-C1 1.273(4). Selected bond angles (deg): C1-S-C4 100.6(2), C1-N1-C9 122.2(3), C2-N2-C3 121.8(3), N2-C3-C4 116.8(3), C3-C4-S 111.5(2), N1-C1-C2 119.6(3).

In contrast with the reaction of 7b, the reduction of compound 12b with LiAlH₄ in THF at room temperature gave 2-(4-chlorophenylimino)-5-oxo-3-(di-*n*-propylamino)thiomorpholine (14) in 87 % yield. Treatment of 12b with P_4S_{10} in benzene gave the corresponding 5-thioxothiazine 15 in 46 % yield. However, only 4-chloroaniline and a complex mixture were isolated by treatment of 12b with 20 % aqueous hydrochloric acid in THF at room temperature. Treatmement of 12b with *m*-CPBA gave a complex mixture from which the starting 12b was recovered in 51 % yield.



The reactions of 1 (R = n-Pr) with 2-bromopropionyl bromide in the presence of pyridine (2 equiv.) in CH₂Cl₂ at 0 °C gave 6-arylimino-2-methyl-5-(di-*n*-propylamino)-2*H*-1,4-thiazin-3-ones (16) as major products and 5-(arylimino)-2-(1-bromoethylidene)-4-(di-*n*-propylamino)thiazolines (17) as minor products. Reaction conditions, yields, and melting points of compounds 16 and 17 are summarized in Table 5. Analytical and spectroscopic data of compounds 16 and 17 are summarized in Table 6 and 7, respectively.



The formation of compounds 16 can be explained by the same mechanism as proposed for compounds 12. We prefer the pathway in which a bromine atom of acyl bromide is displaced by the imino nitrogen of 1 to give an intermediate 18a, which loses HBr to give an intermediate 18b (HSAB principle, vide supra). Nucleophilic displacement of a bromine atom of 18b by the thione sulfur (path a) concomitant with loss of HBr can give compounds 16. However, when the thione sulfur attacks the carbonyl carbon atom in the presence of acid catalyst (HBr) (path b), the formation of bromohydrin 18c would be expected. Dehydration of 18c in the presence of HBr can give rise to a cation 18d, which undergoes deprotonation reaction to give a mixture of (Z)- and (E)-17 (Scheme 3).



Table 5. Reaction conditions, yields, and melting points of compounds 16 and 17

Yield * %
21 (76:24) °
12 (69:31) °
12 (75:25) °

16d	0.709	0.75	90	48	109-110	17d	0

^a Isolated yields. ^b Recrystallized from a mixture of *n*-hexane and dichloromethane. ^c The numbers in the parenthesis represent the ratio of the geometrical isomers, calculated on the basis of the integral values of methyl protons of the ethylidene group.

Table 6. Analytical, ¹H NMR, and IR data of compounds 16

Comp-	Molecular	Analytical	IR (neat)	¹ H NMR (CDCl ₃)
ounds	Formula	Calcd / Found	cm ⁻¹	δ, ppm
16 a	C ₁₇ H ₂₂ N ₄ O ₃ S	C, 62.22 / 62.27	2952, 1670, 1576,	0.60-1.10 (m, 6H, 2CH ₃), 1.45 (d,
		H, 7.25 / 7.23	1542, 1509, 1456,	3H, $J = 8.0$ Hz, CH ₃), 1.40-2.00
		N, 12.09 / 12.06	1432, 1366, 1336,	(m, 4H, 2CH ₂), 3.20-4.00 (m, 4H,
		S, 9.23 / 9.29	1275, 1230, 1160,	2NCH ₂), 3.90 (q, 1H, $J = 8.0$ Hz,
			1099, 1050, 997,	CH), 7.00 (d, 2H, $J = 8.0$ Hz,
			853, 734, 694	ArH), 8.27 (d, 2H, $J = 8.0$ Hz,
				ArH).
16b	$C_{17}H_{22}CIN_3OS$	C, 65.22 / 65.18	2952, 1669, 1595,	0.60-1.10 (m, 6H, 2CH ₃), 1.45 (d,
		H, 7.60 / 7.63	1541, 1470, 1430,	3H, $J = 8.0$ Hz, CH ₃), 1.40-1.95
		N, 12.68 / 12.72	1366, 1318, 1270,	(m, 4H, 2CH ₂), 3.20-3.90 (m, 4H,
		S, 9.67/9.74	1230, 1197, 1157,	2NCH ₂), 3.85 (q, 1H, $J = 8.0$ Hz,
			1085, 1046, 830,	CH), 6.85 (d, 2H, $J = 8.0$ Hz,
			739	ArH), 7.30 (d, 2H, $J = 8.0$
				Hz,ArH).
16c	$C_{18}H_{25}N_3OS$	C, 58.03 / 58.07	2952, 1669, 1590,	0.60-1.20 (m, 6H, 2CH ₃), 1.45 (d
		H, 6.30 / 6.32	1539, 1496, 1453,	$3H, J = 8.0 Hz, CH_3$, 1.40-1.95
		N, 11.94 / 11.99	1429, 1368, 1318,	(m, 4H, 2CH ₂), 2.35 (s, 3H, CH ₃),
		S, 9.11/ 9.18	1274, 1230, 1147,	3.20-3.90 (m, 4H, 2NCH ₂), 3.80
			1045, 1026, 998,	(q, 1H, $J = 8.0$ Hz, CH), 6.80 (d,
			915, 899, 867,	2H, J = 8.0 Hz, ArH), 7.15 (d,
			818, 758, 718	2H, J = 8.0 Hz, ArH).
16d	$C_{18}H_{25}N_3O_2S$	C, 56.34 / 56.30	2952, 1666, 1579,	0.60-1.20 (m, 6H, 2CH ₃), 1.45 (d
		H, 6.12/6.15	1539, 1493, 1450,	$3H, J = 8.0 Hz, CH_3), 1.50-1.95$
		N, 15.46 / 15.50	1366, 1320, 1285,	(m, 4H, 2CH ₂), 3.20-3.85 (m, 4H,
		S, 8.85 / 8.92	1246, 1157, 1098,	2NCH ₂), 3.80 (s, 3H, CH ₃ O), 3.82
			1045, 1026, 995,	(d, 1H, $J = 8.0$ Hz, CH), 6.95 (s,
			910, 894, 832,	4H, ArH) .
			758, 696, 598	

Comp-	Molecular	Analytical	IR (neat)	¹ H NMR (CDCl ₃)
ounds	Formula	Calcd / Found	cm ⁻ '	δ, ppm
17 a	C17H21BrN4O2S	C, 48.01 / 48.07	2952, 1603, 1574,	0.60-1.10 (m, 6H, 2CH ₃), 1.40-
		H, 4.98 / 4.95	1549, 1506, 1451,	1.92 (m, 4H, 2CH ₂), 2.31, 2.60
		N, 13.17 / 13.20	1333, 1261, 1224,	(major) (2s, 3H, CH ₃), 3.40-
		S, 7.54 / 7.60	1162, 1099, 1077,	3.95 (m, 4H, 2NCH ₂), 7.10 (d,
			1042, 1022, 915,	2H, $J = 8.0$ Hz, ArH), 8.27 (d,
			890, 830, 738,	2H, J = 8.0 Hz, ArH).
			699	
17b	C17H21BrClN3S	C, 49.23 / 49.28	2952, 1598, 1574,	0.60-1.10 (m, 6H, 2CH ₃), 1.35-
		Н, 5.10/ 5.13	1549, 1474, 1451,	2.00 (m, 4H, 2CH ₂), 2.32, 2.56
		N, 10.13 / 10.16	1424, 1370, 1222,	(major) (2s, 3H, CH ₃), 3.40-
		S, 7.73 / 7.79	1085, 1042, 1026,	3.90 (m, 4H, 2NCH ₂), 6.95 (d,
			1010, 827	2H, $J = 8.0$ Hz, ArH), 7.35 (d,
				2H, J = 8.0 Hz, ArH
17c	$C_{18}H_{24}BrN_3S$	C, 54.82 / 54.87	2952, 1605, 1586,	0.70-1.10 (m, 6H, 2CH ₃), 1.36-
		H, 6.13 / 6.10	1547, 1494, 1450,	2.00 (m, 4H, 2CH ₂), 2.32, 2.56
		N, 10.65 / 10.63	1424, 1370, 1221,	(major) (2s, 6H, 2CH ₃), 3.50-
		S, 8.13 / 8.20	1096, 1075, 1042,	3.90 (m, 4H, 2NCH ₂), 6.92 (d,
			1024, 818	2H, $J = 8.0$ Hz, ArH), 7.20 (d,
				2H, J = 8.0 Hz, ArH).

Table 7. Analytical, ¹H NMR, and IR data of compounds 17

In the meantime, the reaction of 1a - d with ethyl bromoacetate in CH₂Cl₂ at room temperature gave [(arylimino)(S-ethoxycarbonylmethyl)]methyl-N,N-(di-n-propyl)amidine hydrobromides 19 in good yields. Reaction conditions, yields, and melting points of compounds 19 are summarized in Table 8 and their analytical and spectroscopic data in Table 9.



Attempted purification of compouds 19 by the repeated column chromatography failed but it was successful by employing HPLC using CH_3CN as a solvent.

The structures of compounds 19 were determined on the basis of the spectroscopic data and elemental analyses. It is evident for the amidines to be protonated on the imino nitrogen atom rather than on the amino

nitrogen atom because imino nitrogens are generally more basic than amino nitrogens in amidines.¹³ ¹H NMR spetra of compounds 19 exhibited a broad singlet at δ 4.86 to 6.17 ppm, corresponding to the two protons of the protonated imines.

Table 8. Reaction conditions and yields of compounds 19

Compounds	1	BrCH ₂ CO ₂ Et	Time	Yield *
	mmol	mmol	h	%
19 a	0.334	0.34	24	84
19b	0.447	0.46	1.5	82
19c	0.411	0.42	1	88
19d	0.321	0.34	0.5	71

* Isolated yields.

Table 9. Analytical, ¹H NMR, and IR data of compounds 19

Comp-	Molecular	Analytical	IR (neat)	¹ H NMR (CDCl ₃)
ounds	Formula	Calcd / Found	cm ⁻¹	δ, ppm
19a	C ₁₈ H ₂₇ BrN4O4S	C, 45.48 / 45.55	3312, 2968, 1734,	0.56-0.95 (m, 6H, 2CH ₃), 1.12-
		Н, 5.72/ 5.69	1582, 1510, 1480,	1.85 (m, 7H, 2CH ₂ + CH ₃),
		N, 11.79 / 11.84	1458, 1376, 1338,	2.76-3.41 (m, 4H, 2NCH ₂), 3.85
		S, 6.74/ 6.81	1294, 1262, 1213,	(s, 2H, COCH ₂), 4.22 (q, 2H, J
			1171, 1110, 1050,	= 8.0 Hz, OCH ₂), 4.86 (s, br,
			1026, 859	$2H_{2} = NH_{2}, 6.97 (d, 2H, J = 8.0)$
				Hz, ArH), 8.16 (d, 2H, $J = 8.0$
				Hz, ArH)
19b	C18H27BrClN3O2S	C, 46.51 / 46.47	3296, 2952, 1739,	0.59-1.12 (m, 6H, 2CH ₃), 1.12-
		H, 5.85 / 5.88	1605, 1579, 1478,	2.00 (m, 7H, 2CH ₂ + CH ₃),
		N, 9.04/ 9.09	1370, 1288, 1202,	2.76-3.41 (m, 4H, 2NCH ₂), 3.85
		S, 6.90/6.98	1179, 1150, 1117,	(s, 2H, COCH ₂), 4.20 (q, 2H, J
			1085, 1050, 1026,	= 8.0 Hz, OCH ₂), 5.80 (s, br,
			830	$2H_{2} = NH_{2}, 6.87 (d, 2H, J = 8.0)$
				Hz, ArH), 7.25 (d, 2H, $J = 8.0$
				Hz, ArH)
19c	$C_{19}H_{30}BrN_3O_2S$	C, 51.35 / 51.39	3296, 2952, 1731,	0.55-1.18 (m, 6H, 2CH ₃), 1.18-
		H, 6.80/ 6.83	1648, 1606, 1578,	1.95 (m, 7H, 2CH ₂ + CH ₃), 2.25
		N, 9.45/ 9.42	1474, 1451, 1370,	(s, 3H, CH ₃), 2.76-3.41 (m, 4H,
		S, 7.21 / 7.26	1288, 1149, 1112,	2NCH ₂), 3.82 (s, 2H, COCH ₂),

			1024, 818	4.20 (q, 2H, $J = 8.0$ Hz, OCH ₂), 6.17 (s, br, 2H, = ⁺ NH ₂), 6.83 (d,
				2H, $J = 8.0$ Hz, ArH), 7.05 (d,
				2H, J = 8.0 Hz, ArH)
19d	$C_{19}H_{30}BrN_3O_3S$	C, 49.56 / 49.51	3296, 2952, 1734,	0.58-1.09 (m, 6H, 2CH ₃), 1.09-
		H, 6.57/6.59	1603, 1576, 1496,	1.99 (m, 7H, 2CH ₂ + CH ₃),
		N, 9.13/ 9.16	1454, 1368, 1285,	2.76-3.41 (m, 4H, 2NCH ₂), 3.76
		S, 6.96 / 6.75	1240, 1027, 830	(s, 3H, CH ₃ O), 3.87 (s, 2H,
				COCH ₂), 4.20 (q, 2H, $J = 8.0$
				Hz, OCH ₂), 5.08 (s, br, 2H,
				= ⁺ NH₂), 6.68-6.97 (m, 4H, ArH)

EXPERIMENTAL

N,N-(di-*n*-alkyl)-*N*'-Arylthiocarbamoylamidines were prepared by the literature method.² 2-Bromo-1phenylethanone, 3-chloro-2-propanone, bromoacetyl bromide, 2-bromopropionyl bromide, and ethyl bromoacetate were obtained from Aldrich Chemical Co. Inc.. Thin layer chromatography was carried out on Merck Chromatogram sheet (Kiesel gel 60 F254). Chromatogram was visualized by a mineral UV lamp. Column chromatography was performed using silica gel (Merck, 70-230 mesh) unless otherwise specified. ¹H NMR spetra were obtained with a Brucker AC-80 at 80 MHz, using tetramethylsilane as an internal standard. Infrared (IR) spetra were obtained using a Shimazu IR-470. Mass spectra (MS) were obtained by a VG 12-250 mass spectrometer at 70 eV. HPLC was performed with a C-18 column (μ Bondpak C18, 10 μ m, 7.8 × 300 mm i.d.) and a differential refractometer, using CH₃CN as eluent (flow rate = 0.8 ml / min). Melting points were determined on a Fisher-Johns melting point apparatus and are uncorrected.

General Procedure for the Synthesis of 3-(Di-*n*-alkylamino)-2-arylimino-5-phenyl-2H-1,4-thiazines (7).

To a soluton of N,N-(di-*n*-alkyl)-N'-arylthiocarbamoylamidines 1 (0.49-0.62 mmol) in THF (70 ml) was added K₂CO₃ (0.99-1.27 mmol). A solution of 2-bromo-1-phenylethanone (0.49-0.62 mmol) in THF (50 ml) was dropwise added to the mixture at reflux. Heating was continued for an appropriate time and the reaction mixture was quenched by adding water (50 ml). Evaporation of THF under reduced pressure, followed by extraction with CHCl₃ (3 × 20 ml) gave a residue, which was chromatographed on a silica gel column (2 × 13 cm). Elution with a mixture of *n*-hexane and CH₂Cl₂ (1 : 1) gave compounds 7, unreacted 2-bromo-1phenylethanone and a complex mixture. Consult Table 1 for reaction conditions, yields, and physical properties and Table 2 for the analytical and spectroscopic data of compounds 7.

Hydrolysis of 2-(4-Chlorophenylimino)-5-phenyl-3-(di-*n*-propylamino)-2H-1,4-thiazine (7b) with Aqueous Hydrochloric Acid.

To a solution of 7b (152 mg, 0.382 mmol) in THF (30 ml) was added 20 % aqueous hydrochloric acid (30 ml). The mixture was stirred for 24 h at room temperature. Neutralization of the mixture with aqueous NaOH (5%), followed by evaporation of THF *in vacuo* gave an aqueous solution, which was extracted with

CHCl₃ (3 × 30 ml). The extracts were dried over MgSO₄. Evaporation of the solvent gave a residue, which was chromatographed on a silica gel column (2 × 12 cm). Elution with CH₂Cl₂ gave 2-(4-chlorophenylimino)-5-phenyl-4*H*-1,4-thiazin-3-one (**10**) (64 mg, 53 %): white solid; mp 199-200 °C (*n*-hexane-CH₂Cl₂); IR (KBr) 3336, 3088, 2912, 1670, 1582, 1518, 1478, 1426, 1390, 1301, 1274, 1229, 1168, 1110, 1094, 1066, 1026, 1010, 910, 878, 856, 822, 750, 736, 675, 643, 498, 456, 430 cm⁻¹; ¹H NMR (CDCl₃ + DMSO-d₆) δ 7.35 (d, 2H, *J* = 8.0 Hz, ArH), 7.20-7.62 (m, 3H, ArH), 7.90 (d, 2H, *J* = 8.0 Hz, ArH), 8.00-8.20 (m, 2H, ArH), 8.23 (s, 1H, CH=), 10.47 (s, br, 1H, NH), MS (m/z) 314 (M⁺, 100), 279 (24.5), 272 (15.3), 188 (18.5), 161 (40.9), 102 (84.2). Anal. Calcd for C₁₆H₁₁ClN₂OS: C, 61.05; H, 3.52; N, 8.90; S, 10.18. Found: C, 61.09; H, 3.54; N, 8.87; S, 10.23.

Elution with CH₂Cl₂ gave [*S*-(benzoylmethyl)(4-chlorophenylimino)]methyl-*N*,*N*-(di-*n*-propyl)amidine (11) (28 mg, 18 %): colorless liquid; IR (neat) 3320, 2944, 1696, 1586, 1502, 1445, 1389, 1298, 1085, 1051, 1011, 960, 923, 827, 787, 758, 730, 690 cm⁻¹, ¹H NMR (CDCl₃) δ 0.60-1.05 (m, 6H, 2CH₃), 1.25-1.90 (m, 4H, 2CH₂), 2.32-2.70 (m, 4H, 2NCH₂), 4.44 (s, 2H, CH₂), 7.28 (d, 2H, *J* = 8.0 Hz, ArH), 7.20-7.54 (m, 3H, ArH), 7.55(d, 2H, *J* = 8.0 Hz, ArH), 7.71-7.95 (m, 2H, ArH), 9.04 (s, br, 1H, NH). Anal. Calcd for C₂₂H₂₆ClN₃OS: C, 63.52; H, 6.30; N, 10.10; S, 7.71. Found: C, 63.47; H, 6.32; N, 10.12; S, 7.77.

Elution with CH₂Cl₂ gave an unknown mixture (31 mg).

Reaction of 7b with Lithium Aluminum Hydride (LiAlH4).

To a solution of 7b (144 mg, 0.362 mmol) in THF (30 ml) was added LiAlH₄ (17 mg, 0.45 mmol). The mixture was stirred for 1.5 h at room temperature and then worked up as usual. Only 7b (125 mg, 98 %) was isolable. The reaction did not proceed at reflux temperature.

Reaction of 7b with m-Chloroperbenzoic Acid (m-CPBA).

To a solution of 7b (153 mg, 0.384 mmol) in $CH_2Cl_2(30 \text{ ml})$ was added *m*-CPBA (116 mg, 71.5 %). The color of the solution turned immediately from yellow to dark red. The mixture was stirred for 9 h at room temperature and then washed with aqueous NaHCO₃ solution. The solution was evaporated and the residue was chromatographed on a silica gel column (2 × 15 cm). Elution with a mixture of *n*-hexane and dichloromethane (1 : 1) gave unreacted 7b (41 mg, 27 %) and a complex mixture, which showed many spots on TLC (silica gel, *n*-hexane - CH₂Cl₂).

General Procedure for the Sythesis of 5-(Di-n-alkylamino)-6-arylimino-2H-1,4-thiazin-3-ones (12).

To a solution of a mixture of 1 (0.49-0.63 mmol) and pyridine (0.99-1.5 mmol) in CH_2Cl_2 (50 ml) at 0 °C was added dropwise a solution of bromoacetyl bromide (0.57-1.1 mmol) in CH_2Cl_2 (40 ml) for 65 min to 85 min. The mixture was additionally stirred for an appropriate time, followed by washing with water (5 × 150 ml). Drying over MgSO₄, followed by removal of the solvent in *vacuo* gave a residue, which was chromatographed on a silica gel column (2 × 13 cm). Elution with a mixture of *n*-hexane and EtOAc (1 : 1) gave a mixture of deep blue color. Elution next with the same solvent mixture (1 : 2) gave compounds 12. Consult Table 3 for reaction conditions, yields, and melting points of compounds 12 and Table 4 for their analytical and spectroscopic data.

Reaction of 6-(4-Chlorophenylimino)-5-(di-*n*-propylamino)-2*H*-1,4-thiazin-3-one (12b) with LiAlH₄.

To a solution of **12b** (169 mg, 0.500 mmol) in THF (30 ml) was added a suspension of LiAlH₄ (25 mg, 0.66 mmol) in THF (30 ml) through a dropping funnel for 5 min. The mixture was stirred for 2 h and worked up as usual. Column chromatography (2 × 12 cm) of the mixture using a mixture of *n*-hexane and EtOAc(1 : 1) as an eluent gave 2-(4-chlorophenylimino)-5-oxo-3-(di-*n*-propylamino)thiomorpholine (**14**) (148 mg, 87 %): pale yellow liquid; IR (neat) 3218, 3096, 2952, 2856, 1682, 1616, 1589, 1478, 1466, 1387, 1294, 1181, 1134, 1083, 1038, 1011, 835 cm⁻¹; ¹H NMR (CDCl₃) δ 0.64-1.06 (m, 6H, 2CH₃), 1.32-1.80 (m, 4H, 2CH₂), 2.48-2.90 (m, 4H, 2NCH₃), 3.11 and 4.41 (d, 2H, *J* = 12.0 Hz, CH₂), 4.67 (d, 1H, *J* = 6.0 Hz, CH), 6.79 (d, 2H, *J* = 8.0 Hz, ArH), 7.88 (d, br, 1H, *J* = 6.0 Hz, NH). MS (m/z) 339 (M⁺, 11.7), 240 (65.5), 155 (100). Anal. Calcd for C₁₆H₂₂ClN₃OS: C, 56.54; H, 6.52; N, 12.36; S, 9.43. Found: C, 56.57; H, 6.50; N, 12.39; S, 9.49.

Reaction of 12b with Phosphorus Pentasulfide (P₄S₁₀).

To a solution of **12b** (110 mg, 0.326 mmol) in benzene (30 ml) was added P_4S_{10} (147 mg, 0.331 mmol). The mixture was vigorously stirred for 7 h, followed by washing with water (3 × 50 ml). The organic layer was dried over MgSO₄. Removal of the solvent *in vacuo* gave a residue, which was chromatographed on a silica gel column (3 × 13 cm). Elution with a mixture of *n*-hexane and EtOAc (3 : 1) gave a complex mixture and 6-(4-chlorophenylimino)-5-(di-*n*-propylamino)-2*H*-1,4-thiazine-3-thione (**15**) (53 mg, 46 %): red liquid; IR (neat) 2952, 1586, 1542, 1454, 1427, 1360, 1325, 1248, 1224, 1182, 1142, 1085, 1037, 1022, 1010, 923, 829, 730 cm⁻¹; ¹H NMR (CDCl₃) δ 0.60-1.20 (m, 6H, 2CH₃), 1.40-2.00 (m, 4H, 2CH₂), 3.40-3.86 (m, 4H, 2NCH₂), 3.94 (s, 2H, CH₂), 6.89 (d, 2H, *J* = 8.0 Hz, ArH), 7.36 (d, 2H, *J* = 8.0 Hz, ArH); MS (m/z) 353 (M⁺, 44.1), 323 (39.6), 291 (65.7), 262 (100), 220 (49.6). Anal. Calcd for C₁₆H₂₀ClN₃S₂. C, 54.30; H, 5.70; N, 11.87; S, 18.12. Found: C, 54.27; H, 5.72; N, 11.91; S, 18.17.

Hydrolysis of 12b with Aqueous Hydrochloric Acid

To a solution of 12b (180 mg, 0.533 mmol) in THF (30 ml) was added 20 % hydrochloric acid (30 ml). The mixture was stirred for 24 h, followed by neutralization with aqueous NaOH (5%). The aqueous layer was extracted with CHCl₃ (3×20 ml). The organic extracts were dried over MgSO₄. Removal of the solvent *in vacuo* gave a residue, which was chromatographed on a silica gel column (2×8 cm). Elution with a mixture of *n*-hexane and EtOAc (1 : 1) gave 4-chloroaniline (58 mg, 85 %). Elution with ethanol gave a unknown compound (9 mg) which was insoluble in most of the organic solvents.

General Procedure for the Synthesis of 6-Arylimino-2-methyl-5-(di-*n*-propylamino)-2H-1,4thiazin-3-ones (16) and 5-Arylimino-2-(1-bromoethylidene)4-(di-*n*-propylamino)thiazolines (17).

To a solution of a mixture of 1 (0.54-0.74 mmol) and pyridine (1.2-1.5 mmol) in CH_2Cl_2 (70 ml) at 0° C was added dropwise a solution of 2-bromopropionyl bromide (0.65-0.93 mmol) in CH_2Cl_2 (50 ml) for 1 h. The mixture was additionally stirred for an appropriate time and then washed with water (5 × 50 ml). The mixture was worked up as usual. The reaction mixture was chromatographed on a silica gel column (2 × 15 cm). Elution with a mixture of *n*-hexane and $CH_2Cl_2(4 : 1)$ gave compounds 17 and a complex mixture. Elution with a mixture of n-hexane and EtOAc(1 : 1) gave compounds 16. Consult Table 5 for reaction conditions, yields,

and melting points of compounds 16 and 17. Consult Table 6 and 7 for analytical, and spectroscopic data of compounds 16 and 17, respectively.

General Procedure for the Synthesis of [(Arylimino)(S-ethoxycarbonylmethyl)]methyl-N,N-(di-n-propyl)amidine Hydrobromides (19).

To a solution of 1 (0.32-0.45 mmol) in CH_2Cl_2 (100 ml) was added ethyl bromoacetate (0.34-0.36 mmol). The mixture was stirred for an appropriate time at room temperature. Evaporation of the solvent *in vacuo* gave a crude product. TLC (silica gel) of each product showed one spot with a tail: 19a ($R_f = 0.7 - 0.9$, CH_2Cl_2 : acetone = 2 : 1), 19b ($R_f = 0.4 - 0.6$, CH_2Cl_2 : acetone = 5 : 1), 19c ($R_f = 0.2 - 0.4$, EtOAc : acetone = 1 : 2), 19d ($R_f = 0.2 - 0.4$, EtOAc : acetone = 2 : 7). The crude products were purified by employing HPLC using acetonitrile as an eluent, Consult Table 8 for reaction conditions and yields of compounds 19 and Table 9 for their analytical and spectroscopic data.

Single Crystal X-ray Analyses of 7i and 12i.

The data were collected on an Enraf-Nomius CAD 4 diffractometer using graphite-monochromated Mo-K_{α} radiation. The structures were solved by direct methods and subsequent Fourier maps. Refinements were carried out by full-matrix least-squares techniques. Non-hydrogen atoms were anisotropically refined. Atomic scattering factors were taken from International Tables for X-ray Crystallography, Vol IV, 1974. All calculations and drawings were performed using a Micro VAX II Computer with the SDP system.

	7i	12i
Molecular formula	C20H20ClN3S	C14H16N4O3S
Molecular weight	370.91	320.37
Crystal system	monoclinic	monoclinic
Space group	P2 (1) / n	P2 (1) / c
a, Å	10.2568(12)	8.190(2)
b, Å	13.775(2)	7.586(2)
c, Å	13.679(2)	24.917(8)
a, deg	90.000(13)	90.00(2)
β , deg	98.659(12)	98.23(2)
γ , deg	90.00(2)	90.00(2)
V, Å ³	1910.7(4)	1532.3(7)
Z	4	4
ρ calc., g / cm ³	1.289	1.389
Crystal size, mm	$0.20 \times 0.45 \times 0.55$	$0.25 \times 0.25 \times 0.45$
Absorption coefficient, mm ⁻¹	0.317	0.229
θ range, deg	2.11 - 24.98	1.65 - 23.45
Index ranges	$-12 \le h \le 12, 0 \le k \le 16, 0 \le l \le 16$	$0 \le h \le 9, 0 \le k \le 8, -27 \le l \le 27$
N _b of measured reflections	3504	2070
N_b of reflections used $\{I \ge \sigma(I)\}$	3357	1927
Data to parameter ratio	3356 / 228	1927 / 199
Final R indices	$R_1 = 0.0726$, $wR_2 = 0.2245$	$R_1 = 0.0761$, $wR_2 = 0.1952$
R indices (all data)	$R_1 = 0.1135$, $wR_2 = 0.2417$	$R_1 = 0.0763$, $wR_2 = 0.1957$

Crystal and Refinement Parameters for Compounds 7i and 12i

Atomic coordinates, bond lengths, angles and thermal parameters have been deposited at the Cambridge Crystallographic Data Centre.

Acknowledgements. This paper was financially supported in part by NON DIRECTED RESEARCH FUND, Korea Research Foundation and Center for Biofunctional Molecules (CBM).

REFERENCES AND NOTES

- 1. Lee, H.; Kim, K.; Whang, D.; Kim, K. J. Org. Chem. 1994, 59, 6179-6183.
- 2. Choi, S.-H.; Kim, K. Tetrahedron 1996, 52, 8413-8438.
- (a) Wolfe, S.; Zhang, C.; Johnston, B. D.; Kim, C.-K. Can. J. Chem. 1993, 72, 1066-1075. (b) Wolfe, S.; Stephen, B. PCT Int. Appl. WO 90 07,111, 1990.; Chem. Abstr. 1991, 114, 101534e.
- Takahashi, M.; Yoshioka, K.; Yamazaki, H. PCT Int. Appl. WO 91 18,893, 1991.; Chem. Abstr. 1992, 116, 106323m.
- (a) Stoodley, R. J. in 1,4-Thiazines and Their Dihydro Derivatives, Advances in Heterocyclic Chemistry Vol. 24, Katritzky, A. R.; Boulton, A. J. Eds.; Academic Press: New York, 1979; pp 293-361. (b) Sainsbury, M. in Oxazines, Thiazines and Their Benzo Derivatives, Comprehensive Heterocyclic Chemistry, Vol. 3, Katritzky, A. R.; Rees, C. W. Eds.; Pergamon Press: New York, 1984; Chap. 2.27, pp 995-1038.
- 6. Johnson, C. R.; Sataty, I. J. Med. Chem. 1967, 10, 501-503.
- (a) Reliquet, A.; Besbes, R.; Reliquet, F.; Meslin, J. C. Phosphorus, Sulfur, Silicon Relat. Elem. 1992, 70, 211-217. (b) Reliquet, A.; Besbes, R.; Reliquet, F.; Meslin, J. C. Synthesis, 1991, 543-546.
- Meslin, J. C.; Reliquet, A.; Besbes, R.; Reliquet, F. Fr. Demande FR 2,678,264, 1992; Chem. Abstr. 1993, 119, 28152a.
- 9. Besbes, R.; Reliquet, A.; Reliquet, F.; Meslin, J. C. Phosphorus, Sulfur, Silicon Relat. Elem. 1993, 79, 123-129.
- 10. Foster, R.; Gilchrist, T. L. J. Chem. Soc., Perkin Trans. 1, 1991, 2249-2254.
- 4-Alkyl-2H-1,4-thiazin-3-ones exhibited a singlet at δ 3.28-3.35 ppm, assignable to methylene protons: Hojo, M.; Masuda, R.; Kosaka, S.; Nagase, K. Synthesis, 1979, 272.
- 12. Lowry. T. H.; Richardson, K. S. in *Mechanism and Theory in Organic Chemistry*; 3rd ed. Harper & Row: New York, 1987; Chap. 3, pp 318-322.
- 13. Hafelinger, G. in *The Chemistry of Amidines and Imidates*; Patai, S. Ed.; John Wiley and Sons, Inc.: New York, 1975; Chap. 1, pp 11-18.