Scheme I

changes of ketone solutes. It is obvious that many other reactions and processes are amenable to study in these media.

Acknowledgment. We thank the National Science Foundation (Grant CHE83-01776) for financial support.

Hypervalent Silicon Hydrides: SiH5

David J. Hajdasz and Robert R. Squires*

Department of Chemistry, Purdue University West Lafayette, Indiana 47907

Received February 24, 1986

The rich diversity and practical utility of organosilicon chemistry continues to engender widespread research interests.¹ A key issue in the field involves the role of the low-lying vacant Si d-orbitals in the properties and reactivity of silicon-containing compounds. A familiar manifestation of d-orbital participation is the aptitude of silicon for hypervalency in the form of (10-Si-5) pentavalent anions.² Examples of pentavalent silicon anions have been isolated and structurally characterized,³ and they are frequently invoked as intermediates in nucleophilic addition reactions of organosilanes in solution.⁴⁻⁷ Moreover, several 10-Si-5 negative ions bearing alkyl, alkoxyl, and halogen ligands have been identified as products of gas-phase ion-molecule reactions.⁸⁻¹⁰ We now wish to report the synthesis and characterization of the archetype pentavalent silicon anion, $SiH_5^{-}(1)$, along with several of its simple alkyl derivatives in a flowing afterglow apparatus at 298 K.¹¹

Hydride ion is a potent gas-phase base $(PA = 400.4 \text{ kcal/mol})^{12}$ which can be generated by electron impact on either NH₃ or CH₄. Its direct reaction with SiH₄ in a flow reactor at 0.4 torr has been previously reported by Bohme and co-workers to yield exclusively SiH₃⁻ by proton transfer.¹³ Interestingly, when D⁻ is the reactant

references cited therein.

(3) Stevenson, W. H.; Wilson, S.; Martin, J. C.; Farnham, W. B. J. Am. Chem. Soc. 1985, 107, 6340 and references cited therein. (4) Stevenson, W. H.; Martin, J. C. J. Am. Chem. Soc. 1985, 107, 6352

and reference cited therein.

(5) Sommer, L. H. Stereochemistry, Mechanism and Silicon; McGraw-Hill: New York, 1965.

(6) Corriu, R. J. P.; Dabosi, G.; Martineau, M. J. Organomet. Chem. 1980, 186, 25

(7) Klanberg, F.; Muetterties, E. L. Inorg. Chem. 1968, 7, 155.
(8) (a) Klass, G.; Trenerry, C.; Sheldon, J. C.; Bowie, J. H. Aust. J. Chem.
1981, 34, 519. (b) Sheldon, J. C.; Bowie, J. H.; Hayes, R. N. Nouv, J. Chim.
1984, 8, 79. (c) Sheldon, J. C.; Hayes, R. N.; Bowie, J. H. J. Am. Chem. Soc.

1984, 8, 79. (c) Sheldon, J. C.; Hayes, R. N.; Bowie, J. H. J. Am. Chem. Soc.
1984, 106, 7711. (d) Hayes, R. N.; Bowie, J. H.; Klass, G. J. Chem. Soc., Perkin Trans 2 1984, 1167.
(9) Murphy, M. K.; Beauchamp, J. L. J. Am. Chem. Soc. 1977, 99, 4992.
(10) (a) DePuy, C. H.; Bierbaum, V. M.; Flippin, L. A.; Grabowski, J. J.; Schmitt, R. J.; Sullivan, S. A. J. Am. Chem. Soc. 1980, 102, 5012. (b) Sullivan, S. A.; DePuy, C. H.; Damrauer, R. J. Am. Chem. Soc. 1981, 103, 480. (c) Squires, R. R.; DePuy, C. H. Org. Mass Spectrom. 1982, 17, 187.
(d) Damrauer, R.; DePuy, C. H.; Bierbaum, V. M. Organometallics 1982, 1, 1553. (e) DePuy, C. H.; Damrauer, R. J. Am. Chem. Soc. 1984, 3, 362. (f) DePuy, C. H.; Bierbaum, V. M.; Damrauer, R. J. Am. Chem. Soc. 1984, 106, 4051. 4051

(11) The apparatus is fully described in: Lane, K. R.; Lee, R. E.; Sallans, L.; Squires, R. R. J. Am. Chem. Soc. 1984, 106, 5767. Unless otherwise noted the standard experimental conditions are P(He) = 0.400 torr; F(He) = 190

the standard experimental conditions are P(He) = 0.400 (or; P(He) = 190atm cm³ s⁻¹; $\overline{\nu}$ (He) = 9400 cm s⁻¹; $T = 298 \pm 2$ K. (12) Bartmess, J. E.; McIver, R. T., Jr. In *Gas Phase Ion Chemistry*; Bowers, M. T. Ed.; Vol. 2.; Academic Press: New York, 1979, Chapter 11. (13) Payzant, J. D.; Tanaka, K.; Betowski, L. D.; Bohme, D. K. J. Am. Chem. Soc. 1976, 98, 894.



ion, H⁻ and SiH₂D⁻ also appear as products, implying that reversible addition of the nucleophilic ion to silicon occurs during the course of reactive collisions. We have now found that H^- reacts with alkylsilanes in a flowing afterglow at 0.4 torr total pressure by exclusive addition (eq 1). These reactions are most likely

$$H^- + R_4 Si \xrightarrow{He} R_4 Si H^-$$
 (1)

$$R_4Si = Me_4Si$$
, Et_3SiH , Et_2SiH_2 , $n-C_5H_{11}SiH_3$

saturated termolecular processes as shown by their large bimolecular rate coefficients at 0.4 torr (e.g., 4.5×10^{-9} cm³ s⁻¹ for Et₃SiH) and their pressure-independent product distributions from 0.3 to 0.9 torr. Relative to SiH₄, termolecular addition is facilitated in the larger alkylsilanes by the greater complexity and, thus, longer lifetimes of the intermediate collision complexes.¹⁴ Adduct formation appears to be unique to hydride ion since other strongly basic anions such as NH_2^- (PA = 403.6 kcal/mol)¹² and OH^- (PA = 390.7 kcal/mol)¹² react only by proton abstraction and condensation reactions.¹⁰

The alkylsilyl hydride anions formed by reaction 1 behave as potent hydride donors, reducing a wide variety of substrates in bimolecular hydride transfer reactions such as CO₂, Et₃B, and Fe(CO)₅.¹⁵ Of special significance is the observation of facile hydride transfer among the silanes themselves. In particular, both $n-C_5H_{11}SiH_4^{-}(2)$ and $Et_3SiH_2^{-}(3)$ transfer hydride to SiH₄ to produce the parent ion of the series, SiH_5^- (eq 2). The major

$$Et_{3}SiH_{2}^{-} + SiH_{4} \xrightarrow{68\%} SiH_{3}^{-} + Et_{3}SiH + H_{2}$$

$$\frac{m/z \ 31}{32\%} SiH_{5}^{-} + Et_{3}SiH$$

$$\frac{m/z \ 33}{1}$$
(2)

primary product with $Et_3SiH_2^-$ is SiH_3^- (68%), which is also observed to build up as a secondary product at the expense of SiH5 at higher SiH_4 flow rates. This latter observation indicates that an 'autocatalyzed' decomposition reaction is occurring (eq 3). The

$$\operatorname{SiH}_5^- + \operatorname{SiH}_4 \to \operatorname{SiH}_4 + \operatorname{H}_2 + \operatorname{SiH}_3^- \tag{3}$$

occurrence of such a reaction in the gas phase requires that SiH₅⁻ be thermodynamically unstable with respect to H_2 dissociation but may survive intact under the present conditions by virtue of a kinetic barrier preventing prompt fragmentation. Computational studies of SiH₅⁻ support this conclusion since it has been determined at quite high levels of theory that although H₂ dissociation is ca. 6 kcal/mol exothermic, a D_{3h} trigonal-bipyramidal SiH₅⁻ structure represents a local minimum on the potential energy surface.16-19

Reaction 3 is but one example of a more general protolytic reaction which may occur between silyl hydride anions and Brønsted acids (HX) (eq 4). We have exploited this reaction

$$R_4 SiH^- + HX \rightarrow [R_4 Si \cdots H \cdots H \cdots X] \rightarrow R_4 Si + H_2 + X^-$$
(4)

in the determination of thermochemical data for SiH₅⁻ and two of its alkyl derivatives. Equation 4 may be formulated in terms

⁽¹⁾ For reviews, see: (a) Fessenden, R. J.; Fessenden, J. S. Adv. Orga-nomet. Chem. 1980, 18, 275. (b) Fleming, I. Compr. Org. Chem. 1979, 3, 542. (c) Colvin, E. W. Silicon in Organic Synthesis; Butterworths: London, 1981. (d) West, R. C.; Barton, T. J. J. Chem. Educ. 1980, 57, 334. (e) Barton, T. J. Pure Appl. Chem. 1980, 52, 615. (f) Gladysz, J. A. Acc. Chem. Res. 1984, 17, 326. (g) Schaefer, H. F. Acc. Chem. Res. 1982, 15, 283. (h) Drahnak, T. J.; Michl, J. West, R. J. Am. Chem. Soc. 1981, 103, 1845. (2) Corriu, R. J. P.; Guerin, C. Adv. Organomet. Chem. 1982, 20, 265 and references cited therein

^{(14) (}a) Bohme, D. K.; Dunkin, D. B.; Fehsenfeld, F. C.; Ferguson, E. E. J. Chem. Phys. 1969, 51, 863. (b) Durden, D. A.; Kebarle, P.; Good, A. J.
 Chem. Phys. 1969, 50, 805. (c) Meot-Ner, M. In Gas Phase Ion Chemistry;
 Bowers, M. T. Ed.; Academic Press: New York 1979, Chapter 6.
 (15) Lane, K. R.; Sallans, L.; Squires, R. R. Organometallics 1985, 3, 408.

Table I. Protonolysis Reactions of Silyl Hydride Anions, 298 K

	reference acids ^a					
anion	EtOH (376.1)	MeOH (381.4)	H ₂ O (390.7)	D ₂ O (392.9)	NH ₃ (403.6)	inference ^b
$SiH_{5}^{-}(1)$	+	+	_	_		$18.0 \le HA \le 26.9$
$n-C_{5}H_{11}SiH_{4}^{-}(2)$	+	+	+	+	-	$5.3 \le HA \le 16.3$
$Et_3SiH_2^-(3)$	+	+	+	+	-	$4.7 \le \text{HA} \le 15.6$

 ${}^{a}\Delta H_{acid}(HX)$ shown in parentheses in kcal/mol.¹² A + sign indicates occurrence of protonolysis and formation of the corresponding conjugate base nion. b Estimated limits on the hydride affinity (HA) of the corresponding neutral silane in kcal/mol. Correction factors which take into account anion. changes in rotational and translational entropy have been incorporated.

of the thermochemical cycle in Scheme I, where $HA(R_4Si)$ is the hydride affinity of the silane $D[R_4Si-H^-]$ and $\Delta H_{acid}(HX)$ and $\Delta H_{acid}(H_2)$ are known quantities.¹² Thus, by monitoring the occurrence or nonoccurrence of reaction 4 for a particular silyl hydride and a series of acids of varying acidity, one can derive limits on the hydride affinity of the silane. Table I presents the results of such "bracketing" experiments for three reactant ions. Our experimentally derived limits on the hydride binding energies are in good agreement with the computed values for SiH₄ reported by Baybutt (18.6 kcal/mol, HF),¹⁶ Wilhite and Spialter (16.9 kcal/mol, HF),¹⁷ Keil and Ahlrichs (20.3 kcal/mol, PNO CI),¹⁸ and most recently by Schleyer (18.3 kcal/mol, MP4/ 6-31++G**).¹⁹ Hydride affinities of this magnitude are among the lowest known for any neutral molecule and are comparable to those of NH₃ (8 kcal/mol),²⁰ H₂O (16 kcal/mol),²¹ and benzene (25 kcal/mol).22

The facility with which the silyl hydride ions undergo binary hydride transfer reactions combined with their ready adaptability to isotope labeling makes them ideal systems for investigating the structures and fluxional behavior of isolated 10-Si-5 negative ions. Small Berry pseudorotational barriers have been computed for SiH₅⁻ (ca. 2.7 kcal/mol)^{17,18} and measured experimentally for 10-Si-5 phenyl siliconates in solution.^{4,23} Therefore, one may anticipate that the hydrogens in 1-3 undergo rapid intramolecular scrambling once the ions are formed. To test this, we have generated in situ monodeuterated alkylsilyl hydride ions using D⁻ (from ND_3) in conjunction with reaction 1. Addition of CO_2 to the system as a convenient hydride acceptor is followed by determination of the primary relative yields of HCO₂⁻ and DCO₂⁻ products. In the absence of a kinetic isotope effect for transfer of H⁻ vs. D⁻, chemical equivalence of the hydrogens in the reactant ions would be indicated by statistical yields of HCO₂⁻ and DCO₂⁻. The results for ions 2-d and 3-d are shown below (eq 5 and 6),

$$\operatorname{Et_3SiH} \xrightarrow{D^-} \operatorname{Et_3SiHD^-} \xrightarrow{\operatorname{CO_2}} \operatorname{HCO_2^-}, \operatorname{DCO_2^-} (5)$$

$$3 \cdot d \xrightarrow{68\%} 32\%$$

$$n - C_5 H_{11} \text{SiH}_3 \xrightarrow{D^-} n - C_5 H_{11} \text{SiH}_3 D^- \xrightarrow{CO_2} HCO_2^-, DCO_2^- \quad (6)$$

$$2 - d \xrightarrow{2 - d} 86\% \xrightarrow{1 - 4\%} HCO_2^-, DCO_2^- \quad (6)$$

where the indicated branching ratios have been corrected for the natural isotope distributions in each ion and the contribution to the HCO_2^- yield from the unavoidable presence of unlabeled reactant ion.²⁴ An isotope effect is immediately evident since the determined yields deviate significantly from the 1:1 and 3:1 statistical values. However, the deviation is a constant $k_{\rm H}/k_{\rm D}$ = 2.1 for the two systems, so we may conclude that complete

- (16) Baybutt, P. Mol. Phys. 1975, 29, 389.
 (17) Wilhite, D. L.; Spialter, L. J. Am. Chem. Soc. 1973, 95, 2100.
- (18) Keil, F.; Ahlrichs, R. Chem. Phys. 1975, 8, 384.

- (18) Kell, F.; Anirichs, K. Chem. Phys. 1975, 8, 384.
 (19) Schleyer, P. v. R., private communication.
 (20) (a) Kleingeld, J. C.; Ingemann, S.; Jalonen, J. E.; Nibbering, N. M. M. J. Am. Chem. Soc. 1983, 105, 2474. (b) Kalcher, J.; Rosmus, P. Can. J. Phys. 1984, 62, 1323. (c) Cremer, D.; Kraka, E. J. Phys. Chem. 1986, 90, 33. (d) Coe, J. V.; Snodgrass, J. T.; Freidhoff, C. B.; McHugh, K. M.; Bowen, K. H. J. Chem. Phys., in press.
 (21) (a) Kleingeld, J. C.; Nibbering, N. M. M. Int. J. Mass Spectrom. Ion Phys. 1983, 49, 311. (b) Paulson, J. F.; Henchman, M. J. Bull. Am. Phys. Soc. 1982, 27, 108. Soc. 1982, 27, 108.
 - (22) Lane, K. R.; Sallans, L.; Squires, R. R. J. Am. Chem. Soc., in press.

 - (23) Damrauer, R.; Danahey, S. E. Organometallics, in press.
 (24) Ions 2-d and 3-d undergo secondary H/D exchange reactions with the
- excess neutral silane precursor in the flow tube eventually losing the label.

scrambling of H and D has occurred in the silicon ions prior to the probe reaction with CO₂. Further experiments with $Et_2SiH_2D^-$, SiH_4D^- , and other isotopomers are in progress to confirm this finding.

Acknowledgment. We are grateful to Professor P. v. R. Schleyer for disclosure of his computational results for SiH5⁻ in advance of publication. Support from the National Science Foundation (Grant CHE-8502515) is gratefully acknowledged.

Proton Transfer as a Function of Hydration Number and Temperature: Rate Constants and Product Distributions for $OH^{-}(H_2O)_{0.1,2,3}$ + HF at 200-500 K

Peter M. Hierl,*1,2 Anton F. Ahrens,^{2,3}

Michael Henchman,*4 A. A. Viggiano,⁵ and John F. Paulson Air Force Geophysics Laboratory Hanscom AFB, Massachusetts 01731

D. C. Clary

University Chemical Laboratory Cambridge CB2 1EW, U.K. Received January 2, 1986

Proton-transfer reactions which are exothermic in the gas phase are exceedingly efficient-most reacting at essentially every collision.⁶⁻¹⁰ Several studies have shown how this extreme reactivity is influenced by solvation in the gas phase at 300 K.^{8,9,11-16} Here we report how hydration influences the rate constants and the product distributions of the reaction

$$OH^- + HF = H_2O + F^- \qquad \Delta H^\circ = -19.5 \text{ kcal/mol}^{17}$$
 (1)

throughout the temperature range 200-500 K. Such studies in the gas phase-where the solvation number is systematically changed—examine the role of solvate in the transfer of protons.¹⁸

- (1) SCEEE Fellow/AFOSR Summer Faculty Research Associate, 1984. (2) Permanent address: Department of Chemistry, University of Kansas,
- Lawrence, KS 66045.
- (3) SCEEE Fellow/AFOSR Summer Graduate Research Associate, 1984.
 (4) AFSC-URRP Visiting Professor, 1984-1986. Permanent address: Department of Chemistry, Brandeis University, Waltham, MA 02254.
- (5) System Integration Engineering Inc., Lexington, MA 02173: under contract to AFGL
- (6) Bohme, D. K. In Interactions between Ions and Molecules; Ausloos,
 P., Ed.; Plenum: New York, 1975; p 489.
 (7) Bohme, D. K.; Mackay, G. I.; Schiff, H. I. J. Chem. Phys. 1980, 73,
- 4976.
- (8) Bohme, D. K. Trans. R. Soc. Can., Sect. 3 1981, 19, 265.
 - (9) Bohme, D. K. In Ionic Processes in the Gas Phase; Almoster Ferreira,
- M. A., Ed.; Reidel: Dordrecht, 1984; p 111. (10) Moylan, C. R.; Brauman, J. I. Annu. Rev. Phys. Chem. 1983, 34, 187. (11) Bohme, D. K.; Mackay, G. I.; Tanner, S. D. J. Am. Chem. Soc. 1979,
- 101, 3724. (12) Smith, D.; Adams, N. G.; Henchman, M. J. J. Chem. Phys. 1980, 72, 4951.
- (13) Tanner, S. D.; Mackay, G. I.; Bohme, D. K. Can. J. Chem. 1981, 59,
- (14) Bohme, D. K.; Rakshit, A. B.; Mackay, G. I. J. Am. Chem. Soc. 1982,
- (1) John (100.
 (15) Mackay, G. I.; Rakshit, A. B.; Bohme, D. K. Can. J. Chem. 1982, 60, 2594.
- (16) Raksit, A. B.; Bohme, D. K. Can. J. Chem. 1983, 61, 1683.

(17) Bartmess, J. E.; Molver, R. T., Jr. In *Gas Phase Ion Chemistry*;
Bowers, M. T., Ed.; Academic Press: New York, 1979; Vol. 2, p 90. The value is unaltered by the redetermination of E. A. [OH]. Schulz, P. A.; Mead, R. D.; Jones, P. L.; Lineberger, W. C. J. Chem. Phys. 1982, 77, 1153.