Preparation, Identification, and Reactivity of New Organogold(I) Halide Anion Complexes

O. M. ABU-SALAH* and A. R. AL-OHALY

Department of Chemistry, College of Science, King Saud University, P.O. Box No. 2455, Riyadh, Saudi Arabia

Received April 11, 1983

A previous work showed that copper(I) and silver(I) arylacetylides are fruitful sources of interesting organocopper and silver complexes [1]. In contrast to copper(I) and silver(I), gold(I) complexes have a great tendency to exhibit linear geometry [2]. In gold(I) phenylacetylide $[AuC_2Ph]_{\eta}$, linearity could be achieved by polymerisation through π -bonding between alkyne groups and gold atoms [3]. Treatment of this complex with tertiary phosphines and other neutral ligands gives monomeric acetylides such as Et_3PAuC_2Ph [3].

We report in this letter the reaction of halide ions, Cl⁻, Br⁻, or I⁻ with gold phenylacetylide. The normally insoluble gold complex dissolved within five minutes when added (1:1) to a solution of halide ions in acetone or dichloromethane. White crystals of phenylethynyl gold(I) halide anion complexes [NR₄][XAuC₂Ph] (X = Cl or Br, R = Et; X = I, R = Buⁿ) were obtained in ca. 90% yield on addition of hexane to the above solutions. Elemental analyses and ¹H n.m.r. spectroscopy indicate a 1:1 adduct. Molecular weight measurements, obtained by using a vaporimetric method, for the chloride and iodide complexes gave values characteristic of 1:1 electrolyte (Table I). Therefore, these complexes are monomeric with linear geometry.

TABLE I. Molecular Weight Data.

Complex	Measured M (acetone)	Calculated
[NEt ₄][ClAuC ₂ Ph]	290	463.5
[NBu ⁿ 4][IAuC ₂ Ph]	380	667

The i.r. spectra of these complexes contained a medium to a strong sharp absorption band at ca. 2120 cm⁻¹ assigned for acetylide group. The $\nu(C\equiv C)$ band obtained does not change by altering the type of the halide ion in the complex, and is very close to $\nu(C\equiv C)$ absorption bands in the neutral complexes LAuC₂Ph (L = tertiary phosphines, phosphites, arsines, etc.) [3]. This finding, with anionic complexes,

confirms the availability of poor back-bonding from gold to π^* orbitals of the alkyne group described previously [3].

These complexes are stable and can be handled in the air in the solid state. After several weeks a slight change in colour of the salt is observed, probably due to slight sensitivity to light; this change, however, does not affect the other physical properties. Addition of water to acetone solution of the organogold(I) chloride or bromide regenerates the yellow gold phenylacetylide. Recently organogold(I) complexes $[PPh_4][RC_2AuC_2R]$ (R = H, Me, Ph) were reported [4] to be stable in contrast to their analogous potassium salts (R = H, Ph) [5]. The organogold(I) complexes $[NR_4][XAuC_6F_5]$ (X = Cl, R = Et; X = Br, $R = Bu^n$) were prepared by the reaction of $[NR_4][X]$ with $[thtAuC_6F_5]$ (tht = tetrahydrothiophen) in the presence of the corresponding concentrated HX [6, 7]. Nevertheless, the same method [7] failed to give the analogous organogold(I) iodide anion complex [IAuC₆F₅]⁻. Lately, the complex anion [IAu- C_6F_5] was prepared by the reaction of QI{(Q = NBu₄,(PPh₃)₂N,Ph₃BzP} and thtAuC₆F₅ in ethanol or dichloromethane [8, 9].

When organogold(I) chloride and bromide complexes were treated with [NEt₄][I], the iodide ions displaced both chloride and bromide ions. The major product was the corresponding linear organogold(I) iodide, as indicated by analytical data and i.r. spectroscopy. A minor product of an organogold(I) iodide complex with different stoichiometry was also obtained. The latter is currently being studied to determine its structure. The reaction of the complex [Cl-AuC₂Ph] with PPh₃ gave PPh₃AuC₂Ph in very high yield. On the other hand, the reaction of the complex [IAuC₂Ph] with PPh₃ gave different results. The first crop was PPh₃AuI (34% yield) and the second product was identified as [(PPh₃)₂N][PhC₂AuC₂Ph] (43% yield) [4]. The latter reaction represents a novel method for the preparation of the organogold complex [PhC₂AuC₂Ph]⁻.

Acknowledgements

We thank Mr. M. K. Mahmoud and Mr. M. Ja'far for technical assistance.

References

- 1 (a) O. M. Abu-Salah, M. I. Bruce, S. A. Bezman and M. R. Churchill, J. Chem. Soc., Chem. Commun., 858 (1972).
 - (b) O. M. Abu-Salah and M. I. Bruce, J. Chem. Soc., Dalton Trans., 2311 (1975).

^{*}Author to whom correspondence should be addressed.

- (c) O. M. Abu-Salah, M. I. Bruce, M. R. Churchill and B. G. DeBoer, J. Chem. Soc., Chem. Commun., 688 (1974).
- 2 R. J. Puddephatt, 'The Chemistry of Gold', Elsevier, 11 (1978) and references therein.
- 3 G. E. Coates and C. Parkin, J. Chem. Soc., 3220 (1962).
- 4 R. Nast, P. Schneller and A. Hengefeld, J. Organometal. Chem., 214, 273 (1981).
- 5 R. Nast and U. Kirner, Z. Anorg. Allg. Chem., 330,
- 311 (1964).
- 6 R. Uson, A. Laguna and J. Vicente, J. Chem. Soc., Chem. Commun., 353 (1976).
- 7 R. Uson, A. Laguna and J. Vicente, J. Organometal. Chem., 131, 471 (1977).
- 8 R. Uson, A. Laguna, J. Garcia and M. Laguna, *Inorg. Chim. Acta*, 37(2), 201 (1979).
- 9 R. Uson, A. Laguna, M. Laguna and V. Perez, Synth. React. Inorg. Met.-Org. Chem., 11(4), 361 (1981).