

Use of Thallium(I) Ethoxide in Suzuki Cross Coupling Reactions


Scott A. Frank, Hou Chen, Roxanne K. Kunz, Matthew J. Schnaderbeck, and William R. Roush*

Department of Chemistry, University of Michigan, Ann Arbor, Michigan 48109

roush@umich.edu

Received June 22, 2000

ABSTRACT



Thallium(I) ethoxide promotes Suzuki cross couplings for a range of vinyl- and arylboronic acids with vinyl and aryl halide partners in good to excellent yields. This reagent offers distinct advantages over thallium(I) hydroxide in terms of commercial availability, stability, and ease of use.

The Suzuki reaction is an important method for the synthesis of conjugated olefins, styrenes and biphenyls,^{1–3} and is extremely useful in the context of complex natural products synthesis.^{4–8} In the original version of the reaction,^{2,9} a vinyl (or aryl) halide is combined with a vinyl (or aryl) boronic acid (or ester) or dialkylborane in the presence of bases such as NaOMe or K_2CO_3 and a $\text{Pd}(0)$ catalyst in an organic solvent at temperatures between 60 and 90 °C for 2–24 h. In 1987 Kishi reported remarkable rate accelerations when the Suzuki reaction was performed using TIOH as the base, in which case the cross couplings were complete within minutes at ambient temperature.⁴ We have found the TIOH method to be superior to many other procedures especially

when 1,1-dibromoolefins or 1-iodo-1-trimethylsilylalkenes, which easily undergo base-induced elimination of HBr or HI , are used as the substrates.^{7,10–12}

Despite its great utility in the Suzuki reaction, aqueous solutions of TIOH are both air and light sensitive and have an inferior shelf life. Moreover, commercial sources of the reagent have virtually disappeared, and to our knowledge only a single supplier of solid TIOH currently exists.¹³ Dissolution of solid TIOH in water gives an initially clear solution from which a brown-black precipitate separates even under careful storage conditions.¹⁴ Moreover, we have

(1) Miyaura, N.; Yamada, K.; Sugimoto, H.; Suzuki, A. *J. Am. Chem. Soc.* **1985**, *107*, 972.

(2) Miyaura, N.; Suzuki, A. *Chem. Rev.* **1995**, *95*, 2457.

(3) Suzuki, A. *J. Organomet. Chem.* **1999**, *576*, 147.

(4) Uenishi, J.-i.; Beau, J.-M.; Armstrong, R. W.; Kishi, Y. *J. Am. Chem. Soc.* **1987**, *109*, 4756.

(5) Nicolaou, K. C.; Ramphal, J. Y.; Palazon, J. M.; Spanevello, R. A. *Angew. Chem., Int. Ed. Engl.* **1989**, *28*, 587.

(6) Evans, D. A.; Ng, H. P.; Rieger, D. L. *J. Am. Chem. Soc.* **1993**, *115*, 11446.

(7) Roush, W. R.; Sciotti, R. J. *J. Am. Chem. Soc.* **1998**, *120*, 7411.

(8) Scheidt, K. A.; Tasaka, A.; Bannister, T. D.; Wendt, M. D.; Roush, W. R. *Angew. Chem., Int. Ed.* **1999**, *38*, 1652.

(9) Miyaura, N.; Yamada, K.; Suzuki, A. *Tetrahedron Lett.* **1979**, 3437.

(10) Roush, W. R.; Moriarty, K. J.; Brown, B. B. *Tetrahedron Lett.* **1990**, *31*, 6509.

(11) Roush, W. R.; Kageyama, M.; Riva, R.; Brown, B. B.; Warmus, J. S.; Moriarty, K. J. *J. Org. Chem.* **1991**, *56*, 1192.

(12) Roush, W. R.; Brown, B. B. *J. Am. Chem. Soc.* **1993**, *115*, 2268.

(13) While 10% aqueous solutions of TIOH have been available in the past from a variety of commercial vendors, a recent search revealed only a single supplier of solid TIOH . A conversation with a Johnson-Matthey customer service representative indicated that 10% TIOH solutions exhibited both air and light sensitivity and that they had ceased marketing the reagent for reasons of shelf life. While TIOEt is significantly more stable than TIOH , it is air and moisture sensitive and should be handled under inert atmospheres.


(14) The identity of the brown-black precipitate is unclear, but a likely suspect is Ti_2O_3 : Tóth, I.; Györi, B. *Encyclopedia of Inorganic Chemistry*; King, B. R., Ed.; John Wiley and Sons: New York, 1994; Vol. 8, p 4134.

observed that the efficiency of Suzuki reactions promoted by TIOH is dependent on the age of the TIOH solution, with yields decreasing substantially with older batches of reagent. For example, during studies focusing on completion of a total synthesis of kijanolide,¹⁵ the Suzuki coupling of **1** and **2**, using a 0.5 M stock solution of TIOH that had been prepared 1 month earlier and which had been employed successfully in our total synthesis of bafilomycin A,⁸ provided **3** in 71% yield. However, when we repeated this reaction 4 months later using the same TIOH solution (after removing the brown-black precipitate by filtration), the yield of **3** fell to 50%. The next time this experiment was performed, an additional 6 months later, we prepared a fresh 0.5 M solution of TIOH from the same batch of solid TIOH, again with filtration to remove insoluble dark material, but the yield of **3** was still only 52%. The latter observations suggest that TIOH may also be somewhat unstable in the solid state. Thus, we turned our attention to identification of a suitable replacement for TIOH.

Although thallium(I) ethoxide (TIOEt) is an easily handled liquid that is readily available from several commercial suppliers at lower cost than TIOH, we are aware of only one published account in which this reagent has been used successfully as the base in a Suzuki cross coupling reaction.¹⁶ However, there are several examples where TIOEt failed to give acceptable results,^{17,18} including Kishi's original report.⁴ In the later work, Kishi indicated that TIOEt gave only a 5-fold rate enhancement compared to experiments performed using KOH, while TIOH increased the rate a 1000-fold.

In a key initial experiment (Figure 1), we found that the Suzuki cross coupling of **1** and **2** using TIOEt in an aqueous THF solvent system, using conditions which otherwise mirrored the previous experiments performed using the 10% TIOH solutions, provided **3** in 83% yield. Interestingly, there was no significant difference in the rate of coupling of **1** and **2** using either TIOH or TIOEt.

We have demonstrated that TIOEt may be used with a range of functionalized vinyl and aryl halide and vinyl- and arylboronic acid coupling partners, as summarized in Figure 2.¹⁹ As shown in these examples, vinyl iodides (**4**, **11**), 1,1-dibromo olefins (**7**), α -iodo vinylsilanes (**14**), and aryl iodides (**17**, **19**) are excellent coupling partners, and many potentially sensitive functional groups such as methyl esters, geminally alkylated malonate systems, silyl ethers, enones, etc. are fully compatible with these reaction conditions. In the accompanying paper, Chemler and Danishefsky have demonstrated that TIOEt also gives good results in intramolecular *B*-alkyl Suzuki macrocyclization reactions.²⁰



TIX source	Reagent Age	Yield of 3
TIOH (10% stock solution)	1 month old	71%
TIOH (10% stock solution)	5 months old	50%
TIOH (fresh, from solid)	12 months old	52%
TIOEt	--	83%

Figure 1.

The TIOEt-promoted Suzuki reactions summarized in Figures 1 and 2 are very fast, with TIBr or TII precipitating from solution immediately upon addition of TIOEt to a mixture of the vinyl (or aryl) halide and vinyl (or aryl) boronic acid. Although we have not performed any detailed kinetic measurements, qualitatively these reactions appear to be as fast as Suzuki reactions we have previously performed using TIOH. Moreover, these reactions are substantially faster than those performed using AgO. For example, the coupling of **4** and **5** in the presence of AgO in 3:1 THF–H₂O gave only ca. 25% conversion to **6** after a 30 min reaction period, whereas the reaction of **4** and **5** in the presence of TIOEt is complete almost immediately after mixing of all reagents. Kishi reported that AgO is second only to TIOH in terms of rate acceleration (30-fold) of the Suzuki reaction;⁴ however, our results suggest that TIOEt rivals TIOH in this capacity.

We initiated these studies on the assumption that TIOH is an obligatory intermediate⁴ and that TIOEt would be rapidly converted to TIOH under the aqueous reaction conditions. To determine if it is necessary to run the TIOEt-promoted Suzuki cross couplings in the presence of water, the reaction of **4** and **5** was performed in anhydrous THF (Figure 2). Tetraene **6** was obtained in 92% yield, compared to 97% for the original experiment in the 3:1 THF–H₂O solvent system, with no noticeable difference in reaction rate. A better yield of **18** was obtained from the coupling of phenyl

(15) Roush, W. R.; Brown, B. B. *J. Org. Chem.* **1993**, 58, 2162, and references therein.

(16) Humphrey, J. M.; Aggen, J. B.; Chamberlin, A. R. *J. Am. Chem. Soc.* **1996**, 118, 11759.

(17) Sato, M.; Miyaura, N.; Suzuki, A. *Chem. Lett.* **1989**, 1405.

(18) Markó, I. E.; Murphy, F.; Dolan, S. *Tetrahedron Lett.* **1996**, 37, 2507.

(19) All new compounds were fully characterized by ¹H and ¹³C NMR, IR, and high-resolution mass spectroscopy.

(20) Chemler, S. R.; Danishefsky, S. J. *Org. Lett.* **2000**, 2, 2695. We thank the Sloan-Kettering group for sharing their results with us prior to publication.




Figure 2. Representative TIOEt-promoted Suzuki reactions. Unless otherwise noted, all reactions were performed using TIOEt in 3:1 THF–H₂O using the conditions described in text for the coupling of **5** and **7**.

iodide and vinylboronic acid **15** in wet THF (94%) than in anhydrous THF (83%), although the former result was obtained using an inverse addition protocol (addition of a mixture of **17** and $\text{Pd(PPh}_3)_4$ to a mixture of **15** and TIOEt in wet THF; the yield of **18** was only 73% using the normal addition). However, the aqueous reaction conditions were essential to the successful, high-yield coupling of **19** and **20**.

One variable that is critical to the success of these reactions is the solubility of an intermediate generated from the reaction of the vinyl(aryl)boronic acid and TIOEt. For example, exposure of THF solutions of **15** or **20** to TIOEt in the absence of the other reaction components leads to the immediate precipitation of white solids, which we presume to be the thallium ate complexes of the boronic acids.²¹ The precipitate generated from **15** and TIOEt partially dissolves

when H₂O is added; however the solid analogously generated from **20** does not appear to dissolve at all under these conditions, or in other solvents such as MeOH, CH₃CN, or DMF. Therefore, to maximize the yield of biphenyl **21** from the coupling of **19** and **20**, it was necessary to use 5 equiv of the phenylboronic acid in the aqueous solvent system. Attempts to use arylboronic acids **22** and **23** (Figure 3) in




Figure 3. Unsuccessful reaction partners.

these reactions were unsuccessful, presumably due to the insolubility of the boronic acid–TIOEt reaction product. For this reason, an excess of the boronic acid component was utilized in many of the experiments summarized in Figure 2.²²

(21) Dewar, M. J. S.; Jones, R. *J. Am. Chem. Soc.* **1967**, *89*, 2408.

In summary, we have demonstrated that TIOEt is an excellent alternative to TIOH for use in Suzuki cross coupling reactions. This reagent has advantages compared to TIOH in terms of stability and ease of use. Most importantly, the commercial availability of TIOEt makes it significantly more attractive than TIOH for use in this important cross coupling reaction.

Acknowledgment. Financial support provided by the National Institutes of Health to W.R.R. (GM 26782) and an NIH Postdoctoral Fellowship to H.C. (CA 76655) is gratefully acknowledged.

OL0062446

(22) **Representative experimental procedure:** A solution of **5** (0.42 g, 3.28 mmol) and **7** (0.74 g, 1.09 mmol) in THF and water (12 mL, 3:1) was degassed thoroughly (3×) by using the freeze–pump–thaw method. Tetrakis(triphenylphosphine)palladium(0) (0.13 g, 0.11 mmol) was added as a solid, and the yellow solution was stirred for 5 min at ambient temperature. Thallium ethoxide (0.14 mL, 1.96 mmol) was added by syringe, and a bright yellow precipitate immediately formed. (**NOTE:** *thallium salts are generally quite toxic and should be handled with extreme care.*) The reaction was stirred 30 min and then was diluted with ether (25 mL) and a 1 N sodium bisulfate solution (10 mL). The biphasic mixture was filtered through Celite, and the organic layer was separated, washed with brine, and dried over MgSO₄. Concentration of the filtrate and purification of the crude tetraene by column chromatography afforded 0.60 g (81%) of **8**.