Received: October 15, 1974

PHOTOCHEMICAL REACTIONS OF FLUOROXYPERFLUOROALKANES WITH PERFLUOROCYCLOOLEFINS*

M.S. TOY and R.S. STRINGHAM

Stanford Research Institute, Menlo Park, California 94025 (U.S.A.)

SUMMARY

Under photolysis perfluoro-t-butyl hypofluorite readily undergoes one-electron reduction at -20° C in the presence of perfluorocycloolefin, which functions as a fluoroxy fluorine atom acceptor, to yield bis(perfluorot-butyl)peroxide as the major product. The other products are the simple adducts $C_5F_9OC(CF_3)_3$ and $C_6F_{11}OC(CF_3)_3$ from the corresponding perfluorocycloolefins C_5F_8 and C_6F_{10} , the cleavage compounds of $(CF_3)_3COF$ (i.e., CF_3COCF_3 and CF_4), the corresponding perfluorocycloalkanes, C_5F_{10} and C_6F_{12} , a small amount of vicinal perfluorodi-t-butoxycycloalkanes, and a trace of the dimers and oligomers. The formation of the peroxide $(CF_3)_3COOC(CF_3)_3$ confirms the homolysis of the O-F bond in $(CF_3)_3COF$ under photolysis.

The photochemical reactions of CF_3OF and perfluorocycloolefins at $-60\ ^{o}C$ give mainly the simple adducts $C_5F_9OCF_3$ and $C_6F_{10}OCF_3$ from the corresponding perfluorocycloolefins. The other products are perfluorocyclo-alkanes, dimers, and oligomers, but the peroxide CF_3OOCF_3 and perfluoro-dimethoxycycloalkanes are absent.

^{*} Partially presented at the 168th National American Chemical Society Meeting, Atlantic City, New Jersey, U.S.A., September 1974.

This paper examines the differences between the products of reactions of primary and tertiary fluoroxyperfluoroalkanes with perfluorocycloolefins and discusses the mechanisms of these reactions.

INTRODUCTION

This research represents part of an investigation elucidating the nature of addition reactions of fluoroxy groups to unsaturated perfluorocarbon bonds. The objective is to study the dissociation of the covalent O-F bond by homolytic and heterolytic fissions. Recently we reported that neat $(CF_3)_3COF$ is easily added to hexafluoropropene at $-60^{\circ}C$ to form a simple adduct $n-C_3F_7OC(CF_3)_3$.¹ The n-perfluoropropyl adduct suggests the ionic mechanism, the heterolytic fission of the O-F bond. The unique structural orientation of the product, since it is contrary to the usual nucleophilic fluoride ion adducts (i.e., the perfluoroisopropyl deriva-tives),²,³ suggests electrophilic addition involving fluoronium ion.^{1,4}

Although reactions involving electrophilic addition are not common for perfluoroolefins, Miller and Freedman⁵ reported the electrophilic addition of mercuric fluoride to hexafluoropropene in anhydrous hydrogen fluoride. Such an electrophilic addition mechanism is further supported by Aldrich and coworkers, ⁶ who reported that the addition of mercuric fluoride is limited to terminal perfluoroolefins (because of the increased polarity of the terminal double bond) and that nonterminal perfluoroolefins do not react.

This work was initiated to prepare simple adducts of tertiary fluoroxyperfluoroalkanes and nonterminal perfluoroolefins. It was soon found that $(CF_3)_3COF$ readily decomposed into fragments in the presence of perfluorocyclopentene, while the primary fluoroxyperfluoroalkane (CF₃OF) added to perfluorocyclopentene (C₅F₈) to form a simple adduct (C₅F₉OCF₃) as previously reported by Porter and Cady.⁷

This paper describes the photochemical reactions of primary and tertiary fluoroxyperfluoroalkanes with perfluorocycloolefins.

RESULTS AND DISCUSSION

Under photolysis the reaction of $(CF_3)_3COF$ and perfluorocycloolefin $(C_5F_8 \text{ or } C_6F_{10})$ at $-20^{\circ}C$ gives a mixture of products, which includes the peroxide $(CF_3)_3COOC(CF_3)_3$ as the major product and the simple adducts $C_5F_9OC(CF_3)_3$ or $C_6F_{11}OC(CF_3)_3$, cleavage compounds of $(CF_3)_3COF$ (i.e., CF_3COCF_3 and CF_4), perfluorocycloalkanes C_5F_{10} or C_6F_{12} , a small amount of the vicinal perfluorodi-t-butoxycycloalkanes $C_5F_8[OC(CF_3)_3]_2$ or $C_6F_{10}[OC(CF_3)_3]_2$, and a trace of dimers and oligomers. The yields of the peroxide, hexafluoroacetone, and tetrafluoromethane increase with an increase of $(CF_3)_3COF$, while an increase of perfluorocycloolefin increases the other products.

The proposed reaction sequence described below is shown to be consistent with the observed products from the photochemical reactions of $(CF_3)_3COF$ and perfluorocycloolefins. Some possible reaction paths are presented to correlate and explain the results.

Initiation

$$(CF_3)_3 COF \xrightarrow{NV} (CF_3)_3 CO. + F.$$
 (1)

Reaction (1) initiates a series of reactions in the presence of perfluorocycloolefin, which functions as a fluoroxy fluorine atom acceptor. Reaction (2) is at least partially responsible for the enhancement of homolysis of the O-F bond in $(CF_4)_{3}COF$.

$$C_3F_8$$
 (or C_6F_{10}) + $(CF_3)_3COF \xrightarrow{h_{\vee}} C_5F_9$. (or C_6F_{11} .) + $(CF_3)_3CO$. (2)

A simple combination of the two radical products of reaction (2) accounts for the formation of the simple adduct, perfluoro-t-butoxycyclopentane $C_5F_9OC(CF_3)_3$ and perfluoro-t-butoxycyclohexane $C_6F_{11}OC(CF_3)_3$ (see Tables 1 to 4).

When reactions of $(CF_3)_3COF$ and perfluorocycloolefins were carried out at $0^{\circ}C$ without irradiation, the unimolecular chain decomposition of $(CF_3)_3COF$ occurred. The rapid decomposition is initiated by fluoroxy fluorine atom abstraction from the molecule and is then followed by F-scission of perfluoro-t-butoxy radical.

Propagation and termination

$$(CF_3)_3CO \rightarrow (CF_3)_2CO + CF_3.$$
 (3)

$$(CF_3)_3COF + CF_3$$
. $\rightarrow CF_4 + (CF_3)_3CO$. (4)

The chain reactions (3) and (4) proceed at such a rapid rate that they become almost the sole reactions to account for the cleavage products (i.e., hexafluoroacetone and tetrafluoromethane).

When the same reaction was carried out under photolysis at -20 $^{\circ}$ C, the lower temperature slowed the unimolecular chain decomposition of $(CF_3)_3$ COF. This rapid decomposition of the hypofluorite, which is not observed for

Ч	
NBLE	
È	

s	
ETHER	
OPENTYL	
OROCYCL	
PERFLU	
SOME	
g	
SPECTRA	
NMR	

¹⁹ F NMR SPECTRA OF SOME PERFLUORC	CYCLOPENTYL	ETHERS		
Compound	Relative Intensity	Chemical Shift φ (ppm from CFCl ₃)	Coupling Constants J (Hz)	Assignment
F^{4A} F^{2A}	თ	71.6	J(1,6)=14.9±0.2 J(1,2A)=3.2±0.2 J(1,3B)=3.2±0.2	٦
rad radia a	4	127.5	J(2A,3B)=258±3	2A
		131.3		3B
FSB F3B		134.3		4A
	4		J(4A,5B)=260±3	
		135.1		5B
	1	137.4	J(1,6)=14.9±0.2	9
F ^{4A} F ² A F ^{5B} F ³ B OCF ¹	e	67.6	J(1,6)=10.0±0.2 J(1,2A)=6.0±0.2 J(1,3B)=4.6±0.2	1
		126.8		2A
r4A r2A F6	4		J(2A,3B)=260±3	
		137.1		3B
₽sB _F 3B		131,8		4A
	4		J(4A,5B)=260±3	
		135.3		5B
	I	140.7	Complex multiplet	9

TABLE 1 (concluded)

Compound	Relative Intensity	Chemical Shift ϕ (ppm from CFCl ₃)	Coupling Constants J Hz	Assignment
₽4Å ₩5Å	18	70.9		1A 2B
	2	138.2	Broad	2.4 IR
F^{4B} F^{5B} $OC(CF_{3}^{1A})_{3}$		128.2		3A, 5B
F ^{3A} F ^{2A} F ^{1B}	4		J (3A , 5B) =J (5A , 3B) = 250±3	
F_{3B} 0c(cF_{2}^{2B}),		127.6		5A, 3B
2	Ŋ	130.4		4A, 4B
	18	71.1		lA, 2A
F4A F5A	2	135.7	Broad	IB, 2B
		130.1		3B, 5B
F ^{4B} F ^{3B} oc(cr ₁ ^A) ₃	4		J (3A, 3B) =J (5A, 5B) = 260±3	
$F^{3}A$ oc($cF_{2}^{2}A$)		125.3		3A, 5A
Harris Contraction of the second		133.5		4B
4	2		$J(4A, 4B) = 260 \pm 3$	
		129.3		4A

AND DERIVATIVES
ETHERS
PERFLUOROCYCLOPENTYL
SOME
OF
0006
(LKB
SPECTRA
IASS

TABLE 2

- $285 \, C_{5}F_{11} 0^{+} (2.3), \, 231 \, C_{5}F_{9} 0^{+} (6.0), \, 219 \, C_{1}F_{9}^{+} (2.3), \, 212 \, C_{5}F_{8}^{+} (1.1), \, 181 \, C_{4}F_{7}^{+} (6.0), \, 147 \, C_{4}F_{5} 0^{+}$ (9.0), 131 $C_3F_5^+$ (8.0), 119 $C_2F_5^+$ (4.0), 100 $C_2F_4^+$ (4.0), 97 $C_2F_3O^+$ (15.0), 93 $C_3F_3^+$ (5.0), 69 CF_3^+ $466 \ C_9F_{18}O^+ \ (0.0), \ 447 \ C_9F_{1}O^+ \ (3.9), \ 409 \ C_9F_{15}O^+ \ (0.5), \ 347 \ C_7F_{13}O^+ \ (4.1), \ 316 \ C_8F_{12}O^+ \ (3.6), \ (3.6)$ (100.0), 50 CF⁺₂ (3.0), 31 CF⁺ (3.0). C₅F₉OC(CF₃)₃:
- $682 \ C_{13}F_{26}O_{2}^{+} (0.3), \ 663 \ C_{12}F_{25}O_{2}^{+} (9.6), \ 563 \ C_{11}F_{21}O_{2}^{+} (8.3), \ 447 \ C_{9}F_{17}O^{+} (0.7), \ 435 \ C_{8}F_{17}O^{-} (3.3), \ 682 \ C_{13}F_{26}O_{2}^{+} (0.7), \ 435 \ C_{8}F_{17}O^{-} (3.3), \ 682 \ C_{13}F_{26}O_{2}^{+} (0.7), \ 435 \ C_{8}F_{17}O^{-} (3.3), \ 682 \ C_{13}F_{26}O_{2}^{+} (0.7), \ 435 \ C_{12}F_{17}O^{-} (3.3), \ 682 \ C_{13}F_{26}O_{2}^{+} (0.7), \ 682 \ C_{12}F_{17}O^{-} (0.7), \ 682 \ C_{12}F_{17}O^{-} (0.7), \ 682 \ C_{13}F_{17}O^{-} (0.7), \ 682 \ C_{12}F_{17}O^{-} (0.7), \ 682 \ C_{17}O^{-} (0.7), \ 682 \ C_{12}F_{17}O^{-} (0.7), \ 682 \ C_{17}O^{-} (0.7), \ 682 \ C_{12}F_{17}O^{-} (0.7), \ \ 682 \ C_{12}F_{17}O^{-} (0.7), \ \ 682 \ C_{12}F_{$ 112 $C_{3}F_{4}^{+}$ (1.0), 109 $C_{3}F_{3}O^{+}$ (1.0), 100 $C_{2}F_{4}^{+}$ (4.3), 97 $C_{2}F_{3}O^{+}$ (5.3), 93 $C_{3}F_{3}^{+}$ (1.7), 69 CF_{3}^{+} (100), $c_{3}F_{7}^{+}$ (1.7), 162 $c_{4}F_{6}^{+}$ (1.0), 150 $c_{3}F_{6}^{+}$ (0.7), 147 $c_{3}F_{5}O^{+}$ (2.3), 131 $c_{3}F_{3}^{+}$ (33.0), 119 $c_{2}F_{5}^{+}$ (1.7), $347 C_7 F_{13} 0^+ (20.0), 285 C_5 F_{11} 0^+ (1.0), 231 C_5 F_9^+ (0.7), 219 C_4 F_9^+ (6.0), 181 C_4 F_7^+ (2.0), 169 C_6 F_9^+ (0.0), 169 C_7 F_9^+ (0.0), 160 C_7 F_9^+ (0.0), 180 C_7 F_9^+ (0.0), 160 C_7 F_9^+ (0.0), 180 C_7 F_9^+ (0.0), 160 C_7 F_9^+ (0.0), 160 C_7 F_9^+ (0.0), 160 C_7 F_9^+ (0.0), 160 C_7 F_9^+ (0.0), 180 C_7 F_9^+ (0.0), 160 C_7 F_9^+ (0.0), 180 C_7 F_9^+ (0.0), 180 C_7 F_9^+ (0.0), 160 C_7 F_9^+ (0.0), 160 C_7 F_9^+ (0.0), 180 C_7 F_9^+ (0.$ 50 CF_{9}^{+} (1.0), 47 CFO^{+} (1.3), 31 CF^{+} (1.3), 28 CO^{+} or N_{9}^{+} (1.3). $c_{s}F_{s}[oc(cF_{3})_{3}]_{2}:$
 - $c_{s}F_{7}^{+}$ (3.0), 185 $c_{3}F_{7}O^{+}$ (2.0), 181 $c_{4}F_{7}^{+}$ (54), 162 $c_{4}F_{6}^{+}$ (5.0), 143 $c_{4}F_{5}^{+}$ (2.0), 131 $c_{3}F_{5}^{-}$ (100), $316 C_6 F_{12} O^+ (0.0), 297 C_6 F_{11} O^+ (7.0), 231 C_5 F_9^+ (75), 212 C_5 F_8^+ (2.0), 209 C_5 F_7 O^+ (5.0), 193$ 119 $C_{2}F_{5}^{+}$ (2.0), 100 $C_{2}F_{4}^{+}$ (14), 93 $C_{3}F_{3}^{+}$ (6.0), 69 CF_{3}^{+} (26), 32 0_{2}^{+} (8.0), 28 C^{+} or N_{2}^{+} (36). C₆F₉OCF₃:
- $221 C_{6}F_{7}O^{+}(4), 193 C_{5}F_{7}^{+}(10), 181 C_{4}F_{7}^{+}(4), 162 C_{4}F_{6}^{+}(3), 159 C_{4}F_{5}O^{+}(2), 155 C_{5}F_{5}^{+}(2), 143$ $c_{4}F_{5}^{+}(7)$, 135 $c_{2}F_{5}O^{+}(7)$, 131 $c_{3}F_{5}^{+}(99)$, 119 $c_{2}F_{5}^{+}(6)$, 109 $c_{3}F_{3}O^{+}(6)$, 100 $c_{2}F_{4}^{+}(47)$, 69 $cF_{3}^{+}(7)$ $594 C_{12}F_{22}O_2^{+} (14), 506 C_{11}F_{18}O_2^{+} (4), 444 C_{9}F_{16}O_2^{+} (6), 421 C_{10}F_{15}O^{+} (10), 409 C_{9}F_{15}O^{+} (12),$ [c_sr_aocr₃]₂:
- $C_4F_5^+$ (10), 131 $C_3F_5^+$ (91), 119 $C_2F_5^+$ (12), 100 $C_2F_4^+$ (57), 93 $C_3F_3^+$ (13), 69 CF_3^+ (100), 31 CF^+ (4). $462 C_{10} F_{18}^{+} (0.0), 443 C_{10} F_{17}^{+} (15), 393 C_{9} F_{15}^{+} (5), 343 C_{8} F_{13}^{+} (56), 293 C_{7} F_{11}^{+} (10), 243 C_{6} F_{9}^{+} (14), (10), 100 F_{11}^{+} (10), 100 F_$ $205 \ C_{6} \Gamma_{7}^{+}(4), \ 193 \ C_{5} \Gamma_{7}^{+}(12), \ 181 \ C_{4} \Gamma_{7}^{+}(6), \ 169 \ C_{3} \Gamma_{7}^{+}(5), \ 162 \ C_{4} \Gamma_{6}^{+}(4), \ 155 \ C_{5} \Gamma_{5}^{+}(5), \ 143 \ C_{5}^{+}(5), \ 143 \ C_{5} \Gamma_{5}^{+}(5), \ 143 \ C_{5}^{+}(5), \$ [CsFg]2:

(100), 47 CF0⁺ (4), 32 0_2^+ (4), 31 CF⁺ (5), 28 CO⁺ or N_2^+ (15).

Compound	Relative Intensity	Chemical Shift ϕ (ppm from CFCl ₃)	Coupling Constants J Hz	Assignment
	en	67.8		1(e)
F5 F1	1	142.2		1(a)
	2	123.5		2(a), 6(a)
			J(2a,2e),J(6a,6e)=310±5	
	2	135.3		2(e), 6(e)
$F_{(e)}^4$ (a)	2	125.9		3(a), 5(a)
$F_{(e)}$			J(3a,3e),J(5a,5e)=295±5	
\mathbf{F}^{4} \mathbf{F}^{2} \mathbf{F}^{2} \mathbf{F}^{2} \mathbf{F}^{2}	7	137.8		3(e), 5(e)
	1	127.0		4 (a)
			J(4a,4e)=305±5	
	1	138.7		4(e)

¹⁹F NMR SPECTRA OF SOME PERFLUOROCYCLOHEXYL ETHERS

TABLE 3

	Relative	Chemical Shift	Coupling Constants	
Compound	Intensity	ç (ppm from CFC1,)	J Hz	Assignment
	c	c c t		,
	n	0.01	1771	le
	1	135.4		la
	, e	112.8		2a
1 1 1			J(2a,2e)=300±5	
r(a) $r(a)$	1	116.8		2e
Fe Contort	l	116.9		6a
r(e) $f(e)$ $f(e)$ $f(e)$			J(6a,6e)=260±5	
F^3 F (e)	1	122.1		6e
$\left(\begin{array}{c} -\frac{1}{2} \left(a \right) \\ r^{b} \end{array} \right)$	1	128.0		3a
			J(3a,3e)=260±5	
F ⁴ F ³	l	132.6		Зе
(e) (e) (e) (e)	1	138.6		ភូធ
F.4			J(5a,5e)=270±5	
(a) (a)	1	144.3		5e
	I	151.3		4a
			J(4a,4e)=300±5	
	I	155.5		4e
$F_{1,2}^{5}$ OC(CF_{3}^{1}) 3	18	70.7	7±1	la, 2a
(a) (a)	0	139.6		1e, 2e
F5.7.7 F6 F. F1	7	126.2		За, ба
			J(3a,3e),J(6a,6e)=300±5	
F3	7	132.5		3e, 6e
	2	129.7		4a, 5a
14 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1			J(4a,4e),J(5a,5e)=300±5	
$\mathbf{r}(\mathbf{e}) \mathbf{f} = \mathbf{r}(\mathbf{e}) \mathbf{f}(\mathbf{e})$	2	135.3		4e, 5e
$\mathbf{F}_{(a)}^4$ OC (CF ₂ ²) ₃				
(a)				

TABLE 3 (concluded)

MASS SPECTRA (LKB 9000) OF SOME PERFLUOROCYCLOHEXYL ETHERS

TABLE 4

- 516 $C_{10}F_{20}0^{+}$ (0.0), 497 $C_{10}F_{19}0^{+}$ (4.5), 478 $C_{10}F_{18}0^{+}$ (0.6), 459 $C_{10}F_{17}0^{+}$ (1.6), 428 $C_{9}F_{16}0^{+}$ $(5.3), 219 C_4F_9^+ (4.3), 193 C_5F_7^+ (2.0), 181 C_4F_7^+ (10.0), 169 C_3F_7^+ (0.6), 162 C_4F_6^+ (3.1), 150 C_7F_9^+ (10.1), 150$ $(1.3), \ 409 \ C_9F_{15}O^+ \ (9.3), \ 347 \ C_7F_{13}O^+ \ (2.5), \ 285 \ C_5F_{11}O^+ \ (3.1), \ 281 \ C_6F_{11}^+ \ (5.0), \ 231 \ C_5F_9^+$ $C_{3}F_{6}^{+}(0.6)$, 147 $C_{3}F_{8}O^{+}(3.1)$, 131 $C_{3}F_{5}^{+}(100.0)$, 119 $C_{2}F_{5}^{+}(4.4)$, 100 $C_{2}F_{4}^{+}(4.4)$, 97 $C_{2}F_{3}O^{+}(4.4)$ (4.4), 93 $C_3F_3^+$ (3.8), 69 CF_3^+ (62.5), 28 CO^+ or N_2^+ (1.3). C₆F₁, 0C(CF₃)₃:
- $C_{6}F_{10}[0C(CF_{3})_{3}]_{2}: \quad 732 \ C_{14}F_{28}0_{2}^{+} (0.2), \quad 713 \ C_{14}F_{27}0_{2}^{+} (18.2), \quad 694 \ C_{14}F_{26}0_{2}^{+} (0.4), \quad 675 \ C_{15}F_{25}0_{2}^{+} (3.0), \quad 551 \ C_{10}F_{21}0_{2}^{+} (0.5), \quad 510 \ C_{1$ $c_{10}F_{17}O^{+}(1.2), 447 c_{9}F_{17}O^{+}(0.8), 409 c_{9}F_{15}O^{+}(1.2), 397 c_{8}F_{15}O^{+}(1.4), 378 c_{8}F_{14}O^{+}(0.4), 60.4), 60.4$ $200 C_4 F_8^+ (1.2), 193 C_5 F_7^+ (2.0), 181 C_4 F_7^+ (5.0), 162 C_4 F_6^+ (3.2), 159 C_4 F_5 O^+ (2.4), 150 C_3 F_6^+ (3.2), 150 C$ $(0.6), \ 532 \ \dot{C}_{10}F_{20}O_2^+ \ (0.2), \ 513 \ C_{10}F_{19}O_2^+ \ (0.2), \ 497 \ C_{10}F_{19}O^+ \ (0.6), \ 485 \ C_{9}F_{19}O^+ \ (4.2), \ 459 \ C_{10}F_{19}O^+ \ (4.2), \ 459 \ C_{10}F_{10}O^+ \ (4.2), \ 459 \ C_{10}O^+ \ (4.2), \ 459$ $(1.2), 131 C_3F_5^+ (32.0), 119 C_2F_5^+ (2.4), 112 C_3F_4^+ (1.2), 109 C_3F_3^0^+ (1.4), 100 C_2F_4^+ (2.6), 1.2) C_3F_6^+ (2.6), 1.2) C_3F_6^+ (2.6) C_3F_6^- (2.6) C_3$ 97 $C_2^{F_3}O^+$ (2.8), 93 $C_3F_3^+$ (2.4), 69 CF_3^+ (100.0), 47 CFO^+ (2.0), 31 CF^+ (1.4).

 $CF_{3}OF$ in the presence of perfluorocycloolefins, confirms the stability of perfluoroalkoxy radicals in the following order:

$$CF_3O \cdot > (CF_3)_3CO \cdot$$

which agrees with the protonated analog.8,9

Reaction (5) accounts for the formation of perfluorocycloalkanes (C_5F_{10} and C_6F_{12}) and reaction (6) for the dimer of perfluorocycloalkanes.

$$F_{\bullet} + C_{5}F_{9} \cdot (\text{or } C_{5}F_{11} \cdot) \rightarrow C_{5}F_{10} (\text{or } C_{6}F_{12})$$

$$2C_{5}F_{9} \cdot (\text{or } 2C_{6}F_{11} \cdot) \rightarrow (C_{5}F_{9})_{2} (\text{or } (C_{6}F_{11})_{2})$$
(6)

Reaction (7) must occur to some extent to give radicals $(CF_3)_3 COC_5 F_8$. or $(CF_3)_3 COC_6 F_{10}$,

$$(CF_3)_3CO + C_5F_8 \quad (or \ C_6F_{10}) \rightarrow (CF_3)_3COC_5F_8 \cdot (or \ (CF_3)_3COC_6F_{10} \cdot)$$
(7)

which cross combine with perfluoro-t-butoxy radicals to form the small amount of vicinal perfluorodi-t-butoxycyclopentane $C_5F_8[OC(CF_3)_3]_2$ or vicinal perfluorodi-t-butoxycyclohexane $C_6F_{11}[OC(CF_3)_3]_2$.

$$(CF_3)_3CO. + (CF_3)_3COC_5F_8 \cdot (or (CF_3)_3COC_6F_{10} \cdot) \rightarrow C_5F_8[OC(CF_3)_3]_2$$
$$(or C_6F_{10}[OC(CF_3)_3]_2) (8)$$

The autocombination of perfluoro-t-butoxy radicals (Reaction 9) explains the formation of the main product, bis(perfluoro-t-butyl)peroxide.

$$2(CF_3)_3CO \rightarrow (CF_3)_3COOC(CF_3)_3$$
(9)

Thus the product formation of Reaction (8) can alternatively be expressed as follows:

$$(CF_3)_3 COOC(CF_3)_3 + C_5F_8 (or C_6F_{10}) \xrightarrow{h_{\mathcal{V}}} C_5F_8 [OC(CF_3)_3]_2$$

$$(or C_6F_{10} [OC(CF_3)_3]_2)$$

$$(10)$$

The photochemical reactions of primary fluoroxyperfluoroalkane (CF₃OF) and perfluorocycloolefins (C_5F_8 and C_6F_{10}) form mainly the simple adducts without forming bis(trifluoromethyl)peroxide or perfluorodimethoxycycloalkanes. The difference between the main reaction products of primary and of tertiary fluoroxyperfluoroalkanes can be explained by the shorter life-expectancy of trifluoromethoxy radical compared with perfluoro-t-butoxy radical because of the increased bulky groups surrounding the radical center. The longer lifetime of the perfluoro-t-butoxy radicals enables them to autocombine (Reaction 9), and the absence of CF₃OOCF₃ and perfluorodimethoxycycloalkanes is directly related to the low concentration of CF₃O. radicals. Since the rate of dimerization of alkoxy radicals to form the peroxide is dependent on $[RO.]^2$ due to the second-order reaction,⁸ the low concentration of CF₃O. radicals greatly decreases the CF₃OOCF₃ formation. Consequently, no perfluorodimethoxycycloalkanes are detected as in Reaction (10).

For vicinal perfluorodi-t-butoxycyclopentane, the presence of cisand trans-isomers in a 1:4 ratio is attributed to the involvement of free radical addition mechanisms (Reactions 7 and 8 or 10). The lack of stereospecificity excludes the involvement of tight ion pairs¹⁰ or a four-centered addition mechanism. The trans-addition is favored as a result of steric repulsion caused by the two bulky perfluoro-t-butoxy groups at the vicinal cis-position.

More complex ¹⁹F NMR spectra were obtained for perfluorocyclohexyl ethers (Table 3) than for perfluorocyclopentyl ethers (Table 1).¹¹ Feeney and Sutcliffe reported that, due to the strain and energetic grounds, perfluorocyclohexane and its derivatives have the chair configurations.¹² The bulky CF₃ group of perfluoromethylcyclohexane has been reported as fixed in the equatorial position.^{12,13} Table 3 shows the ¹⁹F NMR of perfluoromethoxy- and -t-butoxy groups at equatorial positions. Although the coupling constants and chemical shifts of perfluoromethoxy- and -methylcyclohexanes¹³ are similar, perfluoro-t-butoxycyclohexane is more complex. This complexity is attributed to some rotational barriers^{14,15} caused by the bulky perfluoro-t-butoxy group. The presence of only one isomer for vicinal perfluorodi-t-butoxycyclohexane is deduced from ¹⁹F NMR (Table 3) as the 1,2-diaxial isomer.

EXPERIMENTAL

Reagents

Perfluoro-t-butyl hypofluorite was prepared by direct fluorination of the sodium salt of perfluoro-t-butanol at -23° C.¹ The liquid product was confirmed by infrared and ¹⁹F NMR spectra as described by Prager and Thompson¹⁶ for (CF₂)₃COF.

Perfluoro-t-butanol, trifluoromethyl hypofluorite, perfluorocyclopentene, and perfluorocyclohexene were obtained from PCR. These reagents were checked by infrared and ¹⁹F NMR spectra and were freed from noncondensible impurities before use by condensation and pumping under vacuum in a trap cooled by liquid nitrogen.

Apparatus

All compounds were handled in quartz reaction vessels attached to a copper vacuum manifold system equipped with Monel valves and Cajon Ultratorr fittings. Pressures were measured by Ashcroft gauges to 75 psia and by a Pennwalt Wallace and Tiernan gauge (0-1550 mm Hg with an accuracy to 0.15 mm Hg). Amounts of volatile reactants were determined by P-V-T measurements, assuming ideal gas behavior.

The IR spectra were measured on a Perkin Elmer 467 spectrometer with a 5-cm Monel gas cell equipped with AgCl windows. The ¹⁹F NMR spectra were determined with a Varian XL-100 spectrometer operating at 94.1 Hz, using CFCl₃ as an external reference at 30° C. Mass spectra were obtained from an LKB 9000 mass spectrometer, which has an ionizing potential of 70 eV and a resolving power of 2000. The gas chromatograph-mass spectra were also run on the LKB 9000 instrument, using a 3 m x 2 mm glass column packed with 1% SE 30 on Gas-Chrom Q at 30° to 80° C and 3 atm nitrogen pressure. The elemental analyses were obtained by a high resolution mass spectrometer (CEC 21-110-B). The molecular weights or parent ions were also determined by a field ionization mass spectrometer.

Reactions of Perfluoro-t-butyl Hypofluorite

(a) With perfluorocyclopentene

Perfluorocyclopentene (2.4 mmol, 0.51 g) was condensed onto an evacuated frozen perfluoro-t-butyl hypofluorite (4.7 mmol, 1.19 g) in a quartz reaction vessel at a temperature of -196° C. The solid mixture was then warmed to -20° C and irradiated by a high pressure Hg arc lamp with a focused beam for 1 hr at -20° C. The liquid was chilled to -40° C

and evacuated. The volatiles (0.33 g, 19% yield) collected in a liquid nitrogen trap were identified by IR spectra as mainly CF_3COCF_3 and CF_4 with characteristic absorptions at 1890 and 2200 cm⁻¹, respectively. In addition small amounts of perfluorocyclopentane and perfluorocyclopentene were found with characteristic absorptions at 546 and 1776 cm⁻¹, respectively.

The residual mixture could not be cleanly separated by trap-

to-trap distillation and was identified by ¹⁹F NMR spectroscopy as $C_5F_9OC(CF_3)_3$ (Table 1, 0.85 mmol, 0.4 g) and the peroxide $(CF_3)_3COOC(CF_3)_3$ (1.3 mmol, 0.6 g). The ¹⁹F NMR of the latter compound shows a single resonance at 69.7 ppm relative to CFCl₃ as the external reference.

The liquid mixture was further analyzed by a gas chromatographmass spectrograph using a 3 m x 2 mm glass column packed with 1% SE-30 on Gas-Chrom Q at 50°C with 3 atm nitrogen carrier gas pressure. Table 2 shows the mass spectra (LKB 9000) of $C_5F_9OC(CF_3)_3$ and $C_5F_8[OC(CF_3)_3]_2$. At higher gas chromatograph temperature (80°C), the mass spectrum showed a trace of oligomers.

The field ionization mass spectrum exhibited two peaks. The strong peak was at mass m/e value of 470, which corresponds to the parent ion or molecular weight of $(CF_3)_3COOC(CF_3)_3$. The other peak was at mass m/e value of 466, which corresponds to the parent ion or molecular weight of $C_5F_9OC(CF_3)_3$.

The characteristic IR absorptions of $C_5F_9OC(CF_3)_3$ are at 1300 cm⁻¹ (s), 1290 cm⁻¹ (s), 1276 cm⁻¹ (s), 1235 cm⁻¹ (s), 1180 cm⁻¹ (m), 1130 cm⁻¹ (m), 1065 cm⁻¹ (w), 1004 cm⁻¹(s), 980 cm⁻¹ (s), 600 cm⁻¹ (w). Those of the peroxide (CF_3)_3COOC(CF_3)_3 are in accord with the literature.¹⁷

Mass spectroscopic molecular weight (CEC 21-110-B) of $C_5F_9OC(CF_3)_3$: Found 446.9685. Calculated for $C_9F_{17}O$ 446.9678. Mass spectroscopic molecular weight (CEC 21-110-B) of $C_5F_8[OC(CF_3)_3]_2$: Found, 681.9477. Calculated for $C_{1,3}F_{2,6}O_2$, 681.9483.

(b) With perfluorocyclohexene

Perfluorocyclohexene (2.0 mmol, 0.52 g) was condensed onto perfluoro-t-butyl hypofluorite (3.9 mmol, 0.99 g) and irradiated in the same manner as described above. The volatiles (0.35 g, 23% yield) were mainly CF_3COCF_3 and CF_4 in the presence of small amounts of perfluorocyclohexane and perfluorocyclohexene. The residual mixture, as deduced from ¹⁹F NMR, contained the peroxide $(CF_3)_3COOC(CF_3)_3$ (1.3 mmol, 0.6 g) and the simple adduct $C_6F_{11}OC(CF_3)_3$ (0.4 mmol, 0.2 g, Table 3) as the major products. The mixture was further analyzed by gas chromatograph-mass spectra. Table shows the mass spectra (LKB 9000) of $C_6F_{11}OC(CF_3)_3$ and $C_6F_{10}[OC(CF_3)_3]_2$. At higher gas chromatograph temperature (80°C) the mass spectrum showed a trace of oligomers.

When the stoichiometry was changed to an excess of perfluorocyclohexene (2.46 g, 9.4 mmol) over perfluoro-t-butyl hypofluorite (0.79 g 3.1 mmol), the amount of volatiles (i.e., CF_3COCF_3 and CF_4) was decreased (0.29 g, 9% yield). The peroxide (0.87 mmol, 0.41 g) was also decreased, while the perfluorocyclohexyl ethers and oligomers were increased.

Reactions of trifluoromethyl hypofluorite

(a) With perfluorocyclopentene

Perfluorocyclopentene (4.0 mmol, 0.848 g) and trifluoromethyl hypofluorite (8.0 mmol, 0.864 g) were irradiated at -60° C for l hr and evacuated at -85° C. The volatile fraction was identified by IR spectrum as pure CF₂OF (3.8 mmol, 0.395 g).

The residual mixture was identified by ¹⁹F NMR spectroscopy (Table 1) as $C_5F_9OCF_3$ (3.3 mmol, 1.04 g), C_5F_{10} (0.3 mmol, 0.075 g), the dimer of the simple adduct $(C_5F_8OCF_3)_2$ (0.13 mmol), and the dimer of perfluorocyclopentane $(C_5F_9)_2$ (0.02 mmol). The liquid mixture was further analyzed by a gas chromatograph-mass spectrograph, using the same column as described above but at 30°C with 3 atm nitrogen carrier gas pressure. Table 2 shows the mass spectra (LKB 9000) of $C_5F_9OCF_3$ and the dimers of the simple adduct $(C_5F_8OCF_3)_2$, and perfluorocyclopentane $(C_5F_9)_2$.

Mass spectroscopic molecular weight (CEC 21-110-B) of $(C_5F_8OCF_3)_2$ Found, 593,9547. Calculated for $C_{12}F_{22}O_2$, 593.9547. Mass spectroscopic molecular weight (CEC 21-110-B) for $(C_5F_9)_2$ is exactly the same as for $C_{10}F_{17}$.

(b) With perfluorocyclohexane

The reaction of perfluorocyclohexene (4.0 mmol, 1.05 g) and trifluoromethyl hypofluorite (8.0 mmol) was carried out in the same manner as the above procedure for CF_3OF and C_5F_8 . The volatile fraction was identified by IR as pure CF_3OF , and the residual liquid was identified by ¹⁹F NMR as quantitative and pure $C_8F_{1,1}OCF_3$ (Table 3).

Mass spectroscopic molecular weight (CEC 21-110-B) of $C_6F_{11}OCF_3$: Found, 346.9749. Calculated for $C_7F_{13}O$, 346.9749.

ACKNOWLEDGEMENTS

The authors wish to thank the Air Force Office of Scientific Research for support of this work, Dr. D. Thomas for mass spectra (LKB 9000 and CEC 21-110-B), Mr. L. Gary for 19 F NMR, and Mr. G. A. St. John for the field ionization mass spectra.

- 1. M. S. Toy and R. S. Stringham, J. Fluorine Chem., 5, in press (paper I).
- 2. J. A. Young, Fluorine Chem. Rev., 1, (1967) 359.
- 3. R. D. Smith, F. C. Fawcett and D. D. Coffman, J. Amer. Chem. Soc., 84 (1962) 4285.
- 4. M. S. Toy and R. S. Stringham, J. Fluorine Chem., 5, in press (paper II).
- 5. W. T. Miller and M. B. Freedman, J. Amer. Chem. Soc., 85 (1963) 180.
- P. E. Aldrich, E. G. Howard, W. J. Linn, W. J. Middleton and W. H. Sharkey, J. Org. Chem., 28 (1963) 184.
- 7. R. S. Porter and G. H. Cady, J. Amer. Chem. Soc., 79 (1957) 5625.
- 8. P. Gray and A. Williams, Chem. Rev., 59 (1959) 239.
- 9. P. Gray, R. Shaw and J.C.T. Thynne, Progr. Reaction Kinetics, 4 (1967) 63.
- D.H.R. Barton, R. H. Hesse, G. P. Jackman, L. Ogunkoya and M. M. Pechet, J. Chem. Soc. Perkin I 7 (1964) 739.
- J. R. van Wazer, <u>Compilation of Reported F¹⁹ NMR Chemical Shifts</u>, Wiley-Interscience, New York (1970).
- 12. J. Feeney and L. H. Sutcliffe, Trans. Faraday Soc., 56 (1960) 1559.
- 13. K. W. Jolley and L. H. Sutcliffe, Trans. Faraday Soc., 64 (1968) 269.
- 14. A. Peake and L. F. Thomas, Trans. Faraday Soc., 62 (1966) 2980.
- 15. G.V.D. Tiers, Proc. Chem. Soc., (1960) 389.
- 16. J. H. Prager and P. G. Thompson, J. Amer. Chem. Soc., 87 (1965) 230.
- 17. D. E. Gould, C. T. Ratcliffe, L. R. Anderson and W. B. Fox, <u>Chem.</u> Comm., (1970) 216.