

Acta Cryst. (1998). **C54**, 103–107**Photocleavage Reactions of Monothioimides**

TAI Y. FU, JOHN R. SCHEFFER AND JAMES TROTTER

Department of Chemistry, University of British Columbia, Vancouver, BC, Canada V6T 1Z1. E-mail: jtrt@xray4.chem.ubc.ca

(Received 29 April 1997; accepted 1 September 1997)

Abstract

The photochemistry and crystal structures of four monothioimides have been studied: *N*-phenyl-3-phenyl-*N*-[phenyl(thiocarbonyl)]propanamide, C₂₂H₁₉NOS, (1a); *N*-(4-bromophenyl)-*N*-[phenyl(thiocarbonyl)]-3-phenylpropanamide, C₂₂H₁₈BrNOS, (1b); *N*-(4-bromo-2,6-dimethylphenyl)-3-phenyl-*N*-[phenyl(thiocarbonyl)]propanamide, C₂₄H₂₂BrNOS, (1c); and *N*-[4-methoxyphenyl(thiocarbonyl)]-*N*-phenyl-3-phenylpropanamide, C₂₃H₂₁NO₂S, (1d). All four molecules adopt similar (*E*_S, *Z*_O) conformations, which are not suitable for photochemical γ -hydrogen abstraction; short intramolecular O...C(S) contacts (2.8 Å) suggest a new mechanism with nucleophilic attack by the carbonyl O atom on the C=S double bond; the presence of water is also essential for the reaction to proceed.

Comment


Acyclic monothioimides of general structure (1) [see scheme below, which shows two conformations of the monothioimides; others are (*Z*, *Z*) and (*E*, *E*)] undergo photochemical reactions when irradiated in solution to give, among other products, thiobenzanilides, (2) (Sakamoto *et al.*, 1991). The reasonable mechanism suggested for this reaction involves Norrish type II γ -

hydrogen abstraction by sulfur (see scheme), followed by 1,4-biradical cleavage.

As part of a determination of the geometrical requirements for intramolecular hydrogen abstraction by sulfur (Fu, Scheffer & Trotter, 1997), the monothioimides (1a)–(1d) have now been synthesized, their photochemistry studied [in solution and in the solid state; the solution photochemistry of (1a) and (1d) had been studied previously by Sakamoto *et al.* (1991)] and crystal structures determined. Irradiation with a 450 W Hanovia lamp of solutions of (1a)–(1d) in anhydrous benzene results in no photochemical reaction (Fu, Scheffer & Trotter, 1994). In benzene which has been allowed to equilibrate with atmospheric moisture, photolysis gives the thiobenzanilides (2a)–(2d) and 3-phenylpropanoic acid. Irradiations in benzene containing 10% methanol give (2a)–(2d) and methyl 3-phenylpropanoate. Photolysis of (1a)–(1d) in the solid state with rigorous exclusion of moisture again results in no photoreaction; when no special attempts are made to exclude moisture, irradiation leads to the same two photoproducts that are formed in benzene solution.

The crystal structure determinations of the monothioimides (1a)–(1d) show that all four molecules adopt similar (*E*_S, *Z*_O) conformations in the solid state (Fig. 1). *MMX* calculations (*PCModel*; Serena Software, 1988) indicate that this is the most stable conformation, presumably as a result of more favourable alignment of the carbonyl and thiocarbonyl dipoles, and smaller 1,3-steric repulsions (Fu, 1994). The (*E*_O, *Z*_S) conformation (scheme above), which is required for γ -hydrogen abstraction by sulfur, is higher in energy by 1.5 kJ mol^{−1}, with the (*Z*, *Z*) and (*E*, *E*) conformations intermediate in energy [1 kJ mol^{−1} above (*E*_S, *Z*_O)]. In the observed (*E*_S, *Z*_O) conformation, the shortest C=S... γ -H contacts [S1...H (on C15); Fig. 1] are 5.06, 5.14, 5.07 and 5.07 Å, for (1a)–(1d), respectively, considerably greater than the sum of the van der Waals radii for S and H (3.00 Å), and too distant to allow for hydrogen abstraction (Fu *et al.*, 1997). The shortest S... δ -H contacts are even longer (6.2–6.6 Å). To verify the lack of hydrogen abstraction, monothioimide (1a) containing two γ -deuterium atoms was synthesized and subjected to benzene-solution and solid-state photolysis under non-anhydrous conditions. In each case, the 3-phenylpropanoic acid photoproduct shows no loss of the deuterium label (γ -hydrogen abstraction would lead to a monodeuterated acid). Further corroboration is obtained by irradiating unlabelled (1a) in CH₃OD/benzene, which gives undeuterated methyl 3-phenylpropionate.

Hence it is established that photocleavage of monothioimides does not involve initial γ -hydrogen abstraction by sulfur, and another mechanism must be sought. A clue to a reasonable mechanistic possibility is found in the observed intramolecular contacts between the O atoms of the carbonyl (CO) groups and the C atoms of the thiocarbonyl (CS) groups: O1...C1 = 2.862 (3),






Fig. 1. Views of the molecules of (1a), (1b), (1c) and (1d) (33% probability ellipsoids).

2.769 (3), 2.807 (7) and 2.811 (3) Å, for (1a)–(1d), respectively (Fig. 1). These distances are well below the sum of the van der Waals radii for O and C (3.22 Å), so that the O atom is ideally positioned for excited-state nucleophilic attack on the C=S double bond. A mechanism may therefore be proposed (Fig. 2) involving formation of a 1,3-oxazetidinium ion, (3), which could react with water present in the reaction medium

to form (4), which could then break down, by a process which has been observed previously (Barton *et al.*, 1973), to form the enol of the thiobenzanilide and 3-phenylpropanoic acid. Attempts to trap the intermediate (3) with methyl iodide were unsuccessful.

Having established that photocleavage of the mono-thioimides (1a)–(1d) does not involve γ -hydrogen abstraction, it was of interest to discover whether fully

oxygenated imides behave similarly. To this end, *N*-acetylpropanamide was synthesized and photolysed in CH₃OD; incorporation of one deuterium atom into the methyl propanoate photoproduct indicates that γ -hydrogen abstraction is occurring in this case (Mazzocchi *et al.*, 1980). This is in accord with the preferred (*E,Z*) conformation for such imides in solution (Noe & Raban, 1975), which has the abstracting O and abstracted H in close proximity.




Fig. 2. Proposed mechanism for the photoreaction of the monothioimides.

All four monothioimide molecules, (1*a*)–(1*d*), have normal geometry and dimensions. The thioimide groups are roughly planar, presumably as a result of delocalization of the N-atom lone-pair electrons into the carbonyl and thiocarbonyl π systems. The S=C–N–CO grouping deviates from an exact *anti* (*E*) conformation by 20–30° [torsion angles –158.3 (2), –145.7 (2), –146.8 (4) and –149.2 (2)°; Table 1], and the SC–N–C=O grouping deviates by up to 30° from an exact *syn* (*Z*) conformation [torsion angles 29.9 (3), 23.8 (4), 13.4 (8) and 0.4 (4)°]. The phenylpropyl chains have *anti* conformations in (1*b*) and (1*d*) [C14–C15–C16–C17 torsion angles –167.6 (3) and 178.0 (2)°, respectively], but *gauche* conformations in (1*a*) and (1*c*) [torsion angles –68.6 (3) and –68.4 (7)°, respectively].

Experimental

Details of the synthesis and photochemistry have been deposited with the IUCr.

Compound (1*a*)

Crystal data

C₂₂H₁₉NOS
M_r = 345.46

Cu *K* α radiation
 λ = 1.5418 Å

Orthorhombic

*P*2₁2₁2₁
a = 8.022 (4) Å
b = 38.685 (4) Å
c = 5.847 (3) Å
V = 1815 (1) Å³
Z = 4
D_x = 1.264 Mg m^{–3}
D_m not measured

Data collection

Rigaku AFC-6S diffractometer
 ω –2 θ scans
Absorption correction:
 ψ scans (North, Phillips & Mathews, 1968)
*T*_{min} = 0.633, *T*_{max} = 0.787
2113 measured reflections
2113 independent reflections

Refinement

Refinement on *F*²
R(*F*) = 0.030
wR(*F*²) = 0.085 (0.113 for the enantiomeric structure)
S = 1.71
2113 reflections
227 parameters
H atoms not refined
w = 1/[$\sigma^2(F_o^2)$ + 0.00022(*F_o*²)²]

Cell parameters from 24 reflections
 θ = 44.3–54.3°
 μ = 1.598 mm^{–1}
T = 294 K
Prism
0.30 × 0.15 × 0.15 mm
Orange

1861 reflections with *I* > 3 σ (*I*)
 $\theta_{\max}$ = 77.42°
h = 0 → 10
k = 0 → 48
l = 0 → 6
3 standard reflections every 200 reflections
intensity decay: 0.17%

(Δ/σ)_{max} = 0.019
 $\Delta\rho_{\max}$ = 0.26 e Å^{–3}
 $\Delta\rho_{\min}$ = –0.21 e Å^{–3}
Extinction correction: Zachariasen (1967)
Extinction coefficient: 4.3 (7) × 10^{–6}
Scattering factors from *International Tables for Crystallography* (Vol. C)

Compound (1*b*)

Crystal data

C₂₂H₁₈BrNOS
M_r = 424.35
Monoclinic
*P*2₁/*a*
a = 9.506 (1) Å
b = 17.643 (2) Å
c = 11.960 (1) Å
 β = 91.65 (1)°
V = 2005.1 (4) Å³
Z = 4
D_x = 1.406 Mg m^{–3}
D_m not measured

Data collection

Rigaku AFC-6S diffractometer
 ω –2 θ scans
Absorption correction:
 ψ scans (North, Phillips & Mathews, 1968)
*T*_{min} = 0.171, *T*_{max} = 0.316
4584 measured reflections
4178 independent reflections

Cu *K* α radiation
 λ = 1.5418 Å
Cell parameters from 23 reflections
 θ = 55.6–57.0°
 μ = 3.839 mm^{–1}
T = 294 K
Prism
0.60 × 0.40 × 0.30 mm
Orange

3479 reflections with *I* > 3 σ (*I*)
*R*_{int} = 0.015
 $\theta_{\max}$ = 77.47°
h = 0 → 11
k = 0 → 22
l = –15 → 15
3 standard reflections every 200 reflections
intensity decay: none

RefinementRefinement on F^2 $R(F) = 0.044$ $wR(F^2) = 0.130$ $S = 3.63$

4178 reflections

236 parameters

H atoms not refined

 $w = 1/[\sigma^2(F_o^2) + 0.00006(F_o^2)^2]$ $(\Delta/\sigma)_{\max} = 0.007$ $\Delta\rho_{\max} = 0.83 \text{ e } \text{\AA}^{-3}$

(0.9 Å from Br)

 $\Delta\rho_{\min} = -0.70 \text{ e } \text{\AA}^{-3}$

Extinction correction:

Zachariasen (1967)

Extinction coefficient:

 $8.6(4) \times 10^{-6}$

Scattering factors from

International Tables for Crystallography (Vol. C)**Compound (1c)****Crystal data**C₂₄H₂₂BrNOS $M_r = 452.41$

Orthorhombic

*Pca*2₁ $a = 38.032(3) \text{ \AA}$ $b = 7.373(4) \text{ \AA}$ $c = 7.725(4) \text{ \AA}$ $V = 2166(2) \text{ \AA}^3$ $Z = 4$ $D_x = 1.387 \text{ Mg m}^{-3}$ D_m not measuredMo $K\alpha$ radiation $\lambda = 0.7107 \text{ \AA}$

Cell parameters from 25

reflections

 $\theta = 12.6\text{--}16.3^\circ$ $\mu = 1.983 \text{ mm}^{-1}$ $T = 294 \text{ K}$

Prism

 $0.30 \times 0.20 \times 0.20 \text{ mm}$

Red

Data collection

Rigaku AFC-6S diffractometer

 ω scan

Absorption correction:

 ψ scans (North, Phillips & Mathews, 1968) $T_{\min} = 0.650$, $T_{\max} = 0.673$

2844 measured reflections

2844 independent reflections

1302 reflections with

 $I > 3\sigma(I)$ $\theta_{\max} = 30.00^\circ$ $h = 0 \rightarrow 53$ $k = 0 \rightarrow 9$ $l = 0 \rightarrow 10$

3 standard reflections

every 200 reflections

intensity decay: 3.3%

RefinementRefinement on F^2 $R(F) = 0.033$ $wR(F^2) = 0.082(0.100)$

for opposite polarity)

 $S = 1.18$

2844 reflections

252 parameters

H atoms not refined

 $w = 1/[\sigma^2(F_o^2) + 0.00016(F_o^2)^2]$ $(\Delta/\sigma)_{\max} = 0.004$ $\Delta\rho_{\max} = 0.87 \text{ e } \text{\AA}^{-3}$ $\Delta\rho_{\min} = -2.16 \text{ e } \text{\AA}^{-3}$

(at Br site)

Extinction correction: none

Scattering factors from

International Tables for Crystallography (Vol. C)**Compound (1d)****Crystal data**C₂₃H₂₁NO₂S $M_r = 375.48$

Orthorhombic

Pbca $a = 26.345(6) \text{ \AA}$ $b = 24.531(7) \text{ \AA}$ $c = 6.073(10) \text{ \AA}$ $V = 3925(9) \text{ \AA}^3$ $Z = 8$ $D_x = 1.271 \text{ Mg m}^{-3}$ D_m not measuredMo $K\alpha$ radiation $\lambda = 0.7107 \text{ \AA}$

Cell parameters from 17

reflections

 $\theta = 10.1\text{--}18.9^\circ$ $\mu = 0.182 \text{ mm}^{-1}$ $T = 294 \text{ K}$

Prism

 $0.30 \times 0.30 \times 0.15 \text{ mm}$

Orange-red

Data collection

Rigaku AFC-6S diffractometer

 ω -2 θ scans

Absorption correction:

 ψ scans (North, Phillips

& Mathews, 1968)

 $T_{\min} = 0.904$, $T_{\max} = 1.000$

5741 measured reflections

5741 independent reflections

1604 reflections with

 $I > 3\sigma(I)$ $\theta_{\max} = 30.01^\circ$ $h = 0 \rightarrow 37$ $k = 0 \rightarrow 34$ $l = -8 \rightarrow 0$

3 standard reflections

every 200 reflections

intensity decay: 1.0%

RefinementRefinement on F^2 $R(F) = 0.046$ $wR(F^2) = 0.122$ $S = 1.03$

5741 reflections

244 parameters

H atoms not refined

 $w = 1/[\sigma^2(F_o^2) + 0.00022(F_o^2)^2]$ $(\Delta/\sigma)_{\max} = 0.0013$ $\Delta\rho_{\max} = 0.87 \text{ e } \text{\AA}^{-3}$

(0.6 Å from S)

 $\Delta\rho_{\min} = -1.07 \text{ e } \text{\AA}^{-3}$

(at S site)

Extinction correction: none

Scattering factors from

International Tables for Crystallography (Vol. C)

Table 1. Comparison of selected bond distances (Å) and angles (°)

	(1a)	(1b)	(1c)	(1d)
C—Br		1.897(2)	1.907(5)	
C1=S1	1.643(2)	1.630(2)	1.632(6)	1.628(2)
C14=O1	1.202(3)	1.204(3)	1.194(6)	1.202(2)
N1—C1	1.383(2)	1.398(3)	1.399(7)	1.397(3)
N1—C14	1.431(2)	1.417(3)	1.418(7)	1.433(3)
C1—N1—C8	120.3(2)	121.1(2)	117.5(4)	118.2(2)
C1—N1—C14	123.0(2)	120.2(2)	122.4(4)	122.5(2)
C8—N1—C14	115.8(2)	118.1(2)	119.1(5)	118.2(2)
S1=C1—N1—C14	−158.3(2)	−145.7(2)	−146.8(4)	−149.2(2)
C1—N1—C14=O1	29.9(3)	23.8(4)	13.4(8)	0.4(4)

H atoms were fixed in ideal sites; for (1c), the methyl H-atom sites were based on difference peaks.

For all compounds, data collection: *MSC/AFC Diffractometer Control Software* (Molecular Structure Corporation, 1993); cell refinement: *MSC/AFC Diffractometer Control Software*; data reduction: *TEXSAN* (Molecular Structure Corporation, 1995); program(s) used to solve structure: *SIR92* (Altomare *et al.*, 1993); program(s) used to refine structure: *TEXSAN*; software used to prepare material for publication: *TEXSAN*.

We thank the Natural Sciences and Engineering Research Council of Canada for financial support.

Supplementary data for this paper are available from the IUCr electronic archives (Reference: FG1342). Services for accessing these data are described at the back of the journal. Details of the synthesis and photochemistry of the four title compounds have also been deposited.

References

- Altomare, A., Cascarano, M., Giacovazzo, C. & Guagliardi, A. (1993). *J. Appl. Cryst.* **26**, 343.
 Barton, D. H. R., Bolton, M., Magnus, P. D. & West, P. J. (1973). *J. Chem. Soc. Perkin Trans. 1*, pp. 1580–1583.

- Fu, T. Y. (1994). PhD thesis, University of British Columbia, Vancouver, Canada.
- Fu, T. Y., Scheffer, J. R. & Trotter, J. (1994). *Tetrahedron Lett.* **35**, 3235–3238.
- Fu, T. Y., Scheffer, J. R. & Trotter, J. (1997). *Acta Cryst.* **C53**, 1257–1259.
- Mazzocchi, P. H., Jameson, W., Nishiyama, T. & DeCamp, A. (1980). *Tetrahedron Lett.* **21**, 989–992.
- Molecular Structure Corporation. (1993). *MSC/AFC Diffractometer Control Software*. MSC, 3200 Research Forest Drive, The Woodlands, TX 77381, USA.
- Molecular Structure Corporation. (1995). *TEXSAN. Single Crystal Structure Analysis Software*. Version 1.7. MSC, 3200 Research Forest Drive, The Woodlands, TX 77381, USA.
- Noe, E. A. & Raban, M. (1975). *J. Am. Chem. Soc.* **97**, 5811–5820.
- North, A. C. T., Phillips, D. C. & Mathews, F. S. (1968). *Acta Cryst.* **A24**, 351–359.
- Sakamoto, M., Tohnishi, M., Fujita, T. & Watanabe, S. (1991). *J. Chem. Soc. Perkin Trans. 1*, pp. 347–351.
- Serena Software (1988). *PCModel*. Version 88.00. Serena Software, Burlingame, CA 94010-1904, USA.
- Zachariasen, W. H. (1967). *Acta Cryst.* **23**, 558–564.

Acta Cryst. (1998). **C54**, 107–109

Bis(4-butoxyphenyl) Terephthalate

ROBERTO CENTORE AND ANGELA TUZI

Dipartimento di Chimica, Università di Napoli 'Federico II', Via Mezzocannone 4, 80134 Napoli, Italy. E-mail: centore@chemna.dichi.unina.it

(Received 27 May 1997; accepted 29 July 1997)


Abstract

The crystal and molecular structures of nematogenic bis(4-butoxyphenyl) terephthalate, $C_{28}H_{30}O_6$, are reported. The molecular structure, determined by X-ray diffraction, is very different from that of the same compound reported on the basis of solid-state NMR analysis [Uryu & Kato (1988). *Macromolecules*, **21**, 378–384]. In the structure reported here, the carbonyl group of the ester is, as expected, almost coplanar with the phenyl ring of the terephthalate. The conformation of the aliphatic chain is not fully *trans*-planar; there is a *gauche* bond. Molecules are stacked in layers with a herring-bone type of packing within the layers.

Comment

Terephthalic and *p*-hydroxybenzoic acids are among the most widely used groups for the synthesis of both monomers and polymers which form liquid crystals. Condensation of these groups with various *p,p'*-difunctionalized aromatic compounds leads to linear high axial ratio structures (esters, amides, *etc.*) which

favor the formation of liquid-crystal phases (Ober, Jin & Lenz, 1984). In previously reported structures of esters of terephthalic, *p*-hydroxybenzoic and related aromatic carboxylic acids (Schweizer & Dunitz, 1982; Centore *et al.*, 1991), the carbonyl group of the ester is found to be coplanar or almost coplanar with the phenyl ring to which it is attached. Consistent with this, semi-empirical quantum-mechanical calculations for 'gas-phase' phenyl benzoate (Bicerano & Clark, 1988) indicate an energy about 6.7 kcal mol⁻¹ greater for the 'perpendicular' arrangement of carbonyl group and phenyl ring, compared with the lowest energy coplanar conformation. Deviations from the coplanar arrangement are only observed as a result of great overcrowding, *e.g.* in the crystal structures of 2,6-disubstituted benzoic acids (Anca, Martinez-Carrera & Garcia-Blanco, 1967; Florencio & Smith, 1970). On the basis of solid-state CP/MAS NMR analysis on crystals of the title compound, (I), Uryu & Kato (1988) reported that the carbonyl group of the ester is almost perpendicular to the plane of the terephthalate phenyl ring. They also reported that the planes of the three phenyl rings of the molecule were parallel, which is rare in the crystal structures of phenyl esters of aromatic acids, and that the aliphatic chains were planar zigzag. We undertook the crystal structure determination reported here in order to establish definitively the conformation in the solid state.



The thermal properties of the title compound as prepared by us are consistent with literature data (crystal–nematic transition: $T_m = 460.4$ K, $\Delta H_m = 45.8$ kJ mol⁻¹; nematic–isotropic transition: $T_i = 505.0$ K, $\Delta H_i = 0.45$ kJ mol⁻¹). Particular care was taken to detect possible solid-state polymorphs; samples recrystallized from ethyl acetate [as reported by Uryu & Kato (1988)] and from various other solvents or solvent mixtures (chloroform–ethanol, chloroform–*n*-hexane, toluene) all have identical DSC thermograms.

The molecules in the crystals are centrosymmetric and belong to point group $\bar{1}$ (C_i). Bond lengths and angles have normal values. The conformation of the molecule, however, is considerably different from that proposed by Uryu & Kato (1988). The carbonyl group of the ester is almost coplanar with the phenyl ring of the terephthalate [O3—C11—C12—C13 = 176.3 (2)°], as is usual in terephthalate esters. The dihedral angle between the planes of the two phenyl rings, C5—C10 and C12—C14, is 65.06 (5)° and is mainly due to the torsion angle around the O2—C8 bond. Furthermore, the conformation of the aliphatic chain is not fully *trans*-planar, a *gauche* conformation being observed around the C3—C4 bond.