

View Article Online View Journal

ChemComm

Accepted Manuscript

This article can be cited before page numbers have been issued, to do this please use: J. Yuan, X. Ma, H. Yi, C. Liu and A. Lei, *Chem. Commun.*, 2014, DOI: 10.1039/C4CC05661B.



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about *Accepted Manuscripts* in the **Information for Authors**.

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard <u>Terms & Conditions</u> and the <u>Ethical guidelines</u> still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.



www.rsc.org/chemcomm

RSCPublishing

Journal Name

COMMUNICATION

Cite this: DOI: 10.1039/x0xx00000x

I₂-Catalyzed Oxidative C(sp³)-H/S-H Coupling: Utilizing Alkanes and Mercaptans as the Nucleophiles

Jiwen Yuan,^a Xu Ma^a Hong Yi,^a Chao Liu,^a and Aiwen Lei^{*,a,b}

Received ooth January 2014, Accepted ooth January 2014

DOI: 10.1039/x0xx00000x

www.rsc.org/

By using alkanes and mercaptans as the nucleophiles, di-tertbutyl peroxide (DTBP) as the oxidant, I₂-catalyzed oxidative $C(sp^3)$ -H/S-H coupling was achieved. This protocol provided a novel process to construct $C(sp^3)$ -S bond from commercially available hydrocarbons and mercaptans.

C-S bonds are key structures of many nature products and functional molecules which exhibit different biological activities, hence much attention has been paid to the formation of C-S bond through coupling reactions¹. Classically, coupling reactions of organic halides/organic boronic acid with mercaptans in the presence of bases and catalysts was employed to form C-S bonds². Oxidative coupling between RH and R'H, which shortens the synthesis route due to the elimination of pre-activation of C-H or X-H bonds, has attracted much attention during the past decades'. Although transition-metal-catalyzed $C(sp^2)$ -H and C(sp)-H functionalization were the major force in this field⁴, they were seldom utilized to perform oxidative coupling with S-H bonds⁵ due to the strong poisoning effect of mercaptans. Moreover, up to now, only isolated reports focus on oxidative C(sp³)-H/S-H coupling⁶. Thus, it is still highly desirable to explore the C-S bond formation utilizing C(sp³)-H bonds and mercaptans as the nucleophiles.

 $C(sp^3)$ -H bonds existed extensively in the nature, and most of which were of alkanes. Thus it is a pivotal task to realize the functionalization of the abundant $C(sp^3)$ -H bonds of alkanes. However, they are difficult to be activated in reactions, which may be ascribe to their nonpolarity, which weakened the interaction between the bonds and other species such as transition metals. Consequently, to date, only few reports have described oxidative coupling utilizing simple alkanes in highly selective manner⁷ although they are always easy accessible. Therefore, it is still a crucial but challenging task to achieve the oxidative coupling of the $C(sp^3)$ -H bonds of simple alkanes. Moreover, there is no report focused on the oxidative C-H/S-H coupling of $C(sp^3)$ -H bonds between simple alkanes and mercaptans, so such topic of $C(sp^3)$ -S bonds (Scheme 1).



Scheme 1. New way for C(sp3)-S bond construction

In consequence of the strong coordination of the mercaptans to the transition metals, and the use of metal limited the further application of the protocols in industry and pharmacy. We initiated the investigation with a metal-free protocol⁸. Recently, iodide catalysts are gaining more attention among the oxidative coupling reactions9. Moreover, compared with transition metals, iodide catalysts might exhibit weaker interaction with mercaptans. Thus, this serious of catalysts were picked out to catalyze the oxidative C(sp³)-H/S-H coupling. To figure out the best condition, 4-methyl thiophenol 2a was employed to react with toluene 1a. After a series of condition optimization, I_2 and DTBP were selected to be the catalyst and the oxidant respectively (see supporting information, Table S1). The detailed condition was: 1a (4 mL), 2a (0.3 mmol), DTBP (1.5 mmol), I_2 (0.045 mmol) at 120 °C for 20 h under N₂. The corresponding sulfide was obtained in the yield of 74% under the optimized condition (Scheme 2, **3aa**)¹⁰.

Then we began to investigate the substrate scope of the reaction. First, various aryl mercaptans were examined to react with toluene 1a under optimized condition. Electron-rich thiophenols performed the benzylation smoothly and good yields were obtained (Scheme 2, 3aa, 3af, 3ag). The electrondeficient thiophenol, however, exhibited lower activity than the electron-rich ones (Scheme 2, 3ae). The halo-substituted thiophenols, which allowed further functionalization, were tolerated (Scheme 2, 3ab, 3ac, 3ad, 3ah). The steric hindrance effect was not obvious in the reactions of thiophenols bearing o-Br and p-Br group (Scheme 2, 3ad, 3ah). It was noteworthy that both electron-rich and electron-deficient heterocyclic mercaptans were also suitable for this transformation (Scheme 2, 3ai, 3aj). Moreover, 4-mercapto-phenyl sulfide 2k, a substrate bearing two mercapto group, also react with toluene with good selectivity that only dibenzylated product was

obtained (Scheme 2, **3ak**). Unfortunately, only trace amount of product were detected when aliphatic mercaptans were employed to couple with toluene, which might result from the over-oxidation of the relatively unstable aliphatic mercaptans.



Scheme 2. I₂-promoted oxidative $C(sp^3)$ -H/S-H coupling of toluene with aryl mercaptans. Reaction conditions: **1a** (4 mL), **2** (0.3 mmol), DTBP (1.5 mmol), I₂ (0.045 mmol) at 120 °C for 20 h under N₂. Yields shown are of isolated products. Bn = benzyl. [a] **2k** (0.15 mmol).

Inspired by the encouraging results, the reactivity of methyl arenes was further investigated. 4-methoxy-thiophenol **2f** was selected to react with various methyl arenes due to the easier isolation of the products. The methyl arenes bearing multiple methyls, which enriched the electron density of the arene slightly, exhibited good activity (Scheme 3, **3bf**, **3ff**). However, methyl arenes bearing electron-deficient group such as Cl and Br group exhibited lower activity in the reaction (Scheme 3, **3cf**, **3df**, **3ef**). Steric effect was also not observed since the *o*-Br and *p*-Br toluenes showed similar activity (Scheme 3, **3df**, **3ef**).



Scheme 3. I₂-promoted oxidative C(sp3)-H/S-H coupling of various methyl arenes with 4-methoxy-thiophenol. Reaction conditions: 1 (4 mL), 2f (0.3 mmol), DTBP (1.5 mmol), I₂ (0.045 mmol) at 120 °C for 20 h under N₂. Yields shown are of isolated products.

Since the bond dissociation energy (BDE) of toluene $C(sp^3)$ -H bond (89 Kcal/mol¹¹) was low among the $C(sp^3)$ -H bonds, cyclohexane, another alkane with relatively low $C(sp^3)$ -H bond energy (96 kcal/mol¹²), was tested utilizing the standard condition. To our delight, the corresponding phenyl alkyl sulfide was obtained in the yield of 80% (Scheme 4, **5af**). Moreover, the BDE of $C(sp^3)$ -H bond of acetone was 95

kcal/mol¹³, and as the result, acetone, the most simple ketone, can also perform the $C(sp^3)$ -H/S-H coupling as predicted with lower yield (Scheme 4, **5bf**). With the C-H bond of acetone activated, acetylacetone reacted with 4-methoxy-thiophenol smoothly and the yield rose to 90% (Scheme 4, **5cf**).



Scheme 4. I₂-promoted oxidative $C(sp^3)$ -H/S-H coupling of various various $C(sp^3)$ -H source with 4-methoxy-thiophenol. Reaction conditions: 4 (4 mL), 2f (0.3 mmol), DTBP (1.5 mmol), I₂ (0.045 mmol) at 120 °C for 20 h under N₂. Yields shown are of isolated products.

Furthermore, we wanted to gain some mechanistic insights of this oxidative coupling of alkane $C(sp^3)$ -H bonds with aryl mercaptans. Control experiments were carried out firstly. When I₂ was excluded from the system, the yield of sulfide lowered slightly [Scheme 5, eq(1)]. Which indicated that I₂ was involved among the transformation. Then, benzyl iodide was employed to react with 4-methylthiophenol (see supporting imformation, Scheme S1) and only 30% of the aryl alkyl sulfide was obtained. This result suggested that the iodization/substitution procedure may not be the main access led to the formation of product.

Then in situ IR was selected to monitor the reaction between toluene and 4-methylthiophenol to figure out the role of I₂. We can see that 4-methyl-thiophenol consumed within 3 minutes accompanying the generation of 4-methylphenyl disulfide (see supporting information, Figure S1). The sampling experiment revealed that the disulfide generated very fast with the yield of 79% (Figure 1, disulfide with I_2), and the product did not form until the fully consumption of mercaptans (Figure 1, product with I₂). Slower generation of the disulfide and lower yield of the product were observed when I2 was taken away from the system (Figure 1, disulfide and product without I2). Therefore, we drew the inference that the disulfides may be the intermediate of the reaction, and I2 may act as accelerator of the generation of disulfide¹⁴. The accelerated dimerization might prevent the side reactions of mercaptans, which may result to the slight promotion of the yield. Another control experiment showed that disulfide exhibited similar activity compared with mercaptan [Scheme 5, eq(2)], which further confirmed the deduction.

In view of the previous reports employed methyl arenes as the benzyl source and DTBP as the oxidant, a single electron transfer (SET) process led to the generation of benzyl radical might be one step among the reaction^{7d, 15}. Therefore, a radicaltrapping experiment was carried out using butylated hydroxytoluene (BHT) as the radical trapper. As predicted, the coupling reaction was suppressed by BHT [Scheme 5, eq(3)]. Thus we believed that a radical process was involved among the reaction.

Then, to further confirm our speculation, electron paramagnetic resonance (EPR) experiments were carried out. A complicated spectrum was recorded when 5,5-dimethyl-1-pyrroline N-oxide (DMPO) was added to the reaction of toluene and DTBP [Figure 2(a)]. Moreover, with I_2 or both I_2

Published on 26 September 2014. Downloaded by Florida State University on 29/09/2014 17:36:24

Published on 26 September 2014. Downloaded by Florida State University on 29/09/2014 17:36:24

Journal Name

and 4-methylphenyl disulfide added to above system, similar radical signal was recorded [Figure 2(b, c)]. This result excluded I_2 and disulfide from the radical signal source. When toluene was replaced by cyclohexane, a different signal was recorded [Figure 2(d)], which indicated that the radical signal was related to the C(sp³)-H bonds, and thus the C(sp³) radical generated from the C(sp³)-H bonds may be the intermediate of the reaction.



Figure 1. The sampling experiment of reaction mixture of 1a (4 mL), 2a (0.3 mmol), DTBP (1.5 mmol) at 120 °C with and without I₂ respectively.



Scheme 5. Control experiments and Radical trapping experiment



Figure 2. The electron paramagnetic resonance (EPR) spectra (X band, 9.4GHz, RT) of (a) reaction mixture of DTBP (1.5 mmol) in toluene (2 mL) at 120 °C for 5 h with the addition of DMPO; (b) Reaction mixture of DTBP (1.5 mmol) and I₂ (0.045 mmol) in toluene (2 mL) at 120 °C for 5 h with the addition of DMPO; (c) Reaction mixture of DTBP (1.5 mmol), I₂ (0.045 mmol) and 4-methylphenyl disulfide (0.2 mmol) in toluene (2 mL) at 120 °C for 5 h with the addition of DMPO; (d) Reaction mixture of DTBP in cyclohexane (2 mL) at 120 °C for 5 h with the addition of DMPO.

Furthermore, an intermolecular kinetic isotopic effect (KIE) experiment was carried out, and a noteworthy KIE was observed with the $k_{\rm H}/k_{\rm D} = 3.0$ (Scheme 6). This result indicated that the C(sp³)-H bond activation might be involved in the rate determining step.



Scheme 6. KIE experiment

Based on the experiment results and literature precedent, a plausible mechanism was proposed. Initially, an oxidative dimerization of aryl mercaptans took place quickly to give corresponding disulfide with the I₂ as the catalyst and DTBP as the oxidant. The homocoupling of the sulfur radical, which generated from the S-H bond abstraction by tert-butoxy radical, were believed to be the route¹⁴. The tert-butoxy radical could also abstracted hydrogen from methyl arene to obtained benzyl radical with a lower rate that it did not occur before the completely consumption of the mercaptan. This step was also considered to be the rate determining step (RDS). Afterward, there are two possible pathways. Similar to organic peroxides, the disulfide might act as a sulfur radical pool that it release the sulfur radical slowly through homolysis¹⁶. And the sulfur radical couple with the benzyl radical to afford the product. Another possibility is that radical-substitution occurred between benzyl radical and disulfide^{6b, 10, 17} to give the product and the sulfur radical, which dimerized quickly to regenerate the disulfide.

In summary, we have established an oxidative $C(sp^3)$ -H/S-H coupling of alkanes with mercaptans to afford aryl alkyl sulfides with the employment of cheap and nontoxic I_2 and DTBP in moderate to good yields. This proposal provided a novel process to construct $C(sp^3)$ -S bond using alkanes and mercaptans directly, which eliminate the pre-functionalization of the starting materials and might encourage the further application. The mechanism study suggested that the disulfides were the intermediate and the $C(sp^3)$ -H bond cleavage was the rate-determing step of the transformation.

Acknowledgement

This work was supported by the 973 Program (2011CB80860, 2012CB725302), the National Natural Science Foundation of China (21390400, 21025206, 21272180 and 21302148), and the Research Fund for the Doctoral Program of Higher Education of China (20120141130002) and the Program for Changjiang Scholars and Innovative Research Team in University (IRT1030). The Program of Introducing Talents of Discipline to Universities of China (111 Program) is also appreciated.

Notes and references

^a College of Chemistry and Molecular Sciences, Wuhan University, Wuhan, Hubei 430072, P. R. China. *Fax:* (+86)-27-68754067; *Tel:* (+86)-27-68754672; *E-mail:* <u>aiwenlei@whu.edu.cn</u>;

^b National Research Center for Carbohydrate Synthesis, Jiangxi Normal University, Nanchang, Jiangxi, 330022, P. R. China.

†Electronic Supplementary Information (ESI) available: [details of any supplementary information available should be included here]. See DOI: 10.1039/c000000x

- (a) V. P. Ananikov, S. S. Zalesskiy and I. P. Beletskaya, Curr. Org. 1 • Synth. 2011, 8, 2-52; (b) I. P. Beletskaya and V. P. Ananikov, Chem. Rev. 2011, 111, 1596-1636; (c) C.-F. Lee, Y.-C. Liu and S. S. Badsara, Chem. - Asian J. 2014, 9, 706-722.
- 2 . (a) G. Bastug and S. P. Nolan, J. Org. Chem. 2013, 78, 9303-9308; (b) M. A. Fernandez-Rodriguez, Q. Shen and J. F. Hartwig, J. Am. Chem. Soc. 2006, 128, 2180-2181; (c) H.-J. Xu, Y.-Q. Zhao, T. Feng and Y.-S. Feng, J. Org. Chem. 2012, 77, 2878-2884; (d) R. B. N. Baig and R. S. Varma, Chem. Commun. 2012, 48, 2582-2584.
- (a) Q. Liu, R. Jackstell and M. Beller, Angew. Chem. Int. Ed. 2013, 3. 52, 13871-13873; (b) C. Zhang, C. Tang and N. Jiao, Chem. Soc. Rev. 2012, 41, 3464-3484; (c) C. Liu, H. Zhang, W. Shi and A. Lei, Chem. Rev. 2011, 111, 1780-1824.
- 4 (a) L. Zhou and W. Lu, Chem. Eur. J. 2014, 20, 634-642; (b) G. Song, F. Wang and X. Li, Chem. Soc. Rev. 2012, 41, 3651-3678; (c) A. E. Wendlandt, A. M. Suess and S. S. Stahl, Angew. Chem. Int. Ed. 2011, 50, 11062-11087; (d) Z. Shao and F. Peng, Angew. Chem. Int. Ed. 2010, 49, 9566-9568.
- 5 (a) Y. Yang, W. Dong, Y. Guo and R. M. Rioux, Green Chem. 2013, 15, 3170-3175; (b) S.-i. Fukuzawa, E. Shimizu, Y. Atsuumi, M. Haga and K. Ogata, Tetrahedron Lett. 2009, 50, 2374-2376; (c) X. Chen, X.-S. Hao, C. E. Goodhue and J.-Q. Yu, J. Am. Chem. Soc. 2006, 128, 6790-6791; (d) C.-L. Yi, Y.-T. Huang and C.-F. Lee, Green Chem. 2013, 15, 2476-2484; (e) J.-n. Xiang, Y.-q. Yuan and S.-r. Guo, Synlett 2013, 24, 443-448; (f) T. Uno, T. Inokuma and Y. Takemoto, Chem. Commun. 2012, 48, 1901-1903.
- (a) Y. Liao, P. Jiang, S. Chen, H. Qi and G.-J. Deng, Green Chem. 6 • 2013, 15, 3302-3306; (b) W. Ali, S. Guin, S. K. Rout, A. Gogoi and B. K. Patel, Adv. Synth. Catal. 2014, 10.1002/adsc.201400360.
- 7 · (a) B. L. Tran, B. Li, M. Driess and J. F. Hartwig, J. Am. Chem. Soc. 2014, 136, 2555-2563; (b) A. P. Antonchick and L. Burgmann, Angew. Chem. Int. Ed. 2013, 52, 3267-3271; (c) Z. Li, Y. Zhang, L. Zhang and Z.-Q. Liu, Org. Lett. 2013, 16, 382-385; (d) P. Xie, Y. Xie, B. Qian, H. Zhou, C. Xia and H. Huang, J. Am. Chem. Soc. 2012, 134, 9902-9905; (e) L. Zhou, S. Tang, X. Qi, C. Lin, K. Liu, C. Liu, Y. Lan and A. Lei, Org. Lett. 2014, 16, 3404-3407; (f) T. Hoshikawa and M. Inoue, Chem. Sci. 2013, 4, 3118-3123; (g) J. R. Suarez and J. L. Chiara, Chem. Commun. 2013, 49, 9194-9196; (h) D. Liu, C. Liu, H. Li and A. Lei, Angew. Chem., Int. Ed. 2013, 52, 4453-4456; (i) Y. Zhu and Y. Wei, Chem. Sci. 2014, 5, 2379-2382; (j) X. Guo and C.-J. Li, Org. Lett. 2011, 13, 4977-4979; (k) D. Liu, C. Liu, H. Li and A. Lei, Chem. Commun. 2014, 50, 3623-3626.
- 8 (a) R. Samanta, K. Matcha and A. P. Antonchick, Eur. J. Org. Chem. 2013, 2013, 5769-5804; (b) R. Rohlmann and O. Garcia Mancheno, Synlett 2013, 24, 6-10; (c) S. Wang, C. C. Shu, T. Wang, J. Yu and G. B. Yan, Chin. Chem. Lett. 2012, 23, 643-646; (d) M. Jiang, H.-J. Yang, Y. Li, Z.-Y. Jia and H. Fu, Chin. Chem. Lett. 2014, 25, 715-719; (e) Z. Li, C. Zhang, L. Zhu, C. Liu and C. Li, Org. Chem. Front. 2014, 1, 100-104.
- (a) M. Uyanik and K. Ishihara, ChemCatChem 2012, 4, 177-185; 9. (b) S. Tang, Y. Wu, W. Liao, R. Bai, C. Liu and A. Lei, Chem. Commun. 2014, 50, 4496-4499; (c) Y. Lv, Y. Li, T. Xiong, Y. Lu,

Q. Liu and Q. Zhang, Chem. Commun. 2014, 50, 2367-2369; (d) U. Kloeckner, P. Finkbeiner and B. J. Nachtsheim, J. Org. Chem. 2013, 78, 2751-2756.

- 10 When this article was under review, a work utilizing methyl arenes and disulfides as the coupling reagents to afford thioesters was published online. The different results between our work and this paper may be ascribe to the different concentration of DTBP: S. S. Badsara, Y.-C. Liu, P.-A. Hsieh, J.-W. Zeng, S.-Y. Lu, Y.-W. Liu and C.-F. Lee, Chem. Commun. 2014, 50, 11374-11377.
- 11 D. A. Pratt, J. H. Mills and N. A. Porter, J. Am. Chem. Soc. 2003, 125, 5801-5810.
- 12 · F. Agapito, P. M. Nunes, B. J. Costa Cabral, R. M. Borges dos Santos and J. A. Martinho Simões, J. Org. Chem. 2008, 73, 6213-6223
- 13 A. Cherkasov and M. Jonsson, J. Chem. Inf. Comput. Sci. 2000, 40, 1222-1226
- 14 M. Kirihara, Y. Asai, S. Ogawa, T. Noguchi, A. Hatano and Y. Hirai, Synthesis 2007, 2007, 3286-3289.
- 15 X. Liu, G. Yu, J. Li, D. Wang, Y. Chen, K. Shi and B. Chen, Synlett 2013, 24, 1588-1594.
- 16 (a) A. M. Mak, R. Steudel and M. W. Wong, Chemistry An Asian Journal 2008, 3, 1026-1034; (b) V. Subramanian and K. Seff, J. Am. Chem. Soc. 1980, 102, 1881-1884.
- 17 (a) J. Zhao, H. Fang, J. Han, Y. Pan and G. Li, Adv. Synth. Catal. 2014, 356, 2719-2724; (b) S.-r. Guo, Y.-q. Yuan and J.-n. Xiang, Org. Lett. 2013, 15, 4654-4657; (c) R.-Y. Tang, Y.-X. Xie, Y.-L. Xie, J.-N. Xiang and J.-H. Li, Chem. Commun. 2011, 47, 12867-12869; (d) B. Du, B. Jin and P. Sun, Org. Lett. 2014, 16, 3032-3035; (e) S.-r. Guo, W.-m. He, J.-n. Xiang and Y.-q. Yuan, Chem. Commun. 2014, 50, 8578-8581.

Published on 26 September 2014. Downloaded by Florida State University on 29/09/2014 17:36:24

$C(sp^3)$ + H + H - S - Ar $\xrightarrow{I_2/DTBP}$ $C(sp^3)$ - S - Ar

19 examples, up to 90%

The $C(sp^3)$ -S bond formation was achieved utilizing $C(sp^3)$ -H and S-H as the nucleophiles. Methyl arenes, cycloalkanes and aliphatic ketones exhibited reactivity for this transformation. Mechanism study revealed that $C(sp^3)$ radical and disulfide were the intermediates of the reaction.