

Tetrahedron 54 (1998) 5709-5720

TETRAHEDRON

Diastereoselective Addition of α -Hydroxyalkyl and α -Alkoxyalkyl Radicals to Chiral 4-Methyleneoxazolidin-5-ones

Stephen G. Pyne* and Karl Schafer

Department of Chemistry, University of Wollongong, Wollongong, NSW, 2522, Australia.

Received 14 January 1998; revised 17 March 1998; accepted 19 March 1998

Abstract: A method for preparing optically active homoserine derivatives via the photoinduced radical additions of alcohols and ethers to the chiral 4-methyleneoxazolidin-5-ones 1 and 2 has been achieved. These photoadducts, however, are readily prone to epimerization under the conditions required for their deprotection. © 1998 Elsevier Science Ltd. All rights reserved.

L-Homoserine and its derivatives are useful building blocks for the synthesis of unnatural amino acids, including γ -seleno-, γ -thio- and γ -phosphono-derivatives.^{1,2} Some of these compounds have interesting biological activities.³ In principle, homoserines can be obtained from the addition of α -hydroxyalkyl and α alkoxyalkyl radicals to α,β -didehydroamino acids (eq 1). Optically active versions of these target compounds could, in principle, be obtained if either the α -hydroxyalkyl and α -alkoxyalkyl radical were chiral or the α,β didehydroamino acid was chiral. Indeed, a readily available chiral auxiliary for α -hydroxyalkyl radicals has been recently developed by Garner.⁴ In this paper we report our studies aimed at preparing optically active homoserine derivatives from the photoinduced radical additions of alcohols and ethers to the chiral, cyclic α,β -didehydroamino acid derivatives, the (2S)⁵ and (2R)⁶ 4-methyleneoxazolidin-5-ones 1 and 2, respectively.

(eq 1)



While the photochemical induced addition of alcohols to α,β -unsaturated carbonyl⁷ and sulfur⁸ compounds has been well documented, reports of these reactions involving α,β -didehydroamino acids as

substrates have not been reported as far as we are aware. The addition of radicals to 1 has been reported by Beckwith.⁹ These reactions proceed with moderate to excellent diastereoselectivities, depending upon the nature of the radical, and favoured formation of *trans* 2,4-disubstituted-oxazolidinones (eq 2).



Results and Discussion

In some preliminary studies on the photolytic stability of 1 and 2 upon irradiation in acetone we found that both 1 and 2 gave numerous products after irradiation with a medium pressure mercury lamp (140 W) for 1 h. Only in the case of 2 could any discrete product be isolated, that being the photocyclization product 3 in 10 % isolated yield. The formation of similar tricyclic products from the photocyclization of related *N*-arylenaminoketones¹⁰ and lactones¹¹ is well known. The stereochemistry of 3 was evident from NOESY experiments that showed cross peaks between the signals for the ortho proton H₀ and H9a.



Photoinduced Reactions of 1 and 2 with Alcohols

Photolysis of a solution of 1 in neat alcohol (R^1R^2CHOH) in the presence of the sensitiser benzophenone (1 molar equiv.) for 1h at ambient temperature (water cooled system) gave mixtures of the photoadducts 4 and their cyclized products, the lactones 5 (Scheme 1). When methanol was employed as the solvent then ${}^{1}H$ NMR analysis of the crude photolysis mixture showed the formation of a photoadduct as a 77 : 23 mixture of diastereoisomers (major: δ 6.08 (s, H2), 1.05 (s, Bu^t); minor: δ 6.06 (s, H2), 0.94 (s, Bu^t)). We assume the major and minor diastereoisomers are 4a and its 2,4-cis diastereoisomer, respectively. Attempted purification of 4a on silica gel gave the lactone 5a in low yield (30%), however, none of the photoadduct 4a was isolated. Photolysis of a solution of 1 in ethanol or isopropanol gave the photoadducts 4b (d. r. = 58 : 42) and 4c (d. r. = >98; <2), respectively, from ¹H NMR analysis of the crude reaction mixtures. Purification of these crude photolysis mixtures on silica gel gave pure samples of 4b/5b and 4c/5c, respectively (Scheme 1). The 4R stereochemistry of both diastereoisomers of 4b and that of 4c was evident from NOESY experiments which showed NOE cross peaks between H4 and the C2 But group. The moderate yields obtained from these reactions prompted us to examine the direct hydrolysis/cyclization of the crude photolysis product from the reaction of 1 and isopropanol. Hydrolysis of the crude photoadduct 4c with 6N HCl at 90 °C for 12 h gave, after purification by recrystallization from ethanol/ether, the hydrochloride salt of the α -amino lactone (R)-6 in 90 % overall yield from 1 (Scheme 2). The enantiomeric purity of (R)-6, however, was 56 % based on 1 H NMR analysis of its Mosher amide (R,S)-7 (78: 22 mixture of diastereoisomers). The low ee of 6 indicated that epimerization at the α -carbon of 6 must have occurred during the acid hydrolysis conditions since 4c was essentially diastereomerically pure and the enantiomeric purity of 1 was >98 %.⁶ Unfortunately, less harsh hydrolysis conditions, in which epimerization would be less likely, did not completely hydrolyse the benzamide group of 5c.



Scheme 2



In contrast, the photolysis of 2 in ethanol or isopropanol lead directly to the lactones **5b** and **5c**, respectively in moderate yields (54 %, Scheme 3). The intermediate photoadducts A could not be detected in the crude photolysis product mixtures. The lactone **5b** was a 55 : 45 mixture of diastereoisomers. Acid hydrolysis of the lactone (S)-**5c** gave the hydrochloride salt of the amine (S)-**6** in 90 % (Scheme 4). The S stereochemistry of this salt was evident from the positive sign of its specific rotation which was opposite in sign to (R)-**6** described above in Scheme 2. The enantiomeric purity of (S)-**6** was 16 % from ¹H NMR analysis of its Mosher amide (R,S)-**7** a value much lower than that found for its enantiomer (R)-**6**. The low enantiomeric purity of (S)-**6** might be due to the low diastereoselectivity in the formation of A (R¹ = Me, R² = H), however this compound could not be detected by ¹H NMR and thus its d. r. could not be measured. However, the different magnitudes for the specific rotations of (R)-**6** ([α]_D²⁵ -12 (c 0.25, 6N HCl) and (S)-**6** ([α]_D²¹ +6 (c 0.1, 6N HCl)), suggested that the formation of A (R¹ = R² = Me) was less diastereoselective than the formation of its *tert*-butyl analogue **4c**.



Photoinduced Reactions of 1 and 2 with Ethers

The photolysis products from 1 and the cyclic ethers, 1,3-dioxolane and THF, are shown below in Scheme 5. The photochemical induced reaction of 1 with 1,3-dioxolane gave the photoadduct 8a as a single diastereoisomer in 52 % yield after column chromatography. While that with THF gave a 57 : 43 mixture of diastereoisomeric photoadducts 8b in 55 % yield (Scheme 5). All three adducts (8a and the two diastereoisomers of 8b) had the 2,4-*trans* stereochemistry from NOESY experiments (Scheme 5). The enantiomeric purity of 8a was > 96 % from ¹H NMR analysis of its ester derivative 9 that was prepared from the reaction of 8a and methyl (*R*)-mandelate under mild based catalysed conditions.



Photoinduced reactions of the acyclic ethers, dimethoxymethane and *tert*-butyl methyl ether with 1 gave mixtures of the 2,4-*trans* and 2,4-*cis*-adducts 10 and 11 respectively in poor yields (Scheme 6), while the reaction of diethyl ether and *tert*-butyl ethyl ether with 1 gave a mixture of four and three diastereoisomeric

products, respectively. In these latter two reactions, the major adducts (10c and 10d, respectively) had the 2,4-trans stereochemistry of the oxazolidinone ring (from NOESY studies) but were formed as a mixture of

epimers at the exocyclic stereogenic centre (Scheme 6). A 95 : 5 mixture of **10b** and **11b** was hydrolysed with 6N HCl to give after purification (R)-**12**. The enantiomeric purity of this compound was estimated as 62 % based upon its specific rotation ($[\alpha]_D^{20}$ -20 (c 0.1, 6N HCl) when compared to the literature value ($[\alpha]_D$ -32 (c 0.2, 2N HCl)).¹²

Scheme 6



lit. (ref. 12), [α]_D -32 (c 0.2, 2N HCl)

The two 2,4-*trans* diastereoisomers of **10d** could be separated by column chromatography and the individual isomers were subjected to acid catalysed hydrolysis (Scheme 7). The major diastereoisomer of **10d** gave mainly the *trans*-lactone **13** from NOESY experiments that showed cross peaks between the resonances for the γ -methyl group and the α -proton. The minor diastereoisomer of **10d** gave mainly the *cis*-lactone **14** Scheme **7**



that did not show any NOE between the γ -methyl group and the α -proton. Each hydrolysis reaction, however gave a mixture of the two diastereoisomers 13 and 14 and the ratio of these diastereoisomers changed with time as monitored by ¹ H NMR. For example, immediately after the hydrolysis of the minor diastereoisomer of 10d the ratio of 14 : 13 was 91 : 9, however after the NMR sample had stood for 3 days the ratio had changed to 74 : 26. The ¹H NMR data for (*R*, *R*)-13 and (*S*, *R*)-14 from this study closely matched that reported in the literature for (*S*, *S*)-13¹³⁻¹⁵ or (*R*, *R*)-13^{13,16,17} and (*S*, *R*)-14,^{14, 16-18} respectively.¹⁹

In conclusion, a method for preparing optically active homoserine derivatives via the photoinduced radical additions of alcohols and ethers to the chiral 4-methyleneoxazolidin-5-ones 1 and 2 has been achieved. These photoadducts, however, are prone to epimerization under the conditions required for their deprotection. Other protecting groups at the oxazolidinone nitrogen (e.g. Boc) may allow the synthesis of these compounds in higher enantiomeric purities.²⁰

Experimental

General procedures were as described previously.⁶ All NMR were determined in CDCl₃ solution, unless otherwise indicated, at 300 MHz (¹H NMR) or 77.5 MHz (¹³C NMR). When D₂O was employed as the NMR solvent then 1,4-dioxane (67.8 ppm for ¹³C) was used as an internal reference. Photolyses were carried out in a water cooled pyrex immersion-well reactor with the light from a 140 W mercury lamp. Prior to photolysis, nitrogen was bubbled through the reaction mixture for 0.5 h.

(3R,9aS)-5-oxo-3-phenyloxazolidino[3,4-b]-3aH-4-oxo-9,9a-dihydroisoquinoline (3). The oxazolidinone (2) (100 mg, 0.36 mmol) was dissolved in dry acetone (150ml) and photolysed under an atmosphere of nitrogen for 1 h. After removal of the solvent under reduced pressure the crude residue was chromatographed on silica gel, eluting with 10% ethyl aceate in light petroleum, to afford 3 as a white solid (10 mg, 10%), m.p. 129-132°C (Found : C, 73.3; H, 4.5; N, 4.9%. C17H13NO3 requires C, 73.1; H, 4.7; N, 5.0%). [α]²⁵D -90° (c, 0.08 in CHCl₃). ¹H n.m.r. δ 3.26, app. t, J 12.9 Hz, 1H, H-9α; 3.37, dd, J 15.6, 6.0 Hz, 1H, H-9β; 4.62, dd, J 12.9, 6.0 Hz, 1H, H-9a; 7.18, s, 1H, H-3; 7.31, d, J 7.5 Hz, 1H, H-8; 7.40-7.42, m, 4H, ArH; 7.51-7.55, m, 3H, ArH; 8.01, dd, J 7.5, 1.5 Hz, 1H, H-5.¹³C n.m.r. δ 29.68, CH₂; 52.17, CH; 88.45, CH; 125.85, CH; 127.78, CH; 128.07, CH; 128.17, CH; 128.60, C; 128.93, CH; 129.85, CH; 133.05, CH; 134.61, C; 135.66, C; 160.95, CO; 171.31, CO. Mass spectrum m/z 280 (M + H⁺, 65%), 146 (65), 60 (100).

General Method: Photolysis of (2S)-3-Benzoyl-2-tert-butyl-4-methyleneoxazolidin-5-one (1) in Isopropyl Alcohol. A mixture of the oxazolidinone 1 (100 mg, 0.39 mmol) and benzophenone (70 mg, 0.39 mmol) in dry isopropyl alcohol (150ml) was photolysed under an atmosphere of nitrogen for 1 h. The reaction mixture was concentrated under reduced pressure and chromatographed on silica gel, eluting with 10% ether in light petroleum, to afford 2 compounds. The first compound was isolated as a colourless oil and identified as (2S,4R)-3-benzoyl-2-tert-butyl-4-(2'-hydroxy-2'-methylpropyl)oxazolidin-5-one (4c) (27 mg) in 23% yield. (Found : M^{+•} + H⁺, 320.1827. C18H26NO4 requires 320.1862). $[\alpha]^{22}D^{-30}$ (c, 0.20 in CHCl3). ¹H n.m.r. δ 0.95, s, 9H; 1.19, s, 3H, CH3; 1.21, s, 3H, CH3; 2.00, dd, J 14.7, 3.0 Hz, 1H, H-1' α ; 2.20, dd, J 14.7, 9.6 Hz, 1H, H-1' β ; 3.53, br. s, 1H, OH; 4.57, dd, J 9.9, 3.0 Hz, 1H, H-4; 6.11, s, 1H, H-2; 7.35-7.56, m, 5H, ArH. ¹³C n.m.r. δ 25.03, CH3; 28.47, CH3; 30.41, CH3; 37.11, C; 46.80, CH2; 54.84, CH; 68.79, C; 96.11, CH; 126.51, CH; 128.94, CH; 130.50, CH; 135.44, C; 173.18, CO; 174.22, CO. Mass spectrum m/z 320 (M + H⁺, 63%), 234 (100). The second compound was obtained as a white solid and identified as (**R**)- α -benzamido- $\gamma\gamma$ +

dimethyl-7-butyrolactone (5c) (33 mg, 36%), m.p. 150-153°C (Found : C, 66.7; H, 6.6; N, 5.6%. C_{13H15}NO₃ requires C, 66.9; H, 6.5; N, 6.0%). $[\alpha]^{22}D + 12°$ (c, 0.30 in CHCl₃). ¹H n.m.r. δ 1.40, s, 3H, CH₃; 1.45, s, 3H, CH₃; 2.03, app. t, J 12.3 Hz, 1H, H- β a; 2.66, dd, J 12.6, 9.0 Hz, 1H, H- β b; 4.91-5.02, m, 1H, H- α ; 7.10, br. s, 1H, NH; 7.27-7.71, m, 5H, ArH. ¹³C n.m.r. δ 26.99, CH₃; 28.87, CH₃; 41.99, CH₂; 50.59, CH; 83.08, C; 127.11, CH; 128.49, CH; 131.89, CH; 132.94, C; 167.47, CO; 175.27, CO. Mass spectrum *m*/z 234 (M + H⁺, 100%), 122 (48), 105 (48).

Photolysis of (2R)-3-Benzoyl-4-methylene-2-phenyloxazolidin-5-one (2) in Isopropyl Alcohol. The photolysis of 2 (100 mg, 0.36 mmol) in isopropyl alcohol (150 ml) and benzophenone (65 mg, 0.36 mmol) was carried out as described above. Column chromatography on silica gel, eluting with 10% ether in light petroleum, afforded (S)- α -benzamido- γ , γ -dimethyl- γ -butyrolactone (5c) as the sole product (45 mg, 54%), isolated as a white solid, m.p. 155-158°C. $[\alpha]^{22}D$ - 11° (c, 0.10 in CHCl3). The spectral data of this compound were identical to that reported above for (R)-5c.

Photolysis of (2S)-3-Benzoyl-2-tert-butyl-4-methyleneoxazolidin-5-one (1) in Ethanol. The photolysis of 1 (100mg, 0.36 mmol) in ethanol (150 ml) was carried out as above. Column chromatography, eluting with 10% ether in light petroleum, afforded 2 fractions. Fraction 1 yielded a colourless oil (27 mg, 23%) which was identified as (2S,4R)-3-benzoyl-2-tert-butyl-4-(2'-hydroxypropyl)oxazolidin-5-one (4b), isolated as an inseparable mixture of two diastereomers (58:42). Major isomer: (Found : M⁺⁺ + H⁺, 306.1716. C₁₇H₂₄NO4 requires 306.1705). ¹H n.m.r. δ 0.99, s, 9H; 1.09, d, J 6.3 Hz, 3H, CH₃; 1.87, ddd, J 14.1, 9.9, 3.6 Hz, 1H, H-1'α; 2.10, ddd, J 13.8, 9.6, 2.7 Hz, 1H, H-1'β; 3.60, br. s, 1H, OH; 3.96-4.06, m, 1H, H-2'; 4.45, dd, J 9.9, 3.6 Hz, 1H, H-4; 6.11, s, 1H, H-2; 7.32-7.49, m, 5H, ArH. ¹³C n.m.r. δ 23.53, CH3; 25.12, CH3; 36.99, C; 43.53, CH2; 54.98, CH; 64.58, CH; 95.74, CH; 126.76, CH; 128.76, CH; 130.66, CH; 135.38, C; 172.83, CO; 173.67, CO. Mass spectrum m/z 306 (M + H⁺, 100%), 237 (50), 220 (95), 105 (80). Minor isomer: ¹H n.m.r. δ 0.92, s, 9H; 1.11, d, J 6.3 Hz, 3H, CH3; 1.53, ddd, J 8.7, 6.3, 6.3 Hz, 1H, H-1'α; 2.00, ddd, J 5.7, 5.7, 3.3 Hz, 1H, H-1'β; 3.60, br. s, 1H, OH; 3.96-4.06, m, 1H, H-2'; 4.55, dd, J 9.0, 5.7 Hz, 1H, H-4; 6.08, s, 1H, H-2; 7.32-7.49, m, 5H, ArH. ¹³C n.m.r. δ 22.98, CH₃; 24.81, CH₃; 36.01, C; 43.42, CH₂; 56.57, CH; 64.96, CH; 96.36, CH; 126.55, CH; 128.90, CH; 130.51, CH; 133.50, C; 172.83, CO; 173.67, CO. Fraction 2 yielded a colourless oil (26 mg, 30%) which was identified to be the lactone (**R**)- α -benzamido- γ -methyl- γ butyrolactone (5b), isolated as an inseparable mixture of two diastereomers (55:45). (major isomer): (Found : M^{+•} + H⁺, 220.0977. C₁₂H₁₄NO3 requires 220.0974). ¹H n.m.r. δ 1.46, d, J 6.6 Hz, 3H, CH3; 2.40, ddd, J 12.6, 10.2, 8.4 Hz, 1H, H-βa; 2.57, ddd, J 12.6, 9.6, 2.1 Hz, 1H, H-βb; 4.77-4.92, m, 2H, H-α & H-γ; 7.19, br. d, J 6.0 Hz, 1H, NH; 7.38-7.79, m, 5H, ArH. ¹³C n.m.r. δ 21.21, CH3; 35.79, CH2; 48.89, CH; 75.30, CH; 127.33, CH; 128.39, CH; 132.02, CH; 133.89, C; 167.67, CO; 175.57, CO. Mass spectrum m/z 220 (M + H⁺, 100%), 105 (95). (minor isomer) : ¹H n.m.r. δ 1.48, d, J 6.6 Hz, 3H, CH3; 1.99, ddd, J 12.3, 12.3, 12.0 Hz, 1H, H-βa; 3.01, ddd, J 12.6, 8.4, 5.1 Hz, 1H, H-βb; 4.65, ddd, J 10.8, 6.0, 5.1 Hz, 1H, H-γ; 4.77-4.92, m, 1H, H- α ; 7.11, br. d, J 6.0 Hz, 1H, NH; 7.38-7.79, m, 5H, ArH. ¹³C n.m.r. δ 20.59, CH₃; 38.39, CH₂; 51.31, CH; 75.08, CH; 127.13, CH; 128.59, CH; 132.02, CH; 131.44, C; 167.22, CO; 175.37, CO.

Photolysis of (2R)-3-Benzoyl-4-methylene-2-phenyloxazolidin-5-one (2) in Ethanol. The photolysis of 2 in ethanol was carried out as described above. Column chromatography on silica gel, eluting with 10% ether in light petroleum, afforded a colourless oil which was identified to be an inseparable mixture of two lactone

diastereomers (S)-(5b) in a ratio of 55:45 (43 mg, 54%). The spectral data for the 2 diastereomers was identical to that reported above for (R)-(5b).

Photolysis of (2S)-3-Benzoyl-2-*tert***-butyl-4-methyleneoxazolidin-5-one (1) in Methanol.** The photolysis of 1 in methanol was carried out as above. ¹H n.m.r. analysis of the crude residue showed this to be a mixture of two diastereomers of **(2S,4R)***-3-benzoyl-2-tert-butyl-4-(2'-hydroxyethyl)oxazolidin-5-one* **(4a)** in a ratio of approximately 77 : 23 (H-2 at δ 6.08 and 6.06 ppm, respectively). Column chromatography, eluting with 10% ether in light petroleum, gave only the lactone **(R)***-α-benzamido-γ-butyrolactone* **(5a)** (24 mg, 30%) as a colourless gum. $[\alpha]^{22}D - 20^{\circ}$ (c, 0.10 in CHCl3). (Found : M⁺⁺ + H⁺, 206.0811. C11H12NO3 requires 206.0817). ¹H n.m.r. δ 2.28, app. dq, J 11.4, 6.5, 6.5, 6.5 Hz, 1H, H-βa; 2.92, dddd, J 14.1, 8.4, 5.7, 1.0 Hz, 1H, H-βb; 4.34, ddd, J 11.1, 9.3, 5.7 Hz, 1H, H-α; 4.52, ddd, 9.0, 9.0, 1.0 Hz, 1H, H-βa; 4.77, ddd, J 11.7, 8.7, 6.0 Hz, 1H, H-γb; 6.98, br. d, J 6.0 Hz, 1H, NH; 7.38-7.79, m, 5H, ArH. ¹³C n.m.r. δ 30.44, CH2; 49.68, CH; 66.22, CH2; 127.14, CH; 128.64, CH; 132.09, CH; 133.00, C; 167.77, CO; 175.67, CO. Mass spectrum *m/z* 206 (M + H⁺, 100%), 105 (95).

Enantiomeric Excess Determination of the Lactone (R)-(6c). The crude residue from the photolysis of 1 (100 mg, 0.39 mmol) in isopropyl alcohol was heated at 90 °C in 6M hydrochloric acid (10 ml) for 12 h. The reaction mixture was cooled and washed with dichloromethane (4 x 10 ml). The aqueous fraction was evaporated to dryness. Purification of the residue by recrystallization from ethanol/ether afforded the hydrochloride salt of (R)-c-amino-y-dimethyl-y-butyrolactone ((R)-6) (58 mg, 90%) as a cream solid, m.p. 185-191°C (dec) (Found : C, 40.8; H, 7.5; N, 7.8%. C6H12NO2Cl requires C, 40.5; H, 7.3; N, 7.5%). [α]²²D -12° (c, 0.25 in 6M HCl). ¹H n.m.r. (D₂O) δ 1.40, s, 3H, CH₃; 1.48, s, 3H, CH₃; 2.19, dd, J 12.3, 12.3 Hz, 1H, H-βa; 2.69, dd, J 12.9, 9.3 Hz, 1H, H-βb; 4.59, dd, J 12.0, 9.3 Hz, 1H, H-α. ¹³C n.m.r. (D₂O) δ 27.02, CH3; 28.94, CH3; 39.57, CH2; 50.95, CH; 87.45, C; 174.49, CO. Mass spectrum m/z 130 (M - HCl + H⁺, 100%). A stirred sample of (R)-6 (20 mg, 0.12 mmol) suspended in dry dichloromethane (1 ml) was treated with triethylamine (25 mg, 0.25 mmol) under a nitrogen atmosphere at room temperature. After 10 minutes, 4dimethylaminopyridine (several crystals) and (R)-Mosher chloride (37 mg, 0.14 mmol) were added sequentially. After 3 h, the reaction mixture was diluted with ether (10 ml) and subsequently washed with 3M HCl (5 ml), 10% aqueous sodium carbonate (5 ml) and water (5 ml). The ethereal layer was dried, filtered and concentrated under reduced pressure to afford a pale yellow oil (24 mg, 58%). ¹H n.m.r. analysis of the crude residue showed this to be a mixture of two diastereomers in a ratio of approximately 78:22. Major isomer: (Found : $M^{+*} + H^+$, 346.1264. C16H19NO4F3 requires 346.1266). ¹H n.m.r. δ 1.44, s, 3H, CH3; 1.48, s, 3H, CH3; 2.57, dd, J 12.9, 6.9 Hz, 1H, H-Ba; 2.63, dd, J 12.6, 9.0 Hz, 1H, H-Bb; 3.47, d, J 1.5 Hz, 3H, OCH3; 4.75, ddd, J 11.7, 8.7, 6.6 Hz, 1H, H-α; 7.38-7.79, m, 6H, ArH & NH. ¹³C n.m.r. δ 26.97, CH3; 28.86, CH3; 41.25, CH₂; 50.35, CH₃; 55.11, C; 55.14, CH₃; 55.16, C; 82.84, C; 127.47, CH; 128.59, CH; 129.63, CH; 132.23, C; 167.11, CO; 173.53, CO. Mass spectrum m/z 346 (M + H⁺, 30%). Minor isomer: ¹H n.m.r. δ 1.45, s, 3H, CH3; 1.52, s, 3H, CH3; 1.97, dd, J 12.6, 7.4 Hz, 1H, H-βa; 2.72, dd, J 12.3, 8.4 Hz, 1H, H-βb; 3.34, d, J 1.2 Hz, 3H, OCH3; 4.86, ddd, J 11.7, 8.7, 7.2 Hz, 1H, H-α; 7.38-7.79, m, 6H, ArH & NH. ¹³C n.m.r. δ 26.83, CH3; 28.80, CH3; 41.74, CH2; 50.12, CH3; 54.85, C; 54.87, CH3; 55.08, C; 82.89, C; 127.92, CH; 128.72, CH; 129.69, CH; 131.47, C; 167.03, CO; 173.77, CO.

Enantiomeric Excess Determination of the Lactone (S)-5c. By following the above procedure the lactone (S)-5c (120 mg, 0.52 mmol) was treated with 6M hydrochloric acid (10 ml) to give the hydrochloride salt of

(S)- α -amino- γ -dimethyl- γ -butyrolactone ((S)-6) as a white solid (77 mg, 90%), m.p. 193-197°C (dec). [α]²²_D +6° (c, 0.10 in 6M HCl). The spectral data of (S)-6 were identical to that found for (R)-6. Mosher analysis of (S)-6 (20 mg, 0.12 mmol) was subsequently carried out according to the procedure described above (yield = 70%). ¹H n.m.r. analysis of the crude residue showed this to be a mixture of two diastereomers in a ratio of approximately 58:42.

Photolysis of (2S)-3-Benzoyl-2-tert-butyl-4-methyleneoxazolidin-5-one (1) in 1,3-Dioxolane. The photolysis of 1 (100 mg, 0.39 mmol) in 1,3-dioxolane (150 ml) was carried out as described above. Column chromatography on silica gel, eluting with 10% ether in light petroleum, afforded (2S,4R)-3-benzoyl-2-tert-butyl-4-([1',3'-dioxan-2'-yl]methyl)oxazolidin-5-one (8a) (66 mg, 52%) as a white solid, m.p. 110-113°C (Found : C, 64.9; H, 7.1; N, 4.0%. C18H23NO5 requires C, 64.8; H, 7.0; N, 4.2%). $[\alpha]^{24}D + 15^{\circ}$ (c, 0.10 in CHCl3). ¹H n.m.r. δ 1.03, s, 9H; 2.00, ddd, J 13.5, 7.2, 3.3 Hz, 1H, CH_aH_b; 2.28, ddd, J 13.8, 10.5, 3.6 Hz, 1H, CH_aH_b; 2.00, app. dt, J 12.9, 12.9, 6.3 Hz, 1H, H-4'α or H-5'α; 3.66, app. dt, J 12.9, 12.9, 6.9 Hz, 1H, H-4'β or H-5'β; 3.74, app. dt, J 12.9, 12.9, 6.3 Hz, 1H, H-4'α or H-5'α; 3.83, app. dt, J 12.9, 12.9, 6.9 Hz, 1H, H-4'β or H-5'β; 4.35, dd, J 10.5, 3.3 Hz, 1H, H-4; 4.97, dd, J 7.5, 3.6 Hz, 1H, H-2'; 6.11, s, 1H, H-2; 7.35-7.47, m, 5H, ArH. ¹³C n.m.r. δ 25.13, CH3; 36.97, C; 39.23, CH2; 54.37, CH; 64.53, CH2; 64.78, CH2; 95.68, CH; 100.66, CH; 126.66, CH; 128.70, CH; 130.39, CH; 135.46, C; 171.89, CO; 173.67, CO. Mass spectrum *m*/z 334 (M + H⁺, 100%), 184 (48).

Enantiomeric Excess Determination of the Oxazolidinone 8a. A mixture of 8a (16 mg, 0.05 mmol), (*R*)methyl mandelate (9 mg, 0.055 mmol) and triethylamine (6 mg, 0.06 mmol) in dry dichloromethane (1 ml) was stirred at room temperature for 3 days under a nitrogen atmosphere. The reaction mixture was diluted with dichloromethane (5 ml) and washed with 2M HCl (1 x 5ml) and water (3 x 5 ml). The organic fraction was dried, filtered and evaporated to dryness. ¹H n.m.r. analysis of the crude residue showed this to be a single diastereomer (9). Column chromatography on silica gel, eluting with 30% ethyl acetate in light petroleum, yielded a pale yellow gum (15 mg, 73%). (Found : $M^{+o} + H^+$, 346.1276. C16H19NO4 F3 requires 346.1266). ¹H n.m.r. δ 2.21, ddd, J 15.0, 4.8, 4.8 Hz, 1H, H- α ; 2.35, ddd, J 15.0, 6.3, 3.6 Hz, 1H, H- β ; 3.68, s, 3H, CH3; 3.79-3.96, m, 4H, 2 x CH2; 4.80, dd, J 11.4, 6.3 Hz, CH; 5.01, app. t, J 4.5 Hz, 1H, CH; 5.91, s, 1H, CH; 7.29-7.60, m, 11H, ArH & NH. ¹³C n.m.r. δ 34.17, CH2; 52.95, CH3; 53.00, CH; 64.97, CH2; 72.90, CH; 102.38, CH; 126.56, CH; 127.15, CH; 128.47, CH; 128.58, CH; 128.60, CH; 131.91, CH; 133.37, C; 138.21, C; 167.84, CO; 173.76, CO; 174.09, CO. Mass spectrum *m/z* 346 (M + H⁺, 66%).

Photolysis of (2S)-3-Benzoyl-2-tert-butyl-4-methyleneoxazolidin-5-one (1) in Tetrahydrofuran. The photolysis of 1 (100 mg, 0.39 mmol) in tetrahydrofuran (150 ml) was carried out as described above. Column chromatography on silica gel, eluting with 10% ether in light petroleum, afforded an inseparable mixture of two diastereomers of (2S)-3-benzoyl-2-tert-butyl-4-[tetrahydrofuran-2'-yl)methyl]oxazolidin-5-one (8b) (71 mg, 55%) as a white solid, m.p. 105-110°C (Found : C, 68.8; H, 7.6; N, 4.1%. C19H25NO4 requires C, 68.8; H, 7.6; N, 4.2%). Mass spectrum *m*/z 332 (M + H⁺, 100%), 246 (40), 182 (73). Major isomer: ¹H n.m.r. δ 1.03, s, 9H; 1.61-2.05, m, 5H, CH_aH_b, H-3'α/β & H-4'α/β; 2.00, ddd, J 13.9, 10.5, 5.7 Hz, 1H, CH_aH_b; 3.58, app. dt, J 6.6, 6.6 Hz, 1H, H-5'α; 3.74, app. dt, J 6.3, 6.3 Hz, 1H, H-5'β; 3.78-3.86, m, 1H, H-2'; 4.03, dd, J 10.8, 3.0 Hz, 1H, H-4; 6.08, s, 1H, H-2; 7.36-7.48, m, 5H, ArH. ¹³C n.m.r. δ 25.11, CH3; 25.32, CH2; 29.57, CH2; 36.95, C; 40.05, CH2; 55.64, CH; 67.35, CH2; 75.09, CH; 95.46, CH; 126.43, CH; 128.82, CH; 130.39, CH; 135.74, C; 171.86, CO; 173.55, CO. Minor isomer: ¹H n.m.r. δ 1.04, s, 9H; 1.41, ddd, J 15.3, 11.1, 6.9

Hz, 1H, CH_aH_b; 1.61-2.05, m, 5H, CH_aH_b, H-3'α/β & H-4'α/β; 3.28, app. dt, J 7.8, 7.8 Hz, 1H, H-5'α; 3.48, app. dt, J 6.9, 6.9 Hz, 1H, H-5'β; 4.07-4.12, m, 1H, H-2'; 4.37, dd, J 11.1, 3.3 Hz, 1H, H-4; 6.10, s, 1H, H-2; 7.36-7.48, m, 5H, ArH. ¹³C n.m.r. δ 25.16, CH₃; 25.35, CH₂; 31.30, CH₂; 36.87, C; 40.93, CH₂; 54.87, CH; 67.31, CH₂; 74.20, CH; 95.27, CH; 126.63, CH; 128.56, CH; 130.28, CH; 135.53, C;172.26, CO; 173.86, CO.

Photolysis of (2S)-3-Benzoyl-2-tert-butyl-4-methyleneoxazolidin-5-one (1) in Diethyl Ether. The photolysis of 1 (100 mg, 0.39 mmol) in diethyl ether (150 ml) was carried out as described above. Column chromatography on silica gel, eluting with 10% ether in light petroleum, afforded an inseparable mixture of 4 diastereomers (2S,4R)-3-benzoyl-2-tert-butyl-4-(2'-ethoxypropyl)oxazolidin-5-one (10c) and (2S,4S)-3benzoyl-2-tert-butyl-4-(2'-hydroxypropyl)oxazolidin-5-one (11c) (52 mg, 40%) as a pale yellow gum (Found : M^{+•} + H⁺, 334.2012. C₁₉H₂₈NO4 requires 334.2018). Mass spectrum m/z 334 (M + H⁺, 83%), 184 (100). 10c (major isomer): ¹H n.m.r. δ 0.82, d, J 6.3 Hz, 3H, H-3'; 1.05, s, 9H; 1.19, t, J 7.2 Hz, 3H, CH3; 1.95, ddd, J 7.5, 6.0 Hz, 1H, H-1'a; 3.07, ddd, J 9.0, 6.9, Hz, 1H, H-1'β; 3.36, q, J 7.2 Hz, 2H, CH2; 3.63, m, 1H, H-2'; 4.33, dd, J 7.8, 6.3 Hz, 1H, H-4; 6.11, s, 1H, H-2; 7.35-7.49, m, 5H, ArH. ¹³C n.m.r. δ 15.55, CH₃; 19.77, CH3; 25.21, CH3; 36.66, C; 42.74, CH2; 54.53, CH; 63.60, CH2; 70.63, CH; 95.38, CH; 127.12, CH; 128.55, CH; 129.94, CH; 135.29, C; 172.56, CO, 174.23, CO. 10c (minor isomer): ¹H n.m.r. δ 0.69, d, J 6.3 Hz, 3H, H-3'; 1.04, s, 9H; 1.11, t, J 6.9 Hz, 3H, CH3; 1.78, ddd, J 13.5, 8.7, 3.6 Hz, 1H, H-1'a; 2.25, ddd, J 13.8, 10.5, 5.4 Hz, 1H, H-1'β; 3.19-3.45, m, 1H, H-2'; 3.41, q, J 6.9 Hz, 2H, CH2; 4.00, dd, J 10.2, 3.0 Hz, 1H, H-4; 6.08, s, 1H, H-2; 7.35-7.49, m, 5H, ArH. ¹³C n.m.r. δ 15.39, CH₃; 18.14, CH₃; 25.17, CH₃; 36.94, C; 41.69, CH₂; 55.16, CH; 63.61, CH2; 71.15, CH; 95.39, CH; 126.42, CH; 128.90, CH; 130.47, CH; 135.76, C; 172.04, CO; 173.71, CO. 11c (major isomer, in part): ¹H n.m.r. δ6.16, s, 1H, H-2. 11c (minor isomer, in part): ¹H n.m.r. δ 6.17, s, 1H, H-2.

Photolysis of (2S)-3-Benzoyl-2-tert-butyl-4-methyleneoxazolidin-5-one (1) in Dimethoxymethane. The photolysis of 1 (100 mg, 0.39 mmol) in dimethoxymethane (150 ml) was carried out as described above. Column chromatography on silica gel, eluting with 10% ether in light petroleum, afforded an inseparable mixture of 2 diastereomers (2S,4R)-3-benzoyl-2-tert-butyl-4-(2',2'-dimethoxyethyl)oxazolidin-5-one (10a) and (2S, 4S)-3-benzoyl-2-tert-butyl-4-(2',2'-dimethoxyethyl)oxazolidin-5-one (10a) and (2S, 4S)-3-benzoyl-2-tert-butyl-4-(2',2'-dimethoxyethyl)oxazolidin-5-one (11a) (21 mg, 16%) as a pale yellow oil (Found : $M^{+\bullet} + H^+$, 336.1805. C18H26NO5 requires 336.1811). Mass spectrum m/z 336 ($M + H^+$, 30%). 10a (major isomer): ¹H n.m.r. δ 1.03, s, 9H; 2.09-2.18, m, 2H, H-2' α ; 2.94, s, 3H, OCH3; 3.23, s, 3H, OCH3; 4.21, dd, J 9.0, 5.4 Hz, 1H, H-4; 4.51, dd, J 7.5, 5.1 Hz, 1H, H-1'; 6.11, s, 1H, H-2; 7.39-7.48, m, 5H, ArH. ¹³C n.m.r. δ 25.16, CH3; 36.98, C; 37.62, CH2; 52.77, CH3; 53.40, CH3; 54.42, CH; 95.59, CH; 100.71, CH; 126.65, CH; 128.75, CH; 130.51, CH; 134.45, C; 172.04, CO; 173.75, CO. 11a (minor isomer, in part): ¹H n.m.r. δ 1.04, s, 9H; 6.07, s, 1H, H-2.

Photolysis of (2S)-3-Benzoyl-2-tert-butyl-4-methyleneoxazolidin-5-one (1) in tert-Butyl Methyl Ether. The photolysis of 1 (100 mg, 0.39 mmol) in tert-butyl methyl ether (150 ml) was carried out as described above. Column chromatography on silica gel, eluting with 10% ether in light petroleum, afforded an inseparable mixture of the diastereomers (2S,4R)-3-benzoyl-2-tert-butyl-4-(2'-tert-butyloxyethyl)oxazolidin-5-one (10b) and (2S,4S)-3-benzoyl-2-tert-butyl-4-(2'-tert-butyloxyethyl)oxazolidin-5-one (11b) (49 mg, 36%) as a colourless oil (Found : $M^{+*} + H^+$, 348.2185. C20H30NO4 requires 348.2175). Mass spectrum m/z 348 (M + H⁺, 65%). 10b (major isomer): ¹H n.m.r. δ 0.99, s, 9H; 1.05, s, 9H; 2.01-2.20, m, 2H, CH₂; 3.29-3.38, 2H, CH₂; 4.14, dd, J 9.9, 3.3 Hz, 1H, H-4; 6.12, s, 1H, H-2; 7.39-7.48, m, 5H, ArH. ¹³C n.m.r. δ 25.20, CH3; 27.27, CH3; 36.30, CH2; 36.76, C; 54.46, CH; 57.33, CH2; 72.76, C; 95.49, CH; 126.74, CH; 128.71, CH; 130.55, CH; 135.41, C; 172.32, CO; 174.09, CO. **11b** (minor isomer): ¹H n.m.r. δ 1.00, s, 9H; 1.12, s, 9H; 2.01-2.20, m, 2H, CH2; 3.17-3.21, 2H, CH2; 4.41, dd, J 9.9, 3.3 Hz, 1H, H-4; 6.17, s, 1H, H-2; 7.39-7.48, m, 5H, ArH. ¹³C n.m.r. δ 24.82, CH3; 27.10, CH3; 36.52, CH2; 37.04, C; 55.39, CH; 56.54, CH2; 73.12, C; 94.64, CH; 127.52, CH; 128.90, CH; 131.75, CH; 136.24, C; 170.59, CO; 173.14, CO.

Hydrolysis of 10b and 11b. A 95:5 mixture of 10b and 11b (30 mg, 0.086 mmol) in 6M hydrochloric acid (5 ml) was heated at 80 °C for 12 h. The reaction mixture was cooled and washed with dichloromethane (4 x 10 ml). The aqueous fraction was evaporated to dryness. Recrystallization of the crude residue from ethanol/ether afforded the hydrochloride salt of (**R**)-α-amino-γ-butyrolactone ((*R*)-12) as a white solid (11 mg, 90%), m.p. >200°C. (Found : M^{+•} + H⁺, 102.0542. C4H8NO2 requires 102.0555). [α]²⁰D -20° (c, 0.10 in 6M HCl) (lit.¹² [α]²⁰D -32° (c, 0.2 in 2M HCl)). ¹H n.m.r. (D2O) δ 2.32-2.43, m, 1H, H-βa; 2.67-2.76, m, 1H, H-βb; 4.43-4.55, m, 3H, H-α & H-γa/b. ¹³C n.m.r. (D2O) δ 28.00, CH2; 49.78, CH; 68.58, CH2; 175.69, CO. Mass spectrum m/z 102 (M - HCl + H⁺, 100%).

Photolysis of (2S)-3-Benzoyl-2-tert-butyl-4-methyleneoxazolidin-5-one (1) in tert-Butyl Ethyl Ether. The photolysis of 1 (100 mg, 0.39 mmol) in a mixture of tert-butyl ethyl ether (2.5 g, 24.5 mmol) and dry acetonitrile (150 ml) was carried out as described above. Column chromatography on silica gel, eluting with 10% ether in light petroleum, afforded 3 partially separable stereoisomers (2S,4R)-3-benzoyl-2-tert-butyl-4-(2'-tert-butyloxypropyl)oxazolidin-5-one (10d) and (2S, 4S)-3-benzoyl-2-tert-butyl-4-(2'-tertbutyloxypropyl)oxazolidin-5-one (11d) (67 mg) in a combined yield of 48%. Samples of 10d (major) and 10d (minor) were obtained diastereomerically pure (Found: M⁺⁺ + H⁺, 362.2334. C₂₁H₃₂NO4 requires 362.2331). Mass spectrum m/z 362 (M + H⁺, 100%), 307 (40), 220 (45). **10d** (major isomer): colourless oil, $[\alpha]^{23}$ D + 3° (c, 0.12 in CHCl3). ¹H n.m.r. δ 0.89, s, 9H; 1.05, s, 9H; 1.07, d, J 6.3 Hz, 3H, CH3; 1.90-1.96, m, 2H, H-1'\alpha & H-1'\beta; 3.92, app. sextet, J 6.0 Hz, 1H, H-2'; 4.23, dd, J 7.5, 6.9 Hz, 1H, H-4; 6.13, s, 1H, H-2; 7.39-7.49, m, 5H, ArH. ¹³C n.m.r. δ 24.22, CH3; 25.31, CH3; 28.54, CH3; 36.63, C; 44.00, CH2; 54.49, CH; 63.29, CH; 73.06, C; 95.29, CH; 126.97, CH; 128.78, CH; 130.73, CH; 135.28, C; 172.72, CO; 174.61, CO. **10d** (minor isomer): colourless oil, $[\alpha]^{23}D + 8^{\circ}$ (c, 0.12 in CHCl₃). ¹H n.m.r. δ 0.56, d, J 6.0 Hz, 3H, CH₃; 1.05, s, 9H; 1.11, s, 9H; 1.82, ddd, J 13.2, 10.2, 2.7 Hz, 1H, H-1'α; 2.16, ddd, J 12.9, 12.9, 3.3 Hz, 1H, H-1'β; 3.79-3.82, m, 1H, H-2'; 3.85, dd, J 10.2, 3.0 Hz, 1H, H-4; 6.10, s, 1H, H-2; 7.37-7.50, m, 5H, ArH. ¹³C n.m.r. δ 20.60, CH3; 25.14, CH3; 28.10, CH3; 36.90, C; 43.70, CH2; 55.45, CH; 63.31, CH; 74.22, C; 95.43, CH; 126.41, CH; 128.97, CH; 130.57, CH; 135.63, C; 172.13, CO; 173.99, CO. 11d (in part): ¹H n.m.r. δ 1.04, s, 9H; 1.10, s, 9H; 6.08, s, 1H, H-2.

Hydrolysis of the major diastereomer of 10d. A mixture of 10d (major) (15 mg, 0.042 mmol) and 6M hydrochloric acid (5 ml) was heated at 90 °C for 12 h. The reaction mixture was cooled and washed with dichloromethane (4 x 10 ml). The aqueous fraction was evaporated to dryness to afford a pale yellow gum (5.5 mg, 88%). ¹H n.m.r. analysis of the crude residue showed this to be a mixture of 2 diastereomers hydrochloride salts of (**R**,**R**)-α-amino-γ-methyl-γ-butyrolactone ((R, R)-13) and (**R**_α,**S**_γ)-α-amino-γ-methyl-γ-butyrolactone ((R, R)-14) in a ratio of approximately 71 : 29 (Found : M^{+•} - HCl + H⁺, 116.0703. C5H₁₀NO₂ requires 116.0712). Mass spectrum m/z 116 (M - HCl + H⁺, 100%). 13 (major isomer): ¹H n.m.r. (D₂O) δ 1.36, dd, J 6.6 Hz, 1.5 Hz, 3H, CH₃; 2.46-2.52, m, 2H, H-βa & H-βb; 4.50, app. t, J 9.9 Hz, 1H, H-α; 4.43-4.55, m, 1H, H-γ. ¹³C n.m.r. (D₂O) δ 24.01, CH₃; 33.15, CH₂; 49.24, CH; 66.93, CH; 173.12,

CO. 14 (minor isomer): ¹H n.m.r. (D₂O) δ 1.31, dd, J 6.3 Hz, 1.5 Hz, 3H, CH₃; 1.87, ddd, J 12.8, 10.1, 1.5 Hz, 1H, H- β a; 2.75, ddd, J 13.5, 8.7, 5.1 Hz, 1H, H- β b; 4.35, dd, J 11.7, 8.7 Hz, 1H, H- α ; 4.65-4.73, m, 1H, H- γ . ¹³C n.m.r. (D₂O) δ 20.76, CH₃; 35.72, CH₂; 51.34, CH; 68.40, CH; 175.15, CO.

Hydrolysis of the minor diastereomer of 10d. A sample of 10d (minor) (12 mg, 0.033 mmol) was hydrolyzed according to the above procedure to afford a pale yellow gum (4 mg, 80%). ¹H n.m.r. analysis of the crude residue showed this to be a mixture of 14 : 13 in a ratio of approximately 91: 9, respectively. The spectral data of 13 and 14 were identical to that described above.

Acknowledgment: We thank the Australian Research Council for financial support.

References

- 1. Nair, S. A.; Lee, B.; Hangauer, Synthesis 1995, 810.
- 2. Koch, T.; Buchardt, O. Synthesis 1993, 1065.
- 3. Barclay, F.; Chrystal, E.; Gani, D. J. Chem. Soc., Perkin Trans 1 1996, 683.
- Garner, P. P.; Cox, P. B.; Klippenstein, S. J. J. Am. Chem. Soc. 1995, 117, 4183. Garner, P.; Leslie, R.; Anderson, J. T. J. Org. Chem. 1996, 61, 6754.
- 5. Zimmermann, J., Seebach, D., Helv. Chim. Acta. 1987 70, 1104.
- 6. Pyne, S. G., Dikic, B., Gordon, P., Skelton, B. W., White, A. H. Aust. J. Chem. 1993, 46, 73.
- 7. Gomez, A. M.; Mantecon, Valverde, S.; Lopez, J. C. J. Org. Chem. 1997, 62, 6612.
- Xiang, J.; Fuchs, P. L. J. Am. Chem. Soc. 1996, 118, 11986. Clark, A. J.; Rooke, S.; Sparey, T. J.; Taylor, P. C.Tetrahedron Lett. 1996,37, 909. Ogura, K.; Yanagisawa, A.; Fujino, T.; Takahashi, K.Tetrahedron Lett. 1988, 29, 5387.
- 9. Axon, J. R.; Beckwith, A. L. J. J. Chem. Soc., Chem. Commun. 1995, 549.
- 10. Dofour, M.; Gramain, J.-C.; Husson, H.-P.; Sinibaldi, M.-E.; Trion, Y. Tetrahedron Lett. 1989, 30, 3429.
- 11. Ibrahim-Ouali, M.; Sinibaldi, M.-E.; Trion, Y.; Gramain, J.-C. Tetrahedron Lett. 1996, 37, 37.
- 12. Greenstein, J. P.; Winitz, M.Chemistry of the Amino Acids, Wiley: New York, 1961; pp 2612-2616, 2638.
- 13. Jacob, M.; Roumestant, M. L.; Viallefont, P.; Martinez, J. Synlett 1997, 691.
- 14. Gull, R.; Schöllkopf, U. Synthesis 1985, 1052.
- 15. Matzinger, P.; Catalfomo, Ph.; Eugster, C. H. Helv. Chim. Acta, 1972, 55, 1478.
- 16. Harding, K. E.; Marman, T. H.; Nam, D.-H. Tetrahedron 1988, 44, 5605.
- 17. Ariza, J.; Font, J.; Ortuno, R. A. Tetrahedron 1990, 46, 1931.
- 18. Schmeck, C.; Hegedus, L. S. J. Am. Chem. Soc. 1994, 116, 9927.
- 19. For related studies see: Jackson, R. F. W.; Rettie, A. B.; Wood, A.; Wythes, M.J. J. Chem. Soc. Perkin Trans. 1, 1994, 1719.
- For a recent application of 2 for the synthesis of amino acids in high enantiomeric purities see: Pyne, S. G.; Schafer, K.; Skelton, B. W.; White, A. H. J. Chem. Soc., Chem. Commun. 1997, 2267.