<u>LETTERS</u>

Bi(OTf)₃-Mediated Cycloisomerization of γ -Alkynyl Arylketones: Application to the Synthesis of Substituted Furans

Meng-Yang Chang,* Yu-Chieh Cheng, and Yi-Ju Lu

Department of Medicinal and Applied Chemistry, Kaohsiung Medical University, Kaohsiung 807, Taiwan

Supporting Information

ABSTRACT: A novel Bi(OTf)₃-mediated cycloisomerization of γ alkynyl arylketones 4, 7, or 10 with molecular sieve (MS) in MeNO₂ affords 3-substituted furans 3, 8, or 11 at rt for 3 h in moderate to good yields. The method provides mild, less-toxic, atom-economic and efficient conditions. The mechanism has been studied and proposed. Moreover, this route can be enlarged to gram scale.



F unctionalized furan is a versatile synthetic building block in numerous biologically active molecules, pharmaceuticals, and natural products.¹ Development of new routes to substituted furans (1) from readily available starting materials still represents a continuing need in the organic synthetic field. Among these protocols, transition metals (Pd^{2+} , Ag^+/Au^{3+} , In^{3+} , or Zn^{2+}) promoting cycloisomerization of diversified γ -alkynones (or in situ generated from coupling of alkynes and 1,3-dicarbonyl synthons) is a major pathway (see Scheme 1).²⁻⁴ Routes to base

Scheme 1. Metal-Mediated Syntheses of Substituted Furans (1)



(DBU or NaH) mediated cyclization of γ -alkynones have been documented.⁵ However, the preparation of furan derivatives often presents drawbacks (e.g., multistep operations and harsh conditions), and this has encouraged organic researchers to explore more efficient synthetic protocols. To the best of our knowledge, for the synthesis of furans, no literature on Bi(III)-mediated annulation of γ -alkynones has been reported. Bi(III) compounds with nontoxic and environmentally friendly properties has been documented for a wide variety of organic reactions.^{6,7} Bi(III)-promoted reactions of γ -alkenyl,^{7a,b} γ -allenyl,^{7c} or γ -alkynyl^{7d} carbonyl synthons provide diversified skeletons.

In continuation of our investigation of the application of α sulfonyl ketones **2**,⁸ an easy and atom-economic Bi(OTf)₃mediated intramolecular 5-*exo-dig* cycloisomerization of α sulfonyl- γ -alkynyl arylketones **4** (derived from alkylation of **2** with propargylic bromides) is employed to construct the structure of 3-sufonyl furans 3, as shown in Scheme 2. A number of articles have highlighted fascinating developments based on





the skeleton of α -sulfonyl ketones (β -ketosulfones)⁹ and γ alkynones are important intermediates in organic transformations, which can be easily constructed to the other functionalized frameworks such as dihydropyrroles or dihydrofurans.¹⁰ Sulfonyl furans exhibit some potent biological activities, such as an endothelial lipase inhibitor or EP1 receptor antagonist.¹¹

After further comparison of literature on the metal-catalyzed benzannulation of γ -alkynones, we chose model substrate 4a (R = Tol, Ar = Ph, Y = H) to initiate our studies with the screening of metal triflate-mediated reaction conditions. Alkylation of 2a with propargyl bromide afforded 4a in a 90% yield in boiling acetone for 8 h. Initially, the use of various metal triflates as catalysts was investigated for the formation of the 3-sulfonyl furan 3a. Catalytic amounts of AgOTf, Mg(OTf)₂, Cu(OTf)₂, Ni(OTf)₂, Sn(OTf)₂, Zn(OTf)₂, Hg(OTf)₂, Yb(OTf)₃, Sm(OTf)₃, Ga-(OTf)₃, In(OTf)₃, La(OTf)₃, Sc(OTf)₃, Fe(OTf)₃, and Bi-(OTf)₃ provided different results in MeNO₂ at 25 °C for 3 h. Among the catalysts screened, AgOTf, $Hg(OTf)_2$, or $In(OTf)_3$ provided 5a as the sole product in 81%, 86%, or 88% yields, respectively (see Table 1, entries 1-3).^{4a-d} Bi(OTf)₃ produced the desired product 3a in an 80% yield along with a 10% yield of 5a (entry 4). Under these conditions, the recovery of the starting material 4a was only observed for the other 11 metal triflates. Furthermore, the use of other Bi(III) salts was examined for synthesizing 3a and treatment of 4a with BiCl₃, BiBr₃, Bi(OAc)₃, or $Bi_2(SO_4)_3$ produced 5a in good yields (70%, 75%, 68%, and 66%) at 25 °C for 3 h with no isolation of 3a (entries 5–8). In





^{*a*}Reaction was run on a 1.0 mmol scale with 4a, 3 h. ^{*b*}Isolated yield. ^{*c*}72% (for entry 13) or 60% (for entry 15) of 4a was recovered. ^{*d*}60 h (for entry 13); 45 h (for entry 15). ^{*e*}4 Å MS (100 mg) was added. ^{*f*}H₂O (0.1 mL) was added.

comparison with this Bi(III) complex, Bi(OTf)₃ is an optimal catalyst for the generation of **3a**. Controlling the $Bi(OTf)_3$ as the catalyst, the factors of equivalence, concentration, and temperature were studied next. When using 5 mol % of $Bi(OTf)_3$, the isolated yield was similar to 2 mol % (entry 9). No obvious changes occurred when decreasing the reaction concentration (5 \rightarrow 10) or elevating the temperature (25 \rightarrow 100) as shown in entries 10-11, respectively. After changing the solvents (from MeNO₂ to MeCN, dichloroethane, benzene, or 1,4-dioxane), different results were observed (entries 12-15). Entry 16 showed that removal of water decreased the yield of 3a in the presence of molecular sieves (MS). On the contrary, the involvement of water increased the ratio of 4a and 3a (entry 17). The results show that the introduction of water caused competition between intermolecular hydration (for the methyl ketone skeleton) and intramolecular 5-exo-dig addition (for furan skeleton) during the Bi(OTf)3-mediated process. Overall, we believed that the Bi(OTf)₃/MeNO₂/MS system should be an optimal combination for generating 3a.

On the basis of the results, a possible reaction mechanism is shown in Scheme 3. How are 3a and 5a produced? The mechanism should be initiated to form A by complexation of an alkynyl motif of 4a with Bi(OTf)₃, and participation of a carbonyl group could lead to the oxonium cation B via intramolecular 5*exo-dig* annulation. Deprotonation of B should give vinyl bismuth intermediate C. Then, protonation by in situ generated HOTf achieves to alternative oxonium cation D, which following debismuthation is able to provide E and recover Bi(OTf)₃. Isomerization from E forms 3a. However, F should be proposed via the involvement of A with water. After deprotonation of F, tautomerization of G, and then HOTf-promoted debismuthation of β -ketobismuth intermediate H, the removal of Bi(OTf)₃ affords I. Subsequently, the tautomerization of H generates 5a. Scheme 3. Possible Mechanism



With optimized conditions in hand (Table 1, entry 16), we further explored the substrate scope of the reaction, and the results are shown in Table 2. For the Ar and R substituents of 3 (Ar = Ph, 4-FC₆H₄, 4-MeOC₆H₄, 4-MeC₆H₄, 4-CF₃C₆H₄, 4-PhC₆H₄, 2-naphthalene, and 2-furan; R = 4-MeC₆H₄, Ph, Me, 3-MeC₆H₄, 4-fBuC₆H₄, 4-MeOC₆H₄, and 4-FC₆H₄), the aromatic ring, with both electron-withdrawing and electron-donating

Table 2. Synthesis of 3^{a-b}

R S O Ar	$Br_{Actor of Actor $	R O Tol	S O Ph O Ph
2	reflux, 8 h 4	3	5b
		yield (%) ^c	
entry	2, Ar, R; Y	4	3
1	2a , Ph, 4-MeC ₆ H ₄ ; H	4a , 90	3a , 86
2	2b , 4-FC ₆ H ₄ , 4-MeC ₆ H ₄ ; H	4b , 92	3b , 76
3	2c , 4-MeOC ₆ H ₄ , 4-MeC ₆ H ₄ ; H	4c , 88	3 c, 75
4	2d , 4-MeC ₆ H ₄ , 4-MeC ₆ H ₄ ; H	4d , 90	3d , 78
5	2e , 4-CF ₃ C ₆ H ₄ , 4-MeC ₆ H ₄ ; H	4e , 85	3e , 80
6	2f , 4-PhC ₆ H ₄ , 4-MeC ₆ H ₄ ; H	4f , 90	3f , 82
7	2g , 2-naphthalene, 4-MeC ₆ H ₄ ; H	4g , 86	3g , 78
8	2h , 4-FC ₆ H ₄ , Ph; H	4h , 89	3h , 73
9	2i , 4-MeOC ₆ H ₄ , Ph; H	4i , 90	3i , 75
10	2 <i>j</i> , 4-PhC ₆ H ₄ , Ph; H	4 j, 90	3 j, 78
11	2k , 4-MeC ₆ H ₄ , Me; H	4k , 86	3k , 76
12	2l , 4-PhC ₆ H ₄ , Me; H	4l , 89	31 , 80
13	2m , Ph, 3-MeC ₆ H ₄ ; H	4m , 86	3m , 82
14	2n , Ph, 4- <i>t</i> BuC ₆ H ₄ ; H	4n , 85	3n , 78
15	20 , Ph, 4-MeOC ₆ H ₄ ; H	40 , 87	30 , 75
16	2p , Ph, 4-FC ₆ H ₄ ; H	4p , 88	3p , 77
17	2q , 2-thiophene, 4-MeC ₆ H ₄ ; H	4q , 85	3q , 78
18	2r , Ph, 4-MeC ₆ H ₄ ; Me	4r , 84	3r , 62
19	2s , Ph, 4-MeC ₆ H ₄ ; Et	4s , 83	3s , 60
20	2t , Ph, 4-MeC ₆ H ₄ ; Ph	4t , 80	$3t^d$

^{*a*}α-Alkylation was run on a 1.0 mmol scale with **2**, K₂CO₃ (2.9 mmol), propargylic bromides (1.05 mmol), acetone (10 mL), reflux, 8 h. ^{*b*}Cycloisomerization was run on a 1.0 mmol scale with **4**, Bi(OTf)₃ (13 mg, 0.02 mmol), MeNO₂ (5 mL), rt, 3 h. ^{*c*}Isolated yield.

substituents, was well tolerated, providing the desired products 4 and 3 in moderate to good yields. First, by controlling the use of 1.05 equiv of propargylic bromides (Y = H, Me, Et, and Ph), 4a-twith an α -propargyl group was yielded in good yields (entries 1– 20). The cycloisomerization reaction of 4 with the combination of Bi(OTf)₃/MeNO₂/MS was next examined. All entries showed that 3a-t were isolated in a range of 60%-86% yields when Ar and R were the alkyl or aryl groups. For 3a-q, the involvement of different substituents did not affect the reaction procedure and the isolated yield was maintained (entries 1-17). When Y was changed to Me and Et groups, the isolated yields of 3r-s were decreased to 60% and 62%, respectively (entries 18-19). Furthermore, Y was adjusted to the Ph group, 5b was isolated in a 78% yield, and no desired 3t was formed (entry 20). The structures of sulfonyl 2-arylfurans 3e-f, 3h, and 3k were determined by single-crystal X-ray crystallography.¹⁷

Moreover, when 1.56 g of 4a (5.0 mmol) was treated with the combination, 1.17 g of 3a was isolated in a 75% yield. This route can be enlarged to gram scale. To examine the limitation of this route (see Scheme 4), the sulfonyl group was first changed to the





aryl, ethyl ester, and benzoyl substituents. The intramolecular cyclization of 7a-d (prepared from α -alkylation of 6a-d with NaH in THF) with the optimal combination of $Bi(OTf)_3/$ $MeNO_2/MS$ afforded a skeleton of 2,3-diarylfurans 8a-d (Ar = Ph, 4-MeOC₆H₄, 4-FC₆H₄, 2-naphthalene) with similar distribution of yields (77-82%) under the above reaction conditions (see eq 1, Scheme 4). The structure of 8b was determined by single-crystal X-ray crystallography.¹² Furthermore, α -alkylation of 9a-b provided 10a-c in good yields (88%-92%) for generating the skeleton of 2-arylfuran with a 3electronwithdrawing group (see eq 2, Scheme 4). With 10a-c in hand, Bi(OTf)₃-mediated cycloisomerization of γ -alkynones **10a**–**c** with the α -ester and benzoyl groups in the presence of molecular sieve in MeNO₂ at rt for 3 h afforded furans 11a-c in 60-81%.¹³ In particular, **12** was isolated in a 25% yield via an intramolecular aldolization of the resulting benzoyl δ -diketone of 10c. To investigate the unsymmetrical cycloisomerization of β diketones (see eq 3, Scheme 4), the $Bi(OTf)_3$ -mediated reaction of 13 (generated from α -alkylation of 9c in 68% yield) was examined next. Under the above conditions, a major isomer 14a was isolated (65%) along with 10% of 14b because the stable β - phenyl chalcone intermediate of 13 bearing a full conjugation was generated easily.¹⁴

In summary, Bi(OTf)₃-mediated cycloisomerization of α substituted γ -alkynyl arylketones is developed for synthesizing a series of 2-arylfurans in good yields. The structures of the key products were confirmed by X-ray crystallography. This atomeconomic route can be enlarged to gram scale. Further investigation regarding synthetic applications of α -sulfonyl ketones will be conducted and published in due course.

ASSOCIATED CONTENT

Supporting Information

Experimental data and scanned photocopies of ¹H and ¹³C NMR spectral data. This material is available free of charge via the Internet at http://pubs.acs.org.

AUTHOR INFORMATION

Corresponding Author

*E-mail: mychang@kmu.edu.tw.

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENTS

We thank the Ministry of Science and Technology of the Republic of China for financial support (NSC 102-2113-M-037-005-MY2).

REFERENCES

(1) For reviews on syntheses of furans, see (a) Hou, X. L.; Cheung, H. Y.; Hon, T. Y.; Kwan, P. L.; Lo, T. H.; Tong, S. Y. T.; Wong, H. N. C. *Tetrahedron* 1998, *54*, 1955. (b) Dean, F. M. *Adv. Heterocycl. Chem.* 1982, *30*, 167. (c) Donnelly, D. M. X.; Meegan, M. *J. Compd. Heterocycl. Chem.* 1984, *4*, 657. (d) Hou, X. L.; Yang, Z.; Wong, H. N. C. *Progress in Heterocyclic Chemistry*; Gribble, G. W., Gilchrist, T. L., Eds.; Pergamon Press: Oxford, U.K., 2002; Vol. *14*, p 139.

(2) For Pd(II)-mediated cyclizations of 4-alkynones, see (a) Arcadi, A.; Cacchi, S.; Fabrizi, G.; Marinelli, F.; Parisi, L. M. Tetrahedron 2003, 59, 4661. (b) Picquet, M.; Bruneau, C.; Dixneuf, P. H. Tetrahedron 1999, 55, 3937. (c) Cacchi, S.; Fabrizi, G.; Moro, L. J. Org. Chem. 1997, 62, 5327.
(d) Li, Y.; Yu, Z. J. Org. Chem. 2009, 74, 8904. (e) Arcadi, A.; Rossi, E. Tetrahedron Lett. 1996, 37, 6811. (f) Saito, A.; Enomoto, Y.; Hanzawa, Y. Tetrahedron Lett. 2011, 52, 4299.

(3) For Ag(I)/Au(III)-mediated cyclizations of 4-alkynones, see
(a) Arcadi, A.; Alfonsi, M.; Chiarini, M.; Marinelli, F. J. Organomet. Chem. 2009, 694, 576. (b) Belting, V.; Krause, N. Org. Biomol. Chem.
2009, 7, 1221. (c) Wang, T.; Zhang, J. Dalton Trans 2010, 39, 4270.

(4) Other metal-mediated cyclizations of 4-alkynones: for Ag, see (a) Das, R.; Chakraboty, D. *Appl. Organomet. Chem.* **2012**, *26*, 722. For Hg(II), see (b) Imagawa, H.; Kurisaki, T.; Nishizawa, M. Org. Lett. **2004**, *6*, 3679. For In(III), see (c) Feng, X.; Tan, Z.; Chen, D.; Shen, Y.; Guo, C.-C.; Xiang, J.; Zhu, C. *Tetrahedron Lett.* **2008**, *49*, 4110. (d) Tsuji, H.; Yamagata, K.-i.; Ueda, Y.; Nakamura, E. *Synlett* **2011**, 1015. For Zn(II), see (e) Liu, C.-R.; Li, M.-B.; Yang, C.-F.; Tian, S. *Chem.—Eur. J.* **2009**, *15*, 793.

(5) For base-mediated cyclizations of 4-alkynones, see (a) Vieser, R.; Eberbach, W. *Tetrahedron Lett.* **1995**, *36*, 4405. (b) Reddy, C. R.; Reddy, M. D. J. Org. Chem. **2014**, *79*, 106.

(6) For reviews on bismuth-promoted reactions, see (a) Briand, G. G.; Burford, N. Chem. Rev. 1999, 99, 2601. (b) Roux, C. L.; Dubac, J. Synlett
2002, 181. (c) Leonard, N. M.; Wieland, L. C.; Mohan, R. S. Tetrahedron
2002, 58, 8373. (d) Gaspard-Iloughmane, H.; Le Roux, C. Eur. J. Org. Chem. 2004, 2517. (e) Gaspard-Iloughmane, H.; Le Roux, C. Trends
Org. Chem. 2006, 11, 65. (f) Hua, R. Curr. Org. Synth. 2008, 5, 1.
(g) Salvador, J. A. R.; Pinto, R. M. A.; Silvestre, S. M. Curr. Org. Synth.
2009, 6, 426. (h) Salvador, J. A. R.; Pinto, R. M. A.; Silvestre, S. M. Mini-

Organic Letters

Rev. Org. Chem. 2009, 6, 241. (i) Bothwell, J. M.; Krabbez, S. W.; Mohan, R. S. Chem. Soc. Rev. 2011, 40, 4649. (j) Ollevier, T. Org. Biomol. Chem. 2013, 11, 2740.

(7) For the representative Bi(OTf)₃-mediated transformations, see
(a) Kelly, B. D.; Allen, J. M.; Tundel, R. E.; Lambert, T. H. Org. Lett. **2009**, 11, 1381. (b) Godeau, J.; Olivero, S.; Antoniotti, S.; Dunach, E. Org. Lett. **2011**, 13, 3320. (c) Lemiere, G.; Cacciuttolo, B.; Belhassen, E.; Dunach, E. Org. Lett. **2012**, 14, 2750. (d) Komeyama, K.; Takahashi, K.; Takaki, K. Org. Lett. **2008**, 10, 5119. (e) Murai, M.; Hosokawa, N.; Roy, D.; Takai, K. Org. Lett. **2014**, 16, 4134. (f) Murai, M.; Origuchi, K.; Takai, K. Org. Lett. **2014**, 16, 3828. (g) Jaratjaroonphong, J.; Tuengpanya, S.; Ruengsangtongkul, S. J. Org. Chem. **2015**, 80, 559. (h) Cacciuttolo, B.; Poulain-Martini, S.; Fontaine-Vive, F.; Abdo, M. A. H.; El-Kashef, H.; Dunach, E. Eur. J. Org. Chem. **2014**, 7458. (i) Cacciuttolo, B.; Poulain-Martini, S.; Dunach, E. Eur. J. Org. Chem. **2014**, 7458. (b) Cacciuttolo, B.; Poulain-Martini, S.; Dunach, E. Eur. J. Org. Chem. **2014**, 7458. (i) Cacciuttolo, B.; Poulain-Martini, S.; Bunach, E. Eur. J. Org. Chem. **2014**, 7458. (i) Cacciuttolo, B.; Poulain-Martini, S.; Dunach, E. Eur. J. Org. Chem. **2014**, 7458. (i) Cacciuttolo, B.; Poulain-Martini, S.; Dunach, E. Eur. J. Org. Chem. **2014**, 7458. (i) Cacciuttolo, B.; Poulain-Martini, S.; Dunach, E. Eur. J. Org. Chem. **2014**, 7458. (i) Cacciuttolo, B.; Poulain-Martini, S.; Dunach, E. Eur. J. Org. Chem. **2014**, 7458. (i) Cacciuttolo, B.; Poulain-Martini, S.; Dunach, E. Eur. J. Org. Chem. **2014**, 7458. (i) Cacciuttolo, B.; Poulain-Martini, S.; Dunach, E. Eur. J. Org. Chem. **2014**, 7458. (i) Cacciuttolo, B.; Poulain-Martini, S.; Dunach, E. Eur. J. Org. Chem. **2014**, 7458. (i) Cacciuttolo, B.; Poulain-Martini, S.; Dunach, E. Eur. J. Org. Chem. **2015**, 56, 1045. (k) Komeyama, K.; Yamada, T.; Igawa, R.; Takaki, K. Chem. Commun. **2012**, 48, 6372.

(8) (a) Chang, M.-Y.; Chen, Y.-C.; Chan, C.-K. *Tetrahedron* 2014, *70*, 8908. (b) Chang, M.-Y.; Chen, Y.-C.; Chan, C.-K. *Synlett* 2014, *25*, 1739.
(c) Chang, M.-Y.; Cheng, Y.-C.; Lu, Y.-J. Org. *Lett.* 2014, *16*, 6252.

(9) For recent synthesis of α -sulfonyl ketones, see (a) Pospisil, J.; Sato, H. J. Org. Chem. 2011, 76, 2269. (b) Sreedhar, B.; Rawat, V. S. Svnlett 2012, 23, 413. (c) Kumar, A.; Muthyala, M. K. Tetrahedron Lett. 2011, 52, 5368. (d) Suryakiran, N.; Reddy, T. S.; Ashalatha, K.; Lakshman, M.; Venkateswarlu, Y. Tetrahedron Lett. 2006, 47, 3853. (e) Lu, Q.; Zhang, J.; Zhao, G.; Qi, Y.; Wang, H.; Lei, W. J. Am. Chem. Soc. 2013, 135, 11481. (f) Tsui, G. C.; Glenadel, Q.; Lau, C.; Lautens, M. Org. Lett. 2011, 13, 208. (g) Curti, C.; Laget, M.; Carle, A. O.; Gellis, A.; Vanelle, P. Eur. J. Med. Chem. 2007, 42, 880. (h) Bin, J. K.; Lee, J. S.; Kim, K. Org. Lett. 2004, 6, 4297. (i) Zhou, G.; Ting, P. T.; Aslanian, R. G. Tetrahedron Lett. 2010, 51, 939. (j) Tang, X.; Huang, L.; Xu, Y.; Yang, J.; Wu, W.; Jiang, H. Angew. Chem., Int. Ed. 2014, 53, 4205. (k) Chawla, R.; Singh, A. K.; Yadav, L. D. S. Eur. J. Org. Chem. 2014, 10, 2032 and cited references therein. (1) Singh, A. K.; Chawla, R.; Yadav, L. D. S. Tetrahedron Lett. 2014, 55, 4742. (m) Shi, X.; Ren, X.; Ren, Z.; Li, J.; Wang, Y.; Yang, S.; Gu, J.; Gao, Q.; Huang, G. Eur. J. Org. Chem. 2014, 10, 5083. (n) Xuan, J.; Feng, Z.-J.; Chen, J.-R.; Lu, L.-Q.; Xiao, W.-J. Chem.-Eur. J. 2014, 20, 3045.

(10) (a) Sun, B.; Ma, Q.; Wang, Y.; Zhao, Y.; Liao, P.; Bi, X. *Eur. J. Org. Chem.* **2014**, 7552. (b) Ma, Q.; Wang, Y.; Zhao, Y.; Liao, P.; Sun, B.; Bi, X. *Eur. J. Org. Chem.* **2014**, 4999.

(11) For the potential biological activities of sulfonyl furans, see
(a) Goodman, K. B.; Bury, M. J.; Cheung, M.; Cichy-Knight, M. A.; Dowdell, S. E.; Dunn, A. K.; Lee, D.; Lieby, J. A.; Moore, M. L.; Scherzer, D. A.; Sha, D.; Suarez, D. P.; Murphy, D. J.; Harpel, M. R.; Manas, E. S.; McNulty, D. E.; Annan, R. S.; Matico, R. E.; Schwartz, B. K.; Trill, J. J.; Sweitzer, T. D.; Wang, D.-y.; Keller, P. M.; Krawiec, J. A.; Jaye, M. C. *Bioorg. Med. Chem. Lett.* **2009**, *19*, 27. (b) Naganawa, A.; Matsui, T.; Ima, M.; Saito, T.; Murota, M.; Aratani, Y.; Kijima, H.; Yamamoto, H.; Maruyama, T.; Ohuchida, S.; Nakai, H.; Toda, M. *Bioorg. Med. Chem.* **2006**, *14*, 7121. (c) Wendt, M. D.; Geyer, A.; McClellan, W. J.; Rockway, T. W.; Weitzberg, M.; Zhao, X.; Mantei, R.; Stewart, K.; Nienaber, V.; Klinghofer, V.; Giranda, V. L. *Bioorg. Med. Chem. Lett.* **2004**, *14*, 3069. (d) Bouhlel, A.; Curti, C.; Dumetre, A.; Laget, M.; Crozet, M. D.; Azas, N.; Vanelle, P. *Bioorg. Med. Chem.* **2010**, *18*, 7310.

(12) CCDC 1041267 (3e), 1041265 (3f), 1041266 (3h), 1041529 (3k), and 1044470 (8b) contain the supplementary crystallographic data for this paper. This data can be obtained free of charge via www. ccdc.cam.ac.uk/conts/retrieving.html (or from the CCDC, 12 Union Road, Cambridge CB2 1EZ, U.K.; fax, 44–1223–336033; e-mail, deposit@ccdc.cam.ac.uk).

(13) (a) Arcadi, A.; Cerichelli, G.; Chiarini, M.; Di Giuseppe, S.; Marinelli, F. *Tetrahedron Lett.* **2000**, *41*, 9195. (b) Chen, Y.-F.; Wang, H.-F.; Wang, Y.; Luo, Y. C.; Zhu, H.-L.; Xu, P.-F. *Adv. Synth. Catal.* **2010**, 352, 1163.

(14) For **14a**, see (b) Han, X.; Widenhoefer, R. A. *J. Org. Chem.* **2004**, 69, 1738. For **14b**, see (a) Tiecco, M.; Testaferri, L.; Tingoli, M.; Marini, F. *J. Org. Chem.* **1993**, *58*, 1349.