



Synthesis of allylamines in enantiomerically pure form

José M. Concellón,* Beatriz Baragaña and Estela Riego

Departamento de Química Orgánica e Inorgánica, Facultad de Química, Universidad de Oviedo, Julián Clavería, 8, 33071 Oviedo, Spain

Received 11 February 2000; accepted 7 April 2000

Abstract

Enantiopure allylamines are obtained without racemization by reaction of α -amino aldehydes with chloromethylithium followed by lithiation. © 2000 Elsevier Science Ltd. All rights reserved.

Keywords: amines; elimination reactions; enantiomeric purity; lithiation.


Chiral allylamines are synthetic precursors to a number of important classes of compounds, such as 4-aminoallylsilanes,¹ aminoepoxides,² iodocyclocarbamates³ and isoxazolines.⁴ One general method³ for the synthesis of allylic amines involves the Wittig olefination reaction with amino aldehydes.⁵ However, α -amino aldehydes are often susceptible to racemization⁶ and there is a possibility that racemization may also occur during the Wittig reaction.^{2a}

Previously, we have reported the synthesis of chiral *erythro* or *threo* aminoepoxides by epoxidation of *erythro* or *threo* 3-amino-1-haloalkan-2-ols. These enantiopure halohydrins can be obtained by reaction of halomethylithium (generated in situ) with α -amino aldehydes or by reduction of chiral α' -amino α -chloro ketones, respectively.⁷ Herein we wish to report a general one-pot synthesis of enantiopure allylamines by lithiation of chiral 3-amino-1-haloalkan-2-ols, which can be prepared by reaction of chloromethylithium with chiral α -amino aldehydes.

The reaction of amino aldehyde **1** with chloromethylithium (generated in situ from chloroiodomethane and methylithium) at -78°C led to the expected chlorinated lithium alcoholate **2**; this intermediate was in situ lithiated with lithium powder at -45°C affording a β -functionalized organolithium intermediate **3**, which undergoes spontaneous β -elimination yielding, after hydrolysis, the corresponding allylamine **4** (Scheme 1).

Under these reaction conditions, the allylic amines **4** were obtained with no detectable racemization. The enantiomeric purity of product **4b** was determined by chiral HPLC (Chiracel OD-H) analysis,⁸ showing an enantiomeric excess (ee) $> 99\%$. Previously, other chiral HPLC analysis was carried out, using a racemic mixture of **4b** to exclude the possibility of co-elution of both enantiomers.

* Corresponding author. Fax: 34-8-510-3446; e-mail: jmcg@sauron.quimica.uniovi.es



Scheme 1. Synthesis of enantiopure allylamines

The described transformation can also be carried out with functionalized amino aldehydes such as *O*-silylated serinal **1d** and it is noteworthy that this methodology provides directly enantiopure allylamines, from α -amino aldehydes, without isolating the intermediate **2**.

A typical procedure for the synthesis of allylamines **4** is as follows: to a -78°C solution of the corresponding aldehyde **1** (3 mmol) and chloriodomethane (0.436 mL, 6 mmol) in THF (10 mL) was added methyllithium (4 mL of 1.5 M solution in diethyl ether, 6 mmol) dropwise over 5 min and under nitrogen. After stirring at -78°C for 30 min, lithium powder (0.216 g, 30 mmol) was added at -45°C , and the mixture was stirred for 8 h at the same temperature. The resulting mixture was treated with ice and after the usual work-up, crude allylamine **4** was obtained. Column flash chromatography over silica gel (15:1 hexane:ethyl acetate) provided the corresponding pure allylamine **4**.

In conclusion, we have described a direct, easy and simple transformation without racemization, of α -amino aldehydes into enantiopure allylamines.

Acknowledgements

We thank II Plan Regional de Investigación del Principado de Asturias (PBP-PGI98-04) and Ministerio de Educación y Cultura (PB97-1278) for financial support. E.R. thanks the Ministerio de Educación y Cultura for a predoctoral fellowship.

References

1. Franciotti, M.; Mordini, A.; Taddei, M. *Synlett* **1992**, 137–138.
2. (a) Luly, J. R.; Dellaria, J. F.; Plattner, J. J.; Soderquist, J. L.; Yi, N. *J. Org. Chem.* **1987**, *52*, 1487–1492. (b) Romeo, S.; Rich, D. H. *Tetrahedron Lett.* **1993**, *34*, 7187–7190. (c) Albeck, A.; Persky, R. *J. Org. Chem.* **1994**, *59*, 653–657. (d) Branat, J.; Kvarnström, I.; Classon, B.; Samuelson, B.; Nilroth, U.; Danielson, H.; Karlén, A.; Halberg A. *Tetrahedron Lett.* **1997**, *38*, 3483–3486.
3. Kobayashi, S.; Isobe, T.; Ohno, M. *Tetrahedron Lett.* **1984**, *25*, 5079–5082.
4. Nishi, T.; Moreisawa, Y. *Heterocycles* **1989**, *29*, 1835–1842.
5. Reetz, M. T. *Chem. Rev.* **1999**, *99*, 1121–1162.
6. (a) Fehrentz, J. A.; Castro, B. *Synthesis* **1983**, 676–678. (b) Rettle, K. E.; Hommick, C. F.; Ponticello, G. S.; Evans, B. E. *J. Org. Chem.* **1982**, *47*, 3016–3018. (c) Lubell, W. D.; Rapoport, H. *J. Am. Chem. Soc.* **1988**, *110*, 7447–7455.
7. (a) Barluenga, J.; Baragaña, B.; Alonso, A.; Concellón, J. M. *J. Chem. Soc., Chem. Commun.* **1994**, 969–970. (b) Barluenga, J.; Baragaña, B.; Concellón, J. M. *J. Org. Chem.* **1995**, *60*, 6696–6699.
8. UV detector 325 nm, 0.8 mL/min, 400:1 hexane:isopropanol; t_R : 6.38 min.