

Tetrahedron Letters 40 (1999) 21-24

TETRAHEDRON LETTERS

## Manganese Triacetate Mediated Oxidation of Hantzsch 1,4-Dihydropyridines to Pyridines

Rajender S. Varma<sup>\*</sup> and Dalip Kumar

Department of Chemistry and Texas Research Institute for Environmental Studies (TRIES), Sam Houston State University, Huntsville, TX 77341-2117, U. S. A. Fax: (409)-294-1585; E-mail: CHM\_RSV@SHSU.EDU

Received 4 September 1998; revised 12 October 1998; accepted 13 October 1998

Abstract: A general and practical route for the high yield oxidative conversion of readily accessible 1,4-dihydropyridines to the corresponding pyridines is described using a relatively benign oxidant, manganese triacetate. © 1998 Elsevier Science Ltd. All rights reserved.

The oxidation of Hantzsch 1,4-dihydropyridines to the corresponding pyridines has been extensively studied in view of the pertinence of this reaction to the metabolism of Hantzsch esters and the calcium channel blocking drugs used in the treatment of various cardiovascular disorders.<sup>1</sup> The reaction has also been used to study the biologically significant NADH redox processes.<sup>2</sup> Consequently, this aromatization reaction continues to attract the attention of researchers for the discovery of milder and general protocols applicable to a wide range of 1,4-dihydropyridines. Many of the reported reagents<sup>3</sup> involve the use of strong oxidants such as KMnO<sub>4</sub>,<sup>3b</sup> CrO<sub>3</sub>,<sup>3c</sup> HNO<sub>3</sub>,<sup>3d</sup> pyridinium chlorochromate (PCC),<sup>3e</sup> ceric ammonium nitrate (CAN),<sup>3f</sup> bentonite clay-supported manganese dioxide<sup>3g</sup> and more recently, bismuth nitrate Bi(NO<sub>3</sub>)<sub>3</sub>.<sup>3h</sup> However, this aromatization reaction with most of these reagents leads to dealkylation at the 4-position or formation of side products.<sup>3b</sup>,<sup>g</sup> In some other instances only moderate to poor yields are obtained.<sup>3b</sup>,<sup>e</sup>,<sup>g</sup>

In view of the above limitations, we decided to develop a practical and general approach for this oxidative conversion using a mild oxidant, manganese triacetate, which has been increasingly utilized in a wide variety of organic reactions<sup>4</sup> namely lactonization of olefins,<sup>5</sup> oxidative cycloadditions,<sup>6</sup> deoximation of carbonyl compounds<sup>7</sup> and oxidative cyclization of phenolic Schiff's bases.<sup>8</sup> Herein, we report our results on an expeditious manganese triacetatemediated oxidation of 1,4-dihydropyridines to pyridines. Acetic acid is the solvent of choice in this conversion for the optimum yield of products; the reaction in various other solvents such as dichloromethane, methanol and acetonitrile is very slow and does not lead to completion even after 24 h. Manganese triacetate being a one-electron oxidant, two equivalents of manganese triacetate are required for the completion of the reaction that generates pyridine derivatives in excellent yields within 15-90 minutes (Table).

The general applicability, versatility and scope of this reaction is defined by using various substrates which illustrate the tolerance of several substituents namely alkyl, aryl, styryl and heterocycles at the 4-position. The salient features of this reaction are milder reaction conditions, excellent yields of the products and the stability of the substituents at 4-position which are normally dealkylated during aromatization by existing methods e.g. oxidants such as KMnO4,<sup>3b</sup> and clay-MnO<sub>2</sub>.<sup>3g</sup> The general exception, typical of all known oxidants described in the modern literature, is the case of secondary alkyl (isopropyl) and benzyl groups. In view of their electron releasing ability and the stability of the corresponding radicals these substituents are expelled with the formation of dealkylated products.<sup>3g</sup> In the case of reaction with diethyl 4-(4-hydroxy phenyl)-2,6-dimethyl-1,4-dihydropyridine-3,5-dicarboxylate, 1f, desired pyridine, 2f is readily obtained; nitrated side products are normally formed in reaction with bismuth nitrate.<sup>3h</sup> Our results on nearly quantitative conversion of various substrates to the corresponding pyridine derivatives are summarized in the Table.

Presumably, the oxidation of 1,4-dihydropyridine is initiated by a single electron transfer to Mn (III) that produces Mn (II) and a radical cation I which subsequently loses a proton to generate a radical II. The second mole of Mn (III) then oxidizes radical II to the protonated pyridine III. Finally, the pyridinium species III loses a proton to deliver the desired pyridine derivative (Scheme 1).



Scheme 1

## Experimental

Manganese triacetate was obtained from Aldrich Chemical Co. and the 1,4dihydropyridines were prepared according to the literature procedure.<sup>9</sup>

**General Procedure:** The preparation of diethyl 2,6-dimethyl-3,5-pyridinedicarboxylate, 2a, is representative of the general procedure employed. To a solution of manganese triacetate (0.536 g, 2 mmol) in acetic acid (5 mL) was added diethyl 1,4-dihydro-2,6-dimethyl-3,5-pyridinedicarboxylate 1a, (0.253 g, 1 mmol). The reaction mixture was stirred at room temperature for 20 min. After completion of the reaction, as indicated by TLC examination, manganese diacetate was filtered off and the reaction mixture poured into water. The contents were then neutralized by NaHCO<sub>3</sub>, extracted with dichloromethane (2 x 10 mL) and dried over anhydrous sodium sulfate. The solvent was removed by distillation under reduced pressure and the resulting crude product obtained was crystallized from ethanol to afford pure 2a in 98 % yield (0.248 g); m.p. 71 °C (lit.<sup>10</sup> m.p. 70-71 °C).



Entry	R	Time	Yield	m. p. (°C) <sup>a</sup>	
		(min)	(%) <sup>b</sup>	Observed	Reported
2a	Н	20	98	71	70-71 <sup>10</sup>
<b>2</b> b	CH <sub>3</sub>	20	97	liq	liq <sup>11</sup>
2c	CH <sub>3</sub> CH <sub>2</sub>	20	97	liq	liq <sup>10</sup>
2d	C <sub>6</sub> H <sub>5</sub>	35	98	63-64	62-6411
2e	С <sub>6</sub> Н5-СН=СН	60	92	162-63	162-65 <sup>11</sup>
2f	4-HOC <sub>6</sub> H <sub>4</sub>	30	90	171	171-73 <sup>12</sup>
2g	4-CH3C6H4	35	98	72-73	<b>72-7</b> 3 <sup>11</sup>
2h	4-CH3OC6H4	40	94	51-53	51-53 <sup>12</sup>
<b>2i</b>	4-NO2C6H4	60	91	114-15	11513
2j	2-Furyl	90	94	liq	liq <sup>14</sup>
2k	2-Thienyl	90	96	liq	liq <sup>15</sup>

Table: Oxidation of 1,4-dihydropyridines to pyridine derivatives using manganese triacetate

<sup>a</sup>Products exhibited physical and spectral properties in accordance with the assigned structures. <sup>b</sup>Yields refer to the isolated pure products. In conclusion, we have developed a general and practical route for the oxidative conversion of 1,4-dihydropyridines to the corresponding pyridine derivatives in excellent yields using a relatively benign oxidant, manganese triacetate.

## Acknowledgment

We are grateful for financial support to the Texas Advanced Research Program (ARP) in chemistry (Grant # 003606-023) and Texas Research Institute for Environmental Studies (TRIES).

## References

- 1. (a) Janis, R. A.; Triggle, D. J. J. Med. Chem. 1983, 26, 775;
  - (b) Gordeev, M. F.; Patel, D. V.; Gordon, E. M. J. Org. Chem. 1996, 61, 924;
  - (c) Bossert, F.; Vater, W. Med. Res. Rev. 1989, 9, 291;
  - (d) Sausins, A.; Duburs, G. Heterocycles 1988, 27, 269;
  - (e) Kazda, S.; Towart, R. Br. J. Pharmacol. 1981, 72, 582P
- (a) Kill, R. J.; Widdowson, D. A. In *Biorganic Chemistry*; van Tamelen, E. E. Ed. Academic Press (NY) Vol. 4, 1978; pp 239-75;
  - (b) Stout, D. M.; Meyers, A. I. Chem. Rev. 1982, 82, 223.
- 3. (a) Balogh, M.; Hermecz, I.; Meszaros, Z.; Laszlo, P. Helv. Chim. Acta. 1984, 67, 2270;
  - (b) Vanden Eynde, J. J.; D'orazio, R.; Yves Van, H. Tetrahedron 1994, 50, 2479;
  - (c) Grinsteins, E.; Stankevics, B.; Duburs, G. Kim. Geterotsikl. Soedin, 1967, 1118: Chem. Abstr. 69, 77095;
  - (d) Bocker, R. H.; Guengerich, F. P. J. Med. Chem. 1986, 28, 1596;
  - (e)Vanden Eynde, J-J.; Mayence, A.; Maquestiau, A. Tetrahedron, 1992, 48, 463;
  - (f) Pfister, J. R. Synthesis 1990, 689;
  - (g) Delgado, F.; Alvarez, C.; Garcia, O.; Penieres, G; Marquez, C. Synth. Commun. 1991, 21, 2137;
  - (h) Mashraqui, S. H.; Karnik, M. A. Synthesis 1998, 713.
- 4. (a) Criegee, R. Angew. Chem. 1958, 70, 173; (b) Snider, B. B. Chem. Rev. 1996, 96, 339.
- 5. (a) Fritad, W. E.; Peterson, J. R. J. Org. Chem. 1985, 50, 10;
- (b) Fristad, W. E.; Peterson, J. R.; Ernst, J. R. J. Org. Chem. 1985, 50, 3143.
- 6. Snider, B. B.; Han, L.; Xie, C. J. Org. Chem. 1997, 62, 6978.
- 7. Demir, A. S.; Tanyeli, C.; Altinel, E. Tetrahedron Lett. 1997, 38, 7267.
- 8. Varma, R. S.; Kumar, D. J. Heterocyclic Chem. 1998, (in press).
- 9. Bocker, R. H.; Guengerich, F. P. J. Med. Chem. 1986, 29, 1596.
- 10. Engelman, F. Ann. 1885, 231, 37.
- 11. Loev, B.; Snader, K. M. J. Org. Chem. 1965, 30, 1914.
- 12. Emmert, D.; Diefenbach, E.; Eck, R. Ber. 1927, 60, 2220.
- 13. Cook, A. H.; Heilboon, I. M.; Steger, L. J. Chem. Soc. 1943, 413.
- 14. Heiber, F. Ber. 1892, 25, 2405.
- 15. Grischkewitsch-Trochimowski, G.; Mazurewitsch, M. Chem. Zentr. 1912, 6, 1562.