

## Novel Rearrangement of the 9H-Fluorene System into the 1H- and 3H-Fluorene Systems


By N. P. BUU-HOÏ,\* C. MARIE, F. PÉRIN, and P. JACQUIGNON

(Institut de Chimie des Substances Naturelles du C.N.R.S., 91-Gif-sur-Yvette, France)

**Summary** Two dehydroanthrones, 11H-indeno[1,2-*b*]-anthracen-11-one and 7H-indeno[2,1-*a*]-anthracen-7-one, have been obtained in the pyrolysis of 2-*o*-toluoylfluorene—a first example of a rearrangement of the ordinary 9H-fluorene system into the tautomeric 3H- and 1H-fluorene ones.

Pyrolysis of 2-*o*-toluoylfluorene (I) at 380–390° and chromatographic separation (on silica column) of the pyrolysate showed that this Elbs reaction had produced four compounds: (i) two pentacyclic hydrocarbons C<sub>21</sub>H<sub>14</sub> arising from normal dehydration, 13H-indeno[1,2-*b*]-anthracene (II), m.p. 319° (lit.<sup>1</sup> 319°) and 13H-indeno[2,1-*a*]-anthracene (III), m.p. 250°; and (ii) two oxygen-containing compounds C<sub>21</sub>H<sub>12</sub>O, one of them gold-yellow, m.p. 246° (absorption bands in the visible region at 406, 432, and 459 cm<sup>-1</sup>), and the other crimson, m.p. 205° (bands in visible at 403, 413, 466, and 520 cm<sup>-1</sup>), resulting from dehydrogenation of (I). To these isomeric dehydroanthrones we assign the 11H-indeno[1,2-*b*]-anthracen-11-one (IV) and 7H-indeno[2,1-*a*]-anthracen-7-one (V) structures, on the grounds of n.m.r. spectroscopy in C<sub>6</sub>D<sub>6</sub> (which showed the absence of methylene protons), mass spectrometry (for both compounds, base peak *m/e* 280, and fragmentation pattern characteristic of anthrones, *i.e.* loss of carbon monoxide, to give the peak *m/e* 252), and their deep-violet halochromism in sulphuric acid indicative of highly conjugated cyclic ketones.<sup>2</sup> The known differences in absorption spectra between the linear and the angular dibenzofluorenones<sup>3</sup> suggest the linear structure (IV) for

the light-coloured dehydroanthrone and the angular structure (V) for the deeper isomer.



Compounds (IV) and (V) are derivatives of, respectively, 3H- and 1H-fluorene, the hitherto unknown tautomers of fluorene itself, and their isolation demonstrates for the first time the formation of dehydroanthrones in Elbs reactions,<sup>3</sup> and the possibility of rearrangements of the double-bond system of 9H-fluorene in which the central methylene group is involved and of which the driving force is the tendency to undergo total conjugation.

We thank Dr. B. Das of the Mass Spectrometry Department of this Institute for the mass spectra.

(Received, April 20th, 1970; Com. 575.)

<sup>1</sup> E. de Barry Barnett, N. F. Goodway, and J. W. Watson, *Ber.*, 1933, **66**, 1876; O. Kruber, *ibid.*, 1937, **70**, 1556.

<sup>2</sup> Cf. R. H. Martin, *J. Chem. Soc.*, 1941, 679; *Helv. Chim. Acta*, 1947, **30**, 620; G. M. Badger, *J. Chem. Soc.*, 1941, 535.

<sup>3</sup> Anthrones have recently been found to be formed regularly in Elbs reactions: N.P. Buu-Hoï, C. Marie, and P. Jacquignon, *Bull. Soc. chim. France*, 1970, 1012.