

tion of Sn-H bonds. The reaction products were apparently mixtures, containing no detectable amount of triphenylstannane.

Acknowledgment.—The author is grateful to Dr. Eugene G. Rochow for helpful discussions and

advice, and to the National Science Foundation for financial support.

MALLINCKRODT CHEMICAL LABORATORY
HARVARD UNIVERSITY
CAMBRIDGE 38, MASS.

COMMUNICATIONS TO THE EDITOR

THE SYNTHESIS OF AN ARGINYL PEPTIDE

Sir:

The synthesis of peptides containing arginine has presented unusual difficulties, and only a few dipeptides have been reported. No peptide in which the carboxyl group of arginine is linked to another amino acid has been synthesized.¹ It has now been found that the pyrophosphite method² can be used for this purpose.

Carbobenzoxy-L-arginine hydrobromide³ (0.010 mole) and methyl L-leucinate (0.010 mole) were added to 7 cc. of diethyl phosphite, then tetraethyl pyrophosphite (0.020 mole) was added. After 30 minutes heating on a steam-bath, methyl carbobenzoxy-L-arginyl-L-leucinate hydrobromide hydrate was precipitated as a gum by 100 cc. of anhydrous ether. This was crystallized by dissolving in 5 cc. of methanol, bubbling in hydrogen bromide a moment, then adding 40 cc. of anhydrous ether; yield 3.06 g. (57%). A further 0.47 g. was obtained from the original filtrate on dilution with ether, giving 3.53 g. in all (66%). Solution in 10 cc. of methanol and dilution with 40 cc. of water yielded 2.79 g. (52%), m.p. 90–92°, $[\alpha]^{23}_D -20.3^\circ$ (c 2, methanol).

Anal. Calcd. for $C_{21}H_{36}N_6O_6Br$: C, 47.2; H, 6.79; N, 13.1; Br, 15.0. Found: C, 47.4; H, 6.99; N, 13.3; Br, 15.2.

Carbobenzoxy-L-arginyl-L-leucine was obtained by heating 2.83 g. of the methyl ester hydrobromide hydrate on a steam-bath in 28 cc. of *N* hydrobromic acid for an hour, making slightly alkaline with ammonium hydroxide and chilling. The crystalline product was recrystallized from 30 cc. of *N*/6 hydrobromic acid by the addition of ammonium hydroxide; yield 1.20 g. (54%), m.p. 223–224° dec., $[\alpha]^{24}_D -26^\circ$ (c 2, 0.4 *N* HBr). *Anal.* Calcd. for $C_{20}H_{31}N_5O_5$: C, 57.0; H, 7.41; N, 16.6. Found: C, 57.2; H, 7.64; N, 16.6.

L-Arginyl-L-leucine hydrobromide hydrate was obtained by hydrogenation with a palladium catalyst of 1.06 g. of the carbobenzoxy derivative in 40 cc. of water plus 2.5 cc. of 1.04 *N* HBr. A crystalline product obtained by evaporation of the filtered

solution under vacuum was washed out with acetone, then recrystallized by dissolving in 4 cc. of water and slowly adding 40 cc. of acetone; yield 0.78 g. (80%), m.p. 162–163° dec., $[\alpha]^{24}_D +8.6^\circ$ (c 2, water).

Anal. Calcd. for $C_{12}H_{23}N_5O_4Br$: C, 37.3; H, 7.3; N, 18.1; Br, 20.7. Found: C, 37.1; H, 7.2; N, 18.2; Br, 20.7.

Paper chromatography gave an R_f value of 0.58 in a butanol–water–acetic acid (5:4:1) system, and showed the presence of arginine and leucine in an acid hydrolysate.

CHEMOTHERAPY DIVISION
STAMFORD RESEARCH LABORATORIES
AMERICAN CYANAMID COMPANY GEORGE W. ANDERSON
STAMFORD, CONNECTICUT

RECEIVED SEPTEMBER 2, 1953

TWO DIMENSIONAL PHASE TRANSITION OF ETHANE ON SODIUM CHLORIDE

Sir:

The chief evidence for first-order phase transitions of films adsorbed on solid surfaces has been the adsorption isotherms measured by Jura, *et al.*, for *n*-heptane on graphite,¹ ferric oxide² and reduced silver powder.³ Adsorption isotherms for *n*-heptane on the same solids (but not the same samples) have subsequently been repeated by Smith⁴ and by Young, Beebe and Bienes,⁵ who report that their work provides no evidence of first-order phase transitions in those systems. Their papers support the trend to interpret all experimentally observed discontinuities in adsorption isotherms as caused by slow rates or diffusion inside the sample or errors in the apparatus.

The chief remaining evidence for a first-order phase transition of an adsorbed film on a solid surface now devolves on the adsorption isotherms of ethane on sodium chloride and on potassium chloride crystals at 90°K., reported by Ross and Boyd,⁶ and since verified in this laboratory on a different sample of sodium chloride at 90°K. by Mr. W.

(1) G. Jura, W. D. Harkins and E. H. Loeser, *J. Chem. Phys.*, **14**, 344 (1946).

(2) G. Jura, E. H. Loeser, P. R. Basford and W. D. Harkins, *ibid.*, **14**, 117 (1946).

(3) Jura, *et al.*, *ibid.*, **13**, 535 (1945).

(4) R. N. Smith, *THIS JOURNAL*, **74**, 3477 (1952).

(5) D. M. Young, R. A. Beebe and H. Bienes, *Trans. Faraday Soc.*, **49**, 1086 (1953).

(6) S. Ross and G. E. Boyd, "New Observations on Two-Dimensional Condensation Phenomena," MDDC Report 864, 1947.

(1) J. S. Fruton, "Advances in Protein Chemistry," Vol. V, Academic Press, Inc., New York, N. Y., 1949, p. 64.

(2) G. W. Anderson, J. Blodinger and A. D. Welcher, *THIS JOURNAL*, **74**, 5309 (1952).

(3) Prepared by R. Janice Joyce, of these laboratories; m.p. 177–179.5°, $[\alpha]^{24}_D -6.9^\circ$ (c 2, water); calcd. for $C_{14}H_{21}N_4O_4Br$: 20.8% Br. Found: 20.6% Br.

(4) Melting points were taken on a calibrated Fisher-Johns block.