


This article was downloaded by: [The Aga Khan University]

On: 05 December 2014, At: 22:13

Publisher: Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954

Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Phosphorus, Sulfur, and Silicon and the Related Elements

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/gpss20>

SYNTHESIS OF 1, n-BIS (2,5-DIPHENYLPHOSPHOL-1-YL) ALKANES

Sandrine Bousquet^a, Jean-Jacques Brunet^a,
Thierry Courcet^a & Denis Neibecker^a

^a Laboratoire de Chimie de Coordination du CNRS, Unité n°8241, liée par conventions à l'Université Paul Sabatier et à l'Institut National Polytechnique, 205 route de Narbonne, 31077, Toulouse, Cedex, France

Published online: 24 Sep 2006.

To cite this article: Sandrine Bousquet, Jean-Jacques Brunet, Thierry Courcet & Denis Neibecker (1998) SYNTHESIS OF 1, n-BIS (2,5-DIPHENYLPHOSPHOL-1-YL) ALKANES, *Phosphorus, Sulfur, and Silicon and the Related Elements*, 142:1, 117-124, DOI: [10.1080/10426509808029671](https://doi.org/10.1080/10426509808029671)

To link to this article: <http://dx.doi.org/10.1080/10426509808029671>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any

losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

SYNTHESIS OF 1,*n*-BIS (2,5-DIPHENYLPHOSPHOL-1-YL)ALKANES

SANDRINE BOUSQUET, JEAN-JACQUES BRUNET,
THIERRY COURCET and DENIS NEIBECKER*

Laboratoire de Chimie de Coordination du CNRS, Unité n°8241, liée par conventions à l'Université Paul Sabatier et à l'Institut National Polytechnique, 205 route de Narbonne, 31077 Toulouse Cedex (France)

(Received 04 August, 1998)

1,*n*-bis (2,5-diphenylphospholyl)alkanes (*n* = 2–6) are easily prepared in a one pot high-yield reaction and fully characterized by multinuclear NMR spectroscopy.

Keywords: Diphosphines; Phospholes; Diphospholes; Synthesis; NMR

INTRODUCTION

Chelating compounds especially diphosphanes have found extensive applications as ligands in organometallic chemistry and in homogeneously transition-metal catalyzed reactions such as hydrogenation, hydroformylation, cross-coupling, etc...^[1] Many optically-active diphosphanes have been designed for use as ligands in enantioselective catalysis, most of them containing the diphenylphosphanyl moiety.^[2]

Some years ago, we initiated a program aimed at finding applications of phospholes in homogeneous catalysis and reported that 1,2,5-triphenylphosphole is a very efficient ligand for the rhodium-catalyzed hydroformylation of aliphatic, aromatic and functionalized olefins.^[3] Since diphosphanes (diphosphines and diphosphites) have been found to be better ligands than monophosphanes for transition metal-catalyzed hydroformylation of olefins,^[4] we sought to investigate 1,*n*-bis(2,5-

* Corresponding author. Telefax: +33(0) 561 553 003. E.mail: neibeker@lcc-toulouse.fr.

diphenylphosphol-1-yl)alkanes. Many diphosphines, optically active or not, containing two dibenzophosphol-1-yl^[5] groups are known and several diphosphines containing two dinaphthophosphol-1-yl^[6] groups have been recently reported. However examples of simple 1,*n*-bis(phosphol-1-yl)alkanes are less common. Literature mentions the synthesis of 1,2-bis(phosphol-1-yl)ethane (unreported yield),^[7] of 1,*n*-bis(2,3,4,5-tetraphenylphosphol-1-yl)alkanes (*n* = 1, 2; 12–23% yield)^[8] and of 1,*n*-bis(3,4-dimethylphosphol-1-yl)alkanes (*n* = 1–4; 38–65% yield).^[9] In addition we have already described the synthesis of optically active 1,4-diphospholes bearing 2,5-diphenylphosphol-1-yl or 3,4-dimethylphosphol-1-yl groups.^[10] We now wish to report the high-yield synthesis and NMR characterization of the still unknown simple 1, *n*-bis(2,5-diphenylphosphol-1-yl) alkanes.

RESULTS AND DISCUSSION

The synthesis of 1,*n*-bis(2,5-diphenylphosphol-1-yl) alkanes **2–6** was performed from 2,5-diphenylphosphol-1-yllithium **1** using the conventional methodology outlined in scheme 1.


Generation of **1** was achieved, as usual, by cleavage of the exocyclic phosphorus-carbon bond of the readily available 1,2,5-triphenylphosphole^[11] with lithium in THF,^[8] followed by selective destruction of the concomitantly generated phenyllithium by *tert*-butyl chloride.^[12] Analysis of the reaction mixture by ³¹P{¹H} NMR spectroscopy revealed the clean formation of **1** (THF solution, δ = 83.6 ppm) sometimes contaminated by a small amount of 2,3,5-triphenylphosphollythium^[13] (δ = 106 ppm). The reaction of **1** and 1,*n*-dichloroalkanes (*n* = 2–4) either did not proceed at room temperature or gave rise at reflux to a mixture of products (³¹P{¹H} NMR analysis) from which the diphospholes **2–4** could be isolated in low yields (25–30%). However, using 1,*n*-dibromoalkanes (*n* = 2–6) allowed a clean reaction at room temperature and resulted in the isolation of diphospholes **2–6** with the yields reported in scheme 1.

The phospholes **2–6** have been characterized by mass spectrometry, elemental analysis (see experimental section) and thoroughly studied by multinuclear NMR spectroscopy.

³¹P{¹H} NMR data (Table I) indicate that the chemical shift for **2–6** is comparable to that of 1,2,5-triphenylphosphole (δ ³¹P = +3.0 ppm)^[14] and

CONCLUSION

1,*n*-bis(2,5-diphenylphosphol-1-yl)alkanes (*n* = 2–6) can be prepared in good yield by an efficient one pot experimental procedure. Work is in progress to study their coordination chemistry and their potential applications as ligands in transition metal-catalyzed hydrofunctionalisation of olefins.



Diphospholes	2 ^(a)	3 ^(a)	4 ^(a)	5 ^(b)	6 ^(c)
³¹ P{ ¹ H} δ (ppm)	+6	+2.1	+2.5	+3.3	+3.5
H ³	7.10 (d) ³ J _{H-P} = 10 Hz	6.90 (d) ³ J _{H-P} = 10 Hz	7.10 (d) ³ J _{H-P} = 10 Hz	6.95 (d) ³ J _{H-P} = 10 Hz	7.11 (d) ³ J _{H-P} = 10 Hz
H ¹	1.55 (t) ³ J _{H-P} = 6 Hz	1.57 (m)	1.60 (m)	1.51 (m)	1.65 (m)
H ²	1.55 (t) ³ J _{H-P} = 6 Hz	0.80 (m)	0.80 (m)	0.76 (dqt) ^(d) ³ J _{H-H} = 8 Hz ³ J _{H-P} = 8 Hz	0.77 (m)
H ³	-	1.57 (m)	-	0.44 (qt) ^(d) ³ J _{H-H} = 8 Hz	0.67 (m)

(a) Bruker AC 200, in CDCl₃ solution at 200.13 and 81.015 MHz for ¹H and ³¹P{¹H} NMR, respectively.

(b) Bruker AMX 400, in C₆D₆ solution at 400.14 and 161.99 MHz for ¹H and ³¹P{¹H} NMR, respectively. In CDCl₃ solution, H² and H³ are superimposed at 0.82 ppm.

(c) Bruker AMX 400, in CDCl₃ solution at 400.14 and 161.99 MHz for ¹H and ³¹P{¹H} NMR, respectively.

(d) qt = quintuplet.

EXPERIMENTAL SECTION

General Procedure

All sample manipulations were carried out under argon using standard Schlenk tube and vacuum techniques. Solvents and reagents were purified


according to literature procedures^[15] and stored under argon. Argon U (L'Air Liquide) was used after passage through 3-Å molecular sieves. Lithium (wires, Aldrich), *tert*-butyl chloride (Fluka), 1,2-dibromoethane (Prolabo), 1,3-dibromopropane (Fluka), 1,4-dibromobutane, 1,5-dibromopentane and 1,6-dibromohexane (Aldrich), were used as received. 1,2,5-triphenylphosphole was prepared by a reported procedure.^[11]

¹H, ¹³C, and ³¹P NMR spectra were recorded in CDCl₃ or C₆D₆ solutions on Bruker AC 80 (³¹P : 32.43 MHz) (analysis of reaction mixtures), AC 200 (¹H : 200.13 MHz, ³¹P : 81.015 MHz, ¹³C : 50.32 MHz) or AMX 400 (¹H : 400.14 MHz, ³¹P : 161.99 MHz, ¹³C : 100.62 MHz) spectrometers. ¹H and ¹³C NMR chemical shifts are referenced to tetramethylsilane assigning the CDCl₃ resonances at 7.27 and 77.0 ppm, respectively and the C₆D₆ resonances at 7.16 and 128.0 ppm, respectively. ³¹P NMR chemical shifts are referenced to external 85% H₃PO₄ in D₂O.

Mass spectra were obtained on a Nermag R10/10 (EI, 70 eV; CI, NH₃) spectrometer and Elemental analyses were performed on a Perkin Elmer 2400 serie II apparatus. Melting point (uncorrected) were measured on a Buchi SMP 20 apparatus. All the compounds synthesized except **2** show a change in color between 90 and 100°C but melt at the reported temperature.

1,2-bis(2,5-diphenylphosphol-1-yl)ethane (**2**)

A 250-mL Schlenk flask containing a teflon-coated magnetic stirring bar was charged with 1,2,5-triphenylphosphole (2.7 g, 8.6 mmol) and submitted to three vacuum-argon cycles. Freshly distilled THF (50 mL) was syringed into the Schlenk flask and the mixture cooled to 0°C. Small cuts of lithium (0.15 g, 21.6 mmol) were added and the reaction medium stirred at 10–15 °C for 5 h. After filtration through celite into another Schlenk flask to remove unreacted lithium, *tert*-butyl chloride (1.2 mL, 11.2 mmol) was added and the filtrate stirred for 16 h. 1,2-dibromoethane (0.37 mL, 4.3 mmol) was then added and the reaction medium stirred for 5 h. Addition of distilled water (60 mL) gave a two-phase system which was vigorously stirred. Water was decanted with a syringe and THF (20 mL) and water (60 mL) were added. This procedure was repeated twice. The THF solution was evaporated to dryness to leave an orange solid. Recrystallization from methanol afforded a yellow solid which was dried under reduced pressure up to constant weight (1.9 g, 89%). m.p. >245°C. – MS (EI, 70 eV): *m/z* (%) = 499 (23), 498 (69) [M⁺]. 470 (22) [M⁺-C₂H₄], 235 (53). – C₃₄H₂₈P₂ (498.5); calc. C 81.91, H 5.66; found C 81.13, H 5.82.

TABLE II $^{13}\text{C}\{^1\text{H}\}$ NMR data of diphospholes **2-6**

	C^1	C^2	C^3	C^4	C^5	C^6	C^7
2 ^(a)	18.9 (dd) $^1J_{\text{C-P}} = 22 \text{ Hz}$ $^2J_{\text{C-P}} = 4 \text{ Hz}$	149.7 (d) $^1J_{\text{C-P}} = 3 \text{ Hz}$	132.1 (d) $^2J_{\text{C-P}} = 9 \text{ Hz}$	136.2 (d) $^2J_{\text{C-P}} = 17 \text{ Hz}$	125.6 (d) $^3J_{\text{C-P}} = 9 \text{ Hz}$	128.8 (s)	127.0 (s)
3 ^(a)	24.5 (dd) $^1J_{\text{C-P}} = 17 \text{ Hz}$ $^3J_{\text{C-P}} = 6 \text{ Hz}$	150.5 (s)	132.0 (d) $^2J_{\text{C-P}} = 8 \text{ Hz}$	136.2 (d) $^2J_{\text{C-P}} = 17 \text{ Hz}$	125.6 (d) $^3J_{\text{C-P}} = 10 \text{ Hz}$	128.8 (s)	127.0 (s)
4 ^(a)	23.9 (d) $^1J_{\text{C-P}} = 16 \text{ Hz}$	26.3 (d) $^3J_{\text{C-P}} = 6 \text{ Hz}$	131.7 (d) $^2J_{\text{C-P}} = 8 \text{ Hz}$	136.9 (d) $^2J_{\text{C-P}} = 17 \text{ Hz}$	126.2 (d) $^3J_{\text{C-P}} = 9 \text{ Hz}$	128.7 (s)	127.0 (s)
5 ^(b)	23.9 (d) $^1J_{\text{C-P}} = 15 \text{ Hz}$	24.6 (s)	132.6 (d) $^2J_{\text{C-P}} = 8 \text{ Hz}$	138.2 (d) $^2J_{\text{C-P}} = 17 \text{ Hz}$	127.2 (d) $^3J_{\text{C-P}} = 10 \text{ Hz}$	129.6 (s)	127.8 (s)
6 ^(b)	24.0 (d) $^1J_{\text{C-P}} = 15 \text{ Hz}$	24.9 (s)	131.8 (d) $^2J_{\text{C-P}} = 9 \text{ Hz}$	137.1 (d) $^2J_{\text{C-P}} = 17 \text{ Hz}$	126.3 (d) $^3J_{\text{C-P}} = 10 \text{ Hz}$	128.8 (s)	127.1 (s)

(a) : Bruker AC 200, in CDCl_3 solution at 50.32 MHz. (b) : Bruker AMX 400, in CDCl_3 solution at 100.62 MHz.

1,3-bis(2,5-diphenylphosphol-1-yl)propane (3)

The same procedure was followed by using 1,3-dibromopropane (0.43 mL, 4.3 mmol). **3**: yield 1.95 g (91%) yellow solid, m.p. 133°C – MS (EI, 70 eV): m/z (%) = 512 (51) [M^+], 277 (100), 133 (36). – $C_{35}H_{30}P_2$ (512.5): calc. C 82.02, H 5.90; found C 81.44, H 5.89.

1,4-bis(2,5-diphenylphosphol-1-yl)butane (4)

The same procedure was followed by using 1,4-dibromobutane (0.51 mL, 4.3 mmol). **4**: yield 2.15 g (93%) yellow solid, m.p. 142°C – MS (EI, 70 eV): m/z (%) = 526 (51) [M^+], 292 (23), 291 (100), 133 (35). – $C_{36}H_{32}P_2$ (526.6): calc. C 82.11, H 6.13; found C 81.44, H 6.57.

1,5-bis(2,5-diphenylphosphol-1-yl)pentane (5)

The same procedure was followed by using 1,5-dibromopentane (0.919 g, 4.00 mmol). **5** was recrystallized twice from CH_2Cl_2 -MeOH. **5**: yield 1.5 g (70%) yellow solid, m.p. 135°C – MS (CI, NH_3): m/z (%) = 542 (42), 541 (100) [MH^+]. – $C_{37}H_{34}P_2$ (540.6): calc. C 82.20, H 6.34; found C 80.52, H 6.49.

1,6-bis(2,5-diphenylphosphol-1-yl)hexane (6)

The same procedure was followed by using 1,6-dibromohexane (0.977 g, 4.00 mmol). **6** yield 1.04 g (47%) yellow solid, m.p. 133°C – MS (CI, NH_3): m/z (%) = 556 (45), 555 (100) [MH^+]. – $C_{38}H_{36}P_2$ (554.7): calc. C 82.29, H 6.54; found C 82.01, H 6.82.

Acknowledgements

The authors wish to thank the Centre National de la Recherche Scientifique (CNRS) for financial support and the Ministère de l'Éducation Nationale, de l'Enseignement Supérieur et de la Recherche for a fellowship to S.B. G. Commenges and F. Lacassin are gratefully acknowledged for the registration of NMR spectra, L. Noé for performing elemental analyses, C. Claparols and S. Richelme for recording mass spectra.

References

- [1] *Homogeneous Catalysis with Metal Phosphine Complexes*, L.H. Pignolet (Ed) (Plenum Press, New York, 1983); *The Chemistry of Organophosphorus Compounds*, F.R. Hartley and S. Patai (Eds) (John Wiley, New York, 1990) vol. 1; *Applied Homogeneous Catalysis with Organometallic Compounds*, B. Cornils and W.A. Herrmann (Eds) (VCH, Weinheim, 1996) vols 1 and 2.
- [2] H. Brunner and R. Zettlmeier, *Handbook of Enantioselective Catalysis*, (VCH, Weinheim, 1993) vols 1 and 2; *Catalytic Asymmetric Synthesis*, I. Ojima (Ed) (VCH, New York, 1993).
- [3] D. Neibecker and R. Réau, *J. Mol. Catal.*, **57**, 153 (1989).
D. Neibecker, R. Réau and S. Lécolier, *J. Org. Chem.*, **54**, 5208 (1989).
D. Neibecker and R. Réau, *New J. Chem.*, **15**, 279 (1991).
- [4] C.P. Casey and L.M. Petrovich, *J. Am. Chem. Soc.*, **117**, 6007 (1995); G.J.H. Buisman, L.A. van der Veen, P.C.J. Kamer and P.W.N.M. van Leeuwen, *Organometallics*, **16**, 5681 (1997).
- [5] G. Consiglio, S.C.A. Nefkens and A. Borer, *Organometallics*, **10**, 2046 (1991) and references therein.
W.A. Schenk, M. Stubbe and M. Hagel, *J. Organomet. Chem.*, **560**, 257 (1998).
- [6] S. Gladiali, D. Fabbri and L. Kollár, *J. Organomet. Chem.*, **491**, 91 (1995).
- [7] C. Charrier and F. Mathey, *Tetrahedron Lett.*, **28**, 5025 (1987).
- [8] E.H. Bray, I. Caplier and R. Saussez, *Tetrahedron*, **27**, 5523 (1971).
- [9] C. Charrier, N. Maigrot and F. Mathey, *Organometallics*, **6**, 586 (1987).
- [10] J.-J. Brunet, M. Gomez, H. Hajouji and D. Neibecker, *Phosphorus, Sulfur Silicon Relat. Elem.*, **85**, 207 (1993).
- [11] B. Lukas, R.M.G. Roberts, J. Silver and A.S. Wells, *J. Organomet. Chem.*, **256**, 103 (1983).
- [12] A.M. Aguiar, J. Giacini and A. Mills, *J. Org. Chem.*, **27**, 674 (1962).
- [13] C. Charrier, H. Bonnard, F. Mathey and D. Neibecker, *J. Organomet. Chem.*, **231**, 361 (1982).
- [14] F. Mathey, *Chem. Rev.*, **88**, 429 (1988).
- [15] D.D. Perrin, W.L.F. Armarego and D.R. Perrin, "Purification of Laboratory Chemicals", 3rd ed. (Pergamon Press, Oxford, 1988).