

SOLID-SUPPORTED SYNTHESIS OF 5'-mRNA CAP-4 FROM TRYPANOSOMATIDS

Magdalena Lewdorowicz and Jacek Jemielity □ *Department of Biophysics, Institute of Experimental Physics, Warsaw University, Warsaw, Poland*

Ryszard Kierzek □ *Institute of Bioorganic Chemistry, Polish Academy of Sciences, Poznan, Poland*

Michal Shapira □ *Department of Life Sciences, The Ben Gurion University at the Negev, Beer Sheva, Israel*

Janusz Stepinski and Edward Darzynkiewicz □ *Department of Biophysics, Institute of Experimental Physics, Warsaw University, Warsaw, Poland*

□ *The unique structure of 5' mRNA cap from Trypanosomatids is the most modified cap found in nature. Here we present the synthesis of cap-4 ($m^7Gpppm_3^{6,6,2'}Ap_m^{2'}Ap_m^{2'}Cpm_2^{3,2'}Up$) on a disulfide-tethered solid support. This approach allows obtaining cap-4 more efficiently than previously described. Moreover such modified resin could be a useful tool for affinity purification of Leishmania proteins interacting with cap-4. For the final step of synthesis, namely coupling of phosphorylated tetranucleotide with activated 7-methylguanosine 5'-diphosphate two systems were compared. Surprisingly, the coupling in water with Mn^{2+} as a catalyst, gave better results than usually more effective coupling in DMF with $ZnCl_2$.*

Keywords 5' mRNA Cap-4; disulfide linker; *Leishmania*; oligonucleotide synthesis; solid support

INTRODUCTION

Cap structure, that is, m^7GpppN plays important role in various biochemical processes and is used as a tool to study translation, splicing and RNA intercellular transport mechanisms. The recognition of the cap by initiation factor eIF4E is a crucial step in the initiation of protein biosynthesis process. The interactions between cap analogues and eIF4E can be monitored by measuring intrinsic fluorescence quenching of the

This work was supported by grant from Polish Ministry of Science and Higher Education (No. 2 P04A 006 28) and grant from HHMI (No. 55005604).

Address correspondence to Jacek Jemielity, Department of Biophysics, Institute of Experimental Physics, Warsaw University, Zwirki i Wigury 93, Warsaw 02-089, Poland. E-mail: jacekj@biogeo.uw.edu.pl

tryptophan residues.^[1] Since affinities of cap analogues to the eIF4E differ, they are attractive compounds for studying the molecular basis of cap recognition. Moreover, these differences may have also potential therapeutic significance.

The 5' cap structure of *Leishmania*, denoted cap-4, is a complex structure that contains unusual modifications on the first four nucleotides—m⁷Gpppm₃^{6,6,2'}Apm^{2'}Apm^{2'}Cpm₂^{3,2'}U. The two of four nucleosides: N⁶,N⁶,O^{2'}-trimethyladenosine and N³,O^{2'}-dimethyluridine, were previously unknown in nature.^[2] In previous articles, we showed that a synthetically prepared cap-4 analogue was efficiently recognized by an LeishIF4E-1 homologue from *Leishmania*.^[3,4] Moreover, this structure is also efficiently recognized by three of four eIF4E isoforms from *Leishmania*—LeishIF4E-1, LeishIF4E-2, and LeishIF4E-4.

Here we describe the chemical synthesis of cap-4 on a disulfide-tethered solid support (Figure 1). This system can be used for affinity purification of *Leishmania* proteins that interact with cap-4. In order to obtain the cap-4 attached to a resin, we synthesized a disulfide-tethered solid support that would be resistant for ammonolysis.^[5,6] The 5'-phosphorylated tetramer was prepared by the standard phosphoramidite solid phase method. However, since two of the four monomers for solid phase synthesis (phosphoramidites of N⁶,N⁶,O^{2'}-trimethyladenosine and N³,O^{2'}-dimethyluridine) are not available commercially, we synthesized them according to the standard procedures.^[3] As a solid support two commercially available resins were chosen: CPG and TentaGel. The 5'-phosphorylated tetramer was coupled on the resin with an imidazolidine derivative of 7-methylguanosine 5'-diphosphate in the presence of 8-fold excess of ZnCl₂ (in DMF)^[3,7,8] and in the presence of Mn²⁺ (in aqueous solution).^[9]

RESULTS AND DISCUSSION

The disulfide linker allowed the RNA chain assembly without cleavage from the support during ammonolysis of base protecting groups; moreover, such linkers also may be reductively cleaved under mild conditions.^[5,6] The synthesis of disulfide-tethered linker was performed in similar way like described earlier.^[5,6] The amino groups of the standard CPG and TentaGel supports were acylated with dimeric 11-mercaptoundecanoic acid in the presence of N-hydroxysuccinimide, N,N-diisopropylcarbodiimide and 4-(dimethylamino)pyridine. The disulfide bond of dimers was cleaved on solid support by reduction with DTT.

The SH function of resin (**1**) was treated with 2-[(4,4'-dimethoxytrityl)oxy]ethyl 2-pyridyl disulfide and gave modified resin (**2**) ready to use in automated synthesis (Figure 2). According to the dimethoxytrityl (DMTr) cation assay, the loading was 60 μmol/g (for CPG) and 160 μmol/g (for TentaGel).




FIGURE 1 The structure of cap-4 on a disulfide-tethered solid support.

2-[(4,4'-Dimethoxytrityl)oxy]ethyl 2-Pyridyl Disulfide was obtained in a two-step reactions as described earlier involving synthesis of 2-hydroxyethyl 2-pyridyl disulfide from 2,2'-dithiodipyridine and mercaptoethanol and protection of OH-group with 4,4'-dimethoxytrityl chloride.^[5] 5'-O-dimethoxytrityl-N⁶,N⁶,O^{2'}-trimethyladenosine-3'-O-(2-cyanoethyl)-N,N-diisopropylphosphoramidite was prepared as described earlier.^[3] 5'-O-dimethoxytrityl-N³,O^{2'}-dimethyluridine-3'-O-(2-cyanoethyl)-N,N-diisopropylphosphoramidite was obtained from 5'-O-dimethoxytrityl-N³,O^{2'}-dimethyluridine in the presence of *N,N*-diisopropylethylamine and 2-cyanoethoxy *N,N*-diisopropylaminochlorophosphine in anhydrous acetonitrile.

After removal of the DMTr group, the tetranucleotide sequence was assembled by normal phosphoramidite chemistry and the 5'-hydroxy function in tetranucleotide was phosphitylated with 3-DMTrO-2,2-




FIGURE 2 Structure of the supports.

bis(ethoxycarbonyl)propyl 2-cyanoethyl *N,N*-diisopropylphosphoramidite and oxidized to a phosphate triester.^[10] The base and phosphate protections were removed by standard DMTr-off ammonolysis. The HPLC control of the ammonolysis of the protective groups have shown partial loss of tetranucleotide from CPG resin. We finished the ammonolysis when the concentration of deprotected oligonucleotide was about 90% (according to the HPLC analysis). In the case of tetranucleotide on TentGel resin, we haven't observed the cleavage of the product from the resin during ammonolysis. The support-anchored oligonucleotide 5'-phosphate was then treated with 7-methylguanosine 5'-diphosphate imidazolide in DMF in the presence of 8-fold excess of ZnCl_2 (Figure 3) or in aqueous solution in the presence of Mn^{2+} . Surprisingly, the latter method gave better results in coupling reaction.

We monitored the capping reaction using HPLC (the disulfide bond was finally cleaved with DTT and the 2-mercaptoethyl group was allowed to




FIGURE 3 The synthesis of cap-4 on disulfide-tethered solid support.

eliminate in aqueous solution of methanol containing 0.01% triethylamine). On the basis of HPLC the yield of the capping step was 50%, (in the case of TentaGel resin).

The cap-4 on solid support (the structure shown on Figure 1.) will be used in affinity chromatography of eIF4E homologues from *Leishmania*.

REFERENCES

1. Niedzwiecka, A.; Marcotrigiano, J.; Stepinski, J.; Jankowska-Anyszka, M.; Wyslouch-Cieszyńska, A.; Dadlez, M.; Gingras, A.-C.; Mak, P.; Darzynkiewicz, E.; Sonenberg, N. Biophysical studies of eIF4E cap-binding protein: Recognition of mRNA 5' cap structure and synthetic fragments of eIF4G and 4E-BP proteins. *J. Mol. Biol.* **2002**, 319, 615–635.
2. Bangs, J.D.; Crain, P.F.; Hashizume, T.; McCloskey, J.A.; Boothroyd, J.C. Mass spectrometry of mRNA cap 4 from trypanosomatids reveals two novel nucleosides. *J. Biol. Chem.* **1992**, 267, 9805–9815.
3. Lewdorowicz, M.; Yoffe, Y.; Zuberek, J.; Jemielity, J.; Stepinski, J.; Kierzek, R.; Stolarski, R.; Shapira, M.; Darzynkiewicz, E. Chemical synthesis and binding activity of the trypanosomatid cap-4 structure. *RNA* **2004**, 10, 1469–1478.
4. Yoffe, Y.; Zuberek, J.; Lewdorowicz, M.; Zeira, Z.; Keasar, C.; Orr-Dahan, I.; Jankowska-Anyszka, M.; Stepinski, J.; Darzynkiewicz, E.; Shapira, M. Cap-binding activity of an eIF4E homolog from *Leishmania*. *RNA* **2004**, 10, 1764–1775.
5. Salo, H.; Guzaev, A.; Lonnberg, H. Disulfide-tethered solid supports for synthesis of photoluminescent oligonucleotide conjugates: Hydrolytic stability and labeling on the support. *Bioconjugate Chem.* **1998**, 9, 365–371.
6. Jemielity, J.; Heinonen, P.; Lönnberg, H.; Darzynkiewicz, E. A novel approach to solid phase chemical synthesis of oligonucleotide mRNA cap analogs. *Nucleosides, Nucleotides Nucleic Acids* **2005**, 24, 601–605.
7. Jemielity, J.; Fowler, T.; Zuberek, J.; Stepinski, J.; Lewdorowicz, M.; Niedzwiecka, A.; Stolarski, R.; Darzynkiewicz, E.; Rhoads, R.E. Novel “anti-reverse” cap analogs with superior translational properties. *RNA* **2003**, 9, 1108–1122.
8. Stepinski, J.; Waddell, C.; Stolarski, R.; Darzynkiewicz, E.; Rhoads, R.E. Synthesis and properties of mRNAs containing the novel “anti-reverse” cap analogues 7-methyl(3'-O-methyl)GpppG and 7-methyl (3'-deoxy)GpppG. *RNA* **2001**, 7, 1486–1495.
9. Sawai, H.; Wakai, H.; Nakamura-Ozaki, A. Synthesis and reactions of nucleoside 5'-diphosphate imidazolide. A nonenzymatic capping agent for 5'-monophosphorylated oligoribonucleotides in aqueous solution. *J. Org. Chem.* **1999**, 64, 5836–5840.
10. Guzaev, A.; Salo, H.; Azhayev, A.; Lonnberg, H. A new approach for chemical phosphorylation of oligonucleotides at the 5'-terminus. *Tetrahedron* **1995**, 51, 9375–9384.

Copyright of Nucleosides, Nucleotides & Nucleic Acids is the property of Taylor & Francis Ltd and its content may not be copied or emailed to multiple sites or posted to a listserv without the copyright holder's express written permission. However, users may print, download, or email articles for individual use.