

Published on Web 07/18/2008

## Observation of a Hidden Intermediate in the Stille Reaction. Study of the Reversal of the Transmetalation Step

Mónica H. Pérez-Temprano,<sup>†</sup> Ainara Nova,<sup>‡</sup> Juan A. Casares,<sup>\*,†</sup> and Pablo Espinet<sup>\*,†</sup>

IU CINQUIMA/Química Inorgánica, Facultad de Ciencias, Universidad de Valladolid, E-47071 Valladolid, Spain, and Química Física, Edifici C.n. Universitat Autònoma de Barcelona, E-08193 Bellaterra, Catalonia, Spain

Received May 5, 2008; E-mail: espinet@qi.uva.es; casares@qi.uva.es

Among the palladium-catalyzed cross-coupling processes, the Migita–Stille reaction is perhaps the most deeply studied experimentally. The accepted mechanism can follow the open or cyclic pathways shown in Scheme 1. A comprehensive account of the mechanistic studies was published in 2004.<sup>1</sup> Further mechanistic studies,<sup>2–4</sup> including detailed theoretical (DFT) calculations,<sup>5,6</sup> have been published recently.

The steps preceding the transmetalation have been experimentally confirmed by observation of intermediates. Of the two intermediates produced by transmetalation in an *open* mechanism (lower part of Scheme 1), *trans*-[PdR<sup>1</sup>R<sup>2</sup>L<sub>2</sub>] (for which the reductive elimination is rate-determining) was observed by NMR, whereas *cis*-[PdR<sup>1</sup>R<sup>2</sup>L<sub>2</sub>] (for which the reductive elimination) was unambiguously deduced kinetically.<sup>7,8</sup> However, for the *cyclic* mechanism (upper part of Scheme 1), the transmetalation is usually rate-determining<sup>9</sup> since the cis arrangement of the R<sup>1</sup> and R<sup>2</sup> substituents obtained from the transmetalation makes their coupling fast. This has precluded so far the observation of any putative transmetalated intermediate that would further support experimentally the proposed *cyclic* mechanism.

Our recent DFT calculations for the catalyzed coupling of PhBr and Sn(vinyl)Me<sub>3</sub> suggest that the formation of the intermediate *cis*-[PdR<sup>1</sup>R<sup>2</sup>L<sub>2</sub>] from the starting reagents is thermodynamically disfavored, so a successful coupling requires an easy irreversible  $R^1-R^2$  reductive elimination to pull the reaction forward.<sup>5</sup> Moreover, a post-transmetalation intermediate with ISnBu<sub>3</sub> acting as a I donor ligand, [PdR<sup>1</sup>R<sup>2</sup>(AsPh<sub>3</sub>)(ISnBu<sub>3</sub>)] (**I1**), was predicted. The detection of this intermediate would provide strong experimental support to the cyclic mechanism since this intermediate does not appear in the open mechanism.

A reaction of XSnBu<sub>3</sub> with a *cis*-[PdR<sup>1</sup>R<sup>2</sup>L<sub>2</sub>] complex reluctant to reductively eliminate R<sup>1</sup>-R<sup>2</sup> might behave (obeying the microscopic reversibility principle) as the reverse of the Stille reaction, to give *trans*-[PdR<sup>1</sup>XL<sub>2</sub>] and R<sup>2</sup>SnBu<sub>3</sub>. This reverse reaction should allow for study of the hidden post-transmetalation steps in the cyclic mechanism; since in the retro-transmetalation<sup>10</sup> sense the intermediate **I1** precedes the transmetalation step, which is the highest barrier for the process, it might become observable. *cis*-[PdRf<sub>2</sub>(AsPh<sub>3</sub>)<sub>2</sub>] (R<sup>1</sup> = R<sup>2</sup> = Rf = 3,5-C<sub>6</sub>Cl<sub>2</sub>F<sub>3</sub>), **1**,<sup>11</sup> a stable compound that additionally offers easy monitoring of the reactions by <sup>19</sup>F NMR,<sup>12</sup> was chosen to study the retro-transmetalation of the Stille reaction (Scheme 2).

The reaction of **1** and ISnBu<sub>3</sub> (**2**) in THF at 323 K yielded a mixture of *trans*-[PdRfI(AsPh<sub>3</sub>)<sub>2</sub>] (**3**) and RfSnBu<sub>3</sub> (**4**) (Scheme 2). Minor side reactions (some decomposition to Pd(0) or formation of some *trans*-[PdRf<sub>2</sub>(AsPh<sub>3</sub>)<sub>2</sub>], which does not undergo retro-

## Scheme 1



Scheme 2



transmetalation) were irrelevant for the kinetic study of the retrotransmetalation process, as they were observed only at times larger than those used in the kinetic study. The other species detected by <sup>19</sup>F NMR were the pursued intermediate [PdRf<sub>2</sub>(AsPh<sub>3</sub>)(ISnBu<sub>3</sub>)] (**I1**) and the dimer [Pd<sub>2</sub>Rf<sub>2</sub>( $\mu$ -I)<sub>2</sub>(AsPh<sub>3</sub>)<sub>2</sub>] (**5**), already reported.<sup>13</sup> In an independent study, the fast equilibrium **3** + **4** = <sup>1</sup>/<sub>2</sub> **5** + L was fully characterized ( $K_4(323 \text{ K}) = 3.7 \times 10^{-5} \text{ mol L}^{-1}$ ;  $\Delta H^0$ = 19.4 ± 2.0 kcal mol<sup>-1</sup>;  $\Delta S^0 = 39.7 \pm 6.2$  cal K<sup>-1</sup> mol<sup>-1</sup>).<sup>14</sup> These values are important for the retro-transmetalation kinetics since the AsPh<sub>3</sub> released modifies the rate of transmetalation. Moreover, due to the high  $\Delta S^0$  value, **5** is not observed in the NMR spectra at room or lower temperatures. The relevance of these

<sup>&</sup>lt;sup>†</sup> Universidad de Valladolid.

<sup>&</sup>lt;sup>\*</sup> Universitat Autònoma de Barcelona.



*Figure 1.* <sup>19</sup>F NMR spectrum ( $F_{ortho}$  range) of the reaction in Scheme 2, recorded at 230 K, showing the signals of intermediate **I1** and a schematic representation of the through-space  ${}^{19}F^{-19}F$  coupling (dotted curves in the structure). The signal \* belongs to *trans*-[PdRf<sub>2</sub>(AsPh<sub>3</sub>)<sub>2</sub>]. <sup>13</sup>C satellites are observed for the very intense signals of *cis*-[PdRf<sub>2</sub>(AsPh<sub>3</sub>)<sub>2</sub>] and *trans*-[PdRf<sub>1</sub>(AsPh<sub>3</sub>)<sub>2</sub>].

Scheme 3



dimeric species in the Stille reaction, usually carried out at reflux, has been noticed before.<sup>13,15–17</sup>

The predicted intermediate  $[PdRf_2(AsPh_3)(ISnBu_3)]$  (**11**) could be unambiguously detected in the course of the kinetic experiments below 283 K, as a deceptively simple pair of doublets of an AA'BB' spin system, due to the through-space *J* coupling of *ortho*-fluorine atoms of the nonequivalent perhaloaryl groups in conditions of slow rotation of the fluoroaryl group about the C–Pd bond (Figure 1). Irradiation of one of the signals simplifies the other to a sharp singlet, while at 323 K the signals appear as broad triplets. This spectroscopic and dynamic behavior has been reported in depth for other  $[PdRf_2L^1L^2]$  complexes.<sup>12,18,19</sup>

The identity of **I1** was confirmed independently. Attempts to prepare [PdRf<sub>2</sub>(AsPh<sub>3</sub>)(THF)] (Scheme 3) as a possible precursor for **I1** afforded a complex mixture of **1** and **6**–**9** in THF,<sup>18,20</sup> which were easily identified. The addition to this mixture of a large excess of ISnBu<sub>3</sub> gave, as major species, *cis*-[PdRf<sub>2</sub>(AsPh<sub>3</sub>)<sub>2</sub>] (**1**), *cis*-[PdRf<sub>2</sub>(ISnBu<sub>3</sub>)<sub>2</sub>] (**10**), and [PdRf<sub>2</sub>(AsPh<sub>3</sub>)(ISnBu<sub>3</sub>)] (**11**). In addition to the <sup>19</sup>F NMR spectra, the <sup>119</sup>Sn NMR spectrum confirmed three Sn-containing species, appearing at 63.2 ppm (m) (the most intense, ISnBu<sub>3</sub>, in excess in solution), 39.7 (**I1**), and 104.6 ppm (**10**).

The activation parameters for the formation of **I1** were determined by an Eyring study using <sup>19</sup>F NMR (at 376.50 MHz) in the range of 225–270 K. Fortunately, at these temperatures, the slow transmetalation step is not observed within the time of experiment. The study afforded for **TS1** the following values:  $\Delta H^{\ddagger} = 15.0 \pm$ 1.1 kcal mol<sup>-1</sup>;  $\Delta S^{\ddagger} = -17.4 \pm 4.5$  cal K<sup>-1</sup> mol<sup>-1</sup>. The negative value for  $\Delta S^{\ddagger}$  supports an associative substitution of AsPh<sub>3</sub> by



**Figure 2.** Observed concentrations as a function of time of  $(\bullet)$  the palladium reactant 1;  $(\blacksquare)$  the trans product 3 (corrected for the formation of dimer 4). Solid lines are the best fit using Gepasi.

ISnBu<sub>3</sub>. These values were used to calculate  $\Delta G^{\ddagger}$  for the preequilibrium at other temperatures.

In an independent study,  $\Delta G^{\dagger}$  for the transmetalation (TS2) was determined at 323 K by monitoring the variations of concentrations of 1 and 3 during the retro-transmetalation reaction. At this temperature, the concentration of I1 reached a maximum very quickly and then kept steady until the final steps of the reaction. The data of eleven runs were fit to the multistep kinetic model proposed in Scheme 2, by nonlinear least-squares (NLLS) regression, using the program Gepasi.<sup>21</sup> The experimental values for the equilibrium constants  $K_1$  ( $K_1 = k_1/k_{-1}$ ; see Scheme 1) and  $K_4$  were entered into the adjust program. Figure 2 shows the results of a typical run (consumption of reactant and formation of trans product as a function of time) and the curves of the Gepasi fit. Luckily, it was found that the value of  $k_2$  was little sensitive to the value assumed for  $k_3$  (which could not be determined experimentally). Thus, although the Gibbs energy of TS3 could not be assigned, a reliable value of  $\Delta G^{\ddagger}(323 \text{ K}) = 26.5 \text{ kcal/mol} (k_2 = 5.28 \times 10^{-3} \text{ kcal/mol})$  $s^{-1}$ ) was calculated for **TS2**.

DFT calculations were carried out on a model where AsH<sub>3</sub> and ISnMe<sub>3</sub> are used instead of AsPh<sub>3</sub> and ISnBu<sub>3</sub> (the computational details are given in the Supporting Information). The calculated  $\Delta E$ (gas) and  $\Delta E$ (solvent, THF) values for the whole reaction are depicted in Figure 3, along with the experimental  $\Delta G$  values determined in THF, so that calculated and experimental results can be directly compared. Note that the experimental data only reach to **TS2** since, beyond that point, only the formation of **3** + **4** is experimentally observed, along with the formation of dimer **5**. In agreement with the calculated  $\Delta E^{0}$  values of 7–8 kcal/mol for the retro-transmetalation equilibrium **1** + **2** = **3** + **4** (including in **3** the amount corresponding to its dimerization to **5**), the retro-transmetalation reaction is completely shifted to the right, that is, to the formation of the Stille reagents, as observed by <sup>19</sup>F NMR.

For intermediate **I1** and for **TS2** (Figure 4), the total number of particles does not change with respect to the starting system (in each case, taking into account the molecule of ligand released, the exchange is two particles for two particles), and a moderate entropic contribution is expected. Consistent with this prediction, the respective experimental values of their Gibbs energies lie between those calculated for E in the gas phase and in solvent. This



**Figure 3.** Profile of the retro-transmetalation reaction 1 + 2, showing the experimental values in THF and the calculated values  $\Delta E(gas)$  and  $\Delta E$ (solvent) in kcal/mol. The calculated structure of the detected intermediate I1 is also shown.



Figure 4. Calculated geometry (DFT) for the observed intermediate I1, and for the transmetalation transition state TS2. Other calculated geometries are given in the Supporting Information.

coincidence supports the idea that the calculated values give a reasonable estimation of the system as far as entropy changes are small.

In contrast, the *E* values calculated for **TS1** are noticeably lower than the experimental Gibbs energy value, but the calculated E<sub>TS1</sub>(solvent, THF) value (12.5 kcal/mol) is not far from the experimental  $\Delta H^{\dagger}$  value (15.0  $\pm$  1.1 kcal/mol). As discussed above, the experimental data point to an associative transition state (hence with a considerable entropic contribution). Interestingly, a correction for entropy of about 8-10 kcal/mol to the  $E_{TSI}$ (solvent, THF) value would bring the calculation very close to the  $\Delta G^{\dagger}_{TS1}$  value obtained experimentally (20.7 kcal/mol).<sup>22</sup> Thus, in spite of some deviations, arising from the simplification of the system using a calculation model with smaller ligands, the estimation of E(solvent) and the crude correction for entropy (based only in the number of particles) behave remarkably well and should possibly be reliable enough to assign associative versus dissociative processes in other cases where only  $\Delta G^{\dagger}$  values were experimentally available.

In summary, an intermediate of the type cis-[PdR<sup>1</sup>R<sup>2</sup>L(ISnBu<sub>3</sub>)], which was proposed to be the result of a cyclic transmetalation but had never been detected in the direct reaction, has been spectroscopically observed in a retro-transmetalation reaction. The formation of this intermediate is a strong experimental support for the occurrence of cyclic mechanisms in some Migita-Stille reactions.

Acknowledgment. We thank the Ministerio de Ciencia y Tecnología (Project CTQ2007-67411/BQU, CTQ2005-09000-C02-01, and a studentship to M.H.P-T.), INTECAT Consolider Ingenio 2010 (CSD2006-0003 and CSD2007-0006), and the Junta de Castilla y León (VA-060-04) for support.

Supporting Information Available: Details of syntheses, characterization of the complexes, kinetic experiments, and DFT calculations including calculated structures. This material is available free of charge via the Internet at http://pubs.acs.org.

## References

- (1) Espinet, P.; Echavarren, A. M. Angew. Chem., Int. Ed. 2004, 43, 4704-
- (2) Nilsson, P.; Puxty, G.; Wendt, O. F. Organometallics 2006, 27, 1285-1292
- (3)Ariafard, A.; Lin, Z.; Fairlamb, J. S. Organometallics 2006, 27, 5788-5794
- Santos, L. S.; Rosso, G. B.; Pilli, R. A.; Eberlin, M. N. J. Org. Chem. 2007, 72, 5809–5812. (4)Nova, A.; Ujaque, G.; Maseras, F.; Lledos, A.; Espinet, P. J. Am. Chem.
- (5)Soc. 2006, 128, 14571-14588. (6)
- (a) Álvarez, R.; Faza, O. N.; Lopez, C. S.; de Lera, A. R. Org. Lett. 2006, 8, 35-38. (b) Alvarez, R.; Faza, O. N.; de Lera, A. R.; Cardenas, D. J. Adv. Synth. Catal. 2007, 349, 887-906.
- Casado, A. L.; Espinet, P.; Gallego, A. M.; Martínez-Ilarduya, J. M. Chem. Commun. 2001, 339–340. (7)
- Casado, A. L.; Espinet, P.; Gallego, A. M J. Am. Chem. Soc. 2000, 122, (8)11771-11782
- Casado, A. L.; Espinet, P. J. Am. Chem. Soc. 1998, 120, 8978-8985.
- (10) We use the word "retrotransmetalation" to indicate the exchange of an R on Pd for an X on Sn, the reverse of what happens in the so-called transmetalation.
- (11) Espinet, P.; Martínez-Ilarduya, J. M.; Pérez-Briso, C.; Casado, A.; Alonso, M. A. J. Organomet. Chem. **1998**, 551, 9–20. (12) Espinet, P.; Albéniz, A. C.; Casares, J. A.; Martínez-Ilarduya, J. M. Coord.
- Chem. Rev., in press.
- (13) Casares, J. A.; Espinet, P.; Salas, G. *Chem.—Eur. J.* 2002, *8*, 4843–4853.
  (14) From the integrated <sup>19</sup>F spectra recorded between 302.6 and 329.9 K.
- (15) (a) Mann, G.; Shelby, Q.; Roy, A. H.; Hartwig, J. F. Organometallics 2003, 22, 2775–2789. (b) Roy, A. H.; Hartwig, J. F. Organometallics 2004, 23, 1533-1541.
- (16) Ricci, A.; Angeluci, F.; Bassetti, M.; Lo Sterzo, C. J. Am. Chem. Soc. 2002, 124, 1060–1071.
- (17) Amatore, C.; Bahsoun, A. A.; Jutand, A.; Meyer, G.; Ntpede, A. N.; Ricard, L. J. Am. Chem. Soc. 2003, 125, 4212-422
- (18) Casares, J. A.; Espinet, P.; Martínez-Ilarduya, J. M.; Mucientes, J. J.; Salas, G. Inorg. Chem. 2007, 46, 1027-1032.
- (19) (a) Alonso, M. A.; Casares, J. A.; Espinet, P.; Martínez-Ilarduya, J. M.; Pérez-Briso, C. Eur. J. Inorg. Chem. **1998**, 1745–1753. (b) de la Cruz, R.; Espinet, P.; Gallego, A. M.; Martín-Alvarez, J. M.; Martínez-Ilarduya, J. M. J. Organomet. Chem. **2002**, 663, 108–117.
- (20) Similar complexes cis-[PdRf2(solvent)(H2O)] and cis-[PdRf2(H2O)2], as well as a detailed study of equilibria in acetone solution for analogous Ni cis-[NiRf2(H2O)2], complexes cis-[NiRf2(solvent)(H2O)], [NiRf<sub>2</sub>(solvent)L], and cis-[NiRf<sub>2</sub>L(H<sub>2</sub>O)], (L = monodentate ligands,
- (21) (a) Mendes, P. Comput. Appl. Biosci. 1993, 9, 563–571. (b) Mendes, P. Trends Biochem. Sci. 1997, 22, 361–363. (c) Mendes, P.; Kell, D. B. Bioinformatics 1998, 14, 869–883. (d) Martins, A. M.; Mendes, P.; Cordeiro, C.; Freire, A. P. Eur. J. Biochem. 2001, 268, 3930–3936.
- (22) The estimated entropic contribution for only the variation in the number of particles (considered as rigid spheres) is 8-10 kcal/mol; see: Watson, L. A.; Eisenstein, O. J. Chem. Educ. 2002, 79, 1269-1277.

JA802994V