

25-Hydroxycholecalciferol: a Biologically Active Metabolite of Cholecalciferol


By J. W. BLUNT, H. F. DeLUCA,* and H. K. SCHNOES

(Department of Biochemistry, University of Wisconsin, Madison, Wisconsin 53706)

It has long been suspected that cholecalciferol (I) (vitamin D₃) must first be converted into a biologically active metabolite before it could exert its metabolic effects.¹ However, it was not until 1966 that the existence of biologically active metabolites was unequivocally shown in this laboratory.² From these, a major polar metabolite fraction designed as "Peak IV" from its chromatographic behaviour appeared to meet the criteria of the metabolically active form of the vitamin.¹⁻⁴ It has now been possible to isolate the biologically active component of this metabolite fraction from the blood plasma of hogs given 250,000 i.u. cholecalciferol daily for 3½ weeks. This substance is identified as 25-hydroxycholecalciferol (II).

The protein precipitated from 6.8 l. of plasma from 4 hogs was extracted with chloroform-methanol and its crude extract submitted to absorption chromatography followed by partition chromatography, yielding 1.3 mg. of pure (II). This was identified as follows. Its u.v. spectrum displayed a maximum at 265 mμ as for (I). G.l.c. at 240° gave two peaks, representing 25-hydroxy-pyrocholecalciferol and its isopyro-isomer, both of which had u.v. spectra identical to those for the corresponding pyro- and isopyro-cholecalciferols arising from (I) under similar conditions. Thus the similarity in structure between (I) and (II) was established. The mass spectrum of (II) indicated a M.W. of 400, (C₂₇H₄₄O₂), i.e. a cholecalciferdiol. Further, the mass spectrum was very similar to that of (I), both showing a peak at *m/e* 271

(C₁₉H₂₇O) arising from the removal of the side-chain. Hence the extra hydroxyl group in (II) is located on the side-chain. The presence of a peak at *m/e* 59 (C₃H₇O) in the spectrum of (II) but not in the spectrum of (I) suggested the location of the hydroxyl function at C-25. This was confirmed by the 100 Mc./sec. n.m.r. spectrum of (II), which



was identical with the spectrum of (I) except that the doublet at 0.87 p.p.m. (*J* 6.0 c./sec.) due to the secondary C-26,27 methyl groups in (I) was absent and was replaced by a strong singlet peak at 1.20 p.p.m., identical with the peak due to the C-26,27 methyl groups in 25-hydroxycholesterol.

We thank the National Institutes of Health for financial support, Mrs. Yoko Tanaka for her excellent technical assistance, and Miss Martha Petrie for recording the n.m.r. spectra.

(Received, May 13th, 1968; Com. 607.)

¹ H. F. DeLuca, *Vitamins and Hormones*, 1967, **25**, 315.

² J. Lund and H. F. DeLuca, *J. Lipid Res.*, 1966, **7**, 739.

³ H. Morii, J. Lund, P. Neville, and H. F. DeLuca, *Arch. Biochem. Biophys.*, 1967, **120**, 503.

⁴ S. J. Stohs and H. F. DeLuca, *Biochemistry*, 1967, **6**, 3338.