1390

Enantioselective Vinylation of Aldehydes with Chiral β-Silylphosphorus Ylides

Hideo lio,* Akio Fujii, Masahiro Ishii and Takashi Tokoroyama*

Faculty of Science, Osaka City University, Sumiyoshi-ku, Osaka 558, Japan

Both enantiomers of diastereoisomeric [1-(N,N-dimethylaminoethyl)ferrocenyl]diphenyl-(2-diphenylmethylsilylethylidene)phosphorane **3** and **4** are prepared; their reaction with benzaldehyde and aliphatic aldehydes is found to give vinylation products with variable enantiomeric excess (up to 92%).

The development of synthetic methods for obtaining enantiomerically pure compounds from carbonyl compounds is an important theme in current organic chemistry, in which various sophisticated methodologies have been introduced.¹ In connection with our work on the diastereoselective synthesis of allyl alcohol using β -silylphosphorus ylide,² we have explored the ylide reagents effecting asymmetric induction, which is the subject of this paper.

Our previous studies indicated that electron donating aryl substituents on the phosphorus atom of the ylide is expedient for the selective formation of allyl alcohol over the Wittig alkenation reaction.³ We investigated the (dimethylamino)ethylferrocenyl group as a chiral substituent because of its through space electronically donating character and its sterically congesting nature.⁴

Enantiomerically pure ferrocenylphosphonium salts 1 and 2 {(S)(R)-1, m.p. 207 °C (decomp.), $[\alpha]_D^{21}$ +44.9° (*c* 1.18, CHCl₃); (*R*)(*R*)-2, m.p. 135 °C (decomp.), $[\alpha]_D^{21}$ +6.6° (*c* 1.44, CHCl₃)} and their enantiomers are synthesised in two steps from the corresponding optically pure diastereoisomeric *N*,*N*-dimethyl-1-[2-(diphenylphosphino)ferrocenyl]ethyl-

amines (PPFA)⁵ in 54 and 49% yields $[i, MeI-Et_2O, 0 \ ^{\circ}C$ to room temp.; *ii*, butyllithium-tetrahydrofuran (THF), 0 $^{\circ}C$ then ICH₂SiMePh₂, 0 $^{\circ}C$ to room temp.]. The chiral ylides **3**

and 4 were generated *in situ* from 1 and 2, respectively by butyllithium in THF and the reactions with aldehydes were carried out in THF at -78 °C. The allylic silyl ether 5 was obtained exclusively without formation of the Wittig alkena-



Table 1 Reaction of chiral ylides with aldehydes

Entry	Aldehyde	Ylide	Lewis acid	E.e. (%) ^a	Config.	Yield (%)
 1	PhCHO	(R)(S)-3		77	R	68
2		(R)(S)-3	Et ₂ AlOEt	70	R	83
3		(S)(R)-3	Et ₂ AlOEt	70	S	80
4		(S)(S)-4		47	R	24
5		(S)(S)-4	Et ₂ AlOEt	57	R	62
6	PhCH ₂ CH ₂ CHO	(R)(S)-3		9	S	34
7	2 2	(S)(R)-3	Et ₂ AlOEt ^b	25	R	95
8		$(\hat{R})(\hat{R})$ -4	_	85	R	46
9		(S)(S)-4	_	80	S	44
10		(S)(S)-4	Et ₂ AlOEt	70	S	51
11	c-C ₆ H ₁₁ CHO	(S)(R)-3		13	R	27
12	0 11	(S)(R)-3	Et ₂ AlOEt	27	R	94
13		(R)(R)-4		92	R	43

^{*a*} E.e. was determined by ¹H NMR (400 MHz) spectrum of the corresponding [methoxy(trifluoromethyl)phenylacetyl] (MTPA) ester derived in two steps (*i*, Bu₄NF–THF; *ii*, (+)- or (-)-MTPACl–C₅H₅N–CH₂Cl₂). ^{*b*} Lewis acids other than Et₂AlOEt gave similar results (AlEt₃, 27% e.e., 95% yield; AlBuⁱ₃, 23% e.e., 92% yield; Ti(OPrⁱ)₄, 27% e.e., 75% yield; BF₃–Et₂O, 22% e.e., 62% yield.



Fig. 1 X-Ray structure of (S)(R)-1

tion product and the recovered chiral phosphines PPFA could be recycled for the preparation of 1 and 2 (Scheme 1). The results of the reaction are shown in Table 1.

The ylide 3 gave 5 with 77% enantiomeric excess (e.e.) in the reaction with benzaldehyde. Although 3 yielded 5 with low e.e. in the reaction with the aliphatic aldehyde, high asymmetric induction up to 92% e.e. was attained in the reaction with ylide 4 and cyclohexylcarbaldehyde. The addition of diethylaluminium ethoxide did not affect the enantiometric excess but increased the yield of 5. Both (R)(S)-3 and (S)(S)-4 gave the *R* enantiomer in the reaction with benzaldehyde, which means the chirality of 5 is mainly induced by the planar chirality of the ferrocenyl group rather than by the central chirality of the aminoethyl group.⁶

For reference purposes an X-ray single crystallographic analysis of (S)(R)-1 was carried out and the resulting structure

is reproduced in Fig. 1.† If the ylide (S)(R)-3 has a configuration similar to (S)(R)-1 in which C(40)-C(41) is

[†] Crystal data for (S)(R)-1: C₄₁H₄₅NSiPFel, M = 793.59, tetragonal, space group P4₁, a = b = 9.755(4), c = 39.96(1) Å, U = 3803(3) Å³, Z = 4, $D_c = 1.387$ g cm⁻³, $\mu = 12.99$ cm⁻¹. 3414 reflections with 2θ ≤ 50.0° were recorded on a four-circle diffractometer using graphite-monochromated Mo-Kα radiation. Of these 2823 [with $I > 3\sigma(I)$] were judged observed. The structure was solved using Patterson methods. All hydrogen atoms could be located on a difference Fourier synthesis. Full matrix least-squares refinement with anisotropic temperature factors for non-hydrogen atoms and isotropic hydrogens converged to R = 0.035 and $R_w = 0.052$. The maximum and minimum residual electron densities in the final difference Fourier map were 0.82 and −1.25 e Å⁻³. Atomic coordinates, thermal parameters, and bond lengths and angles have been deposited at the Cambridge Crystallographic Data Centre. See Notice to Authors, Issue No. 1.



Fig. 2 Transition state for reaction of ylide (S)(R)-3 with benzaldehyde

oriented antiperiplanar to the ferrocenyl group and the silyl group is tilted towards the *re* face of the planar sp² carbon C(40) [dihedral angles of C(1)–P–C(40)–C(41) and P–C(40)–C(41)–Si are 179.2 and 154.4° in (S)(R)-1, respectively],⁷ then the *Si* face of benzaldehyde would approach the *Re* face of the ylide (S)(R)-3 C(40) to avoid the non-bonding interaction between the phenyl group of the former and the substituent on the phosphorus atom in the latter.⁸ The oxygen atom of benzaldehyde would interact with the tilted silicon atom to form a five-membered pentacoordinate silicate (see Fig. 2), which gives the product with *S* configuration. To the best of our knowledge, this is the first example of central chirality induction using a chiral phosphorus ylide.

We thank Professor K. Hirotsu and Miss I. Miyahara, Faculty of Science, Osaka City University for the X-ray crystallographic analysis. This research was supported in part by a Grant-in-Aid for Scientific Research (No. 63540430) from the Ministry of Education, Science, and Culture of Japan.

Received, 2nd July 1991; Com. 1/03311E

References

- R. Noyori and M. Kitamura, *Angew. Chem., Int. Ed. Engl.*, 1991, 30, 49; M. Yoshioka, T. Kawakita and M. Ohno, *Tetrahedron Lett.*, 1989, 30, 1657; W. Oppolzer and R. N. Nadinov, *Tetrahedron Lett.*, 1988, 29, 5645.
- 2 M. Tsukamoto, H. Iio and T. Tokoroyama, *Tetrahedron Lett.*, 1987, 28, 4561.
- 3 M. Tsukamoto, H. Iio and T. Tokoroyama, *Tetrahedron Lett.*, 1985, **26**, 4471; M. Tsukamoto, H. Iio and T. Tokoroyama, *J. Chem. Soc.*, *Chem. Commun.*, 1986, 880.
- 4 W. E. McEwen, C. E. Sullivan and R. O. Day, *Organometallics*, 1983, **2**, 420; D. W. Allen, *Z. Naturforsch. B*, 1980, **35**, 981.
- 5 T. Hayashi, T. Mise, M. Fukushima, M. Kagotani, N. Ngashima, Y. Hamada, A. Matsumoto, S. Kawakami, M. Konishi, K. Yamamoto and M. Kumada, *Bull. Chem. Soc. Jpn.*, 1980, **53**, 1138.
- 6 A. Togni and S. T. Pastor, J. Org. Chem., 1990, 55, 1649; S. T. Pastor and A. Togni, J. Am. Chem. Soc., 1989, 111, 2333.
- 7 J. C. J. Bart, J. Chem. Soc. (B), 1969, 350.
- 8 M. Schlosser and B. J. Schaub, J. Am. Chem. Soc., 1982, 104, 5821;
 E. Vedejs and C. F. Marth, J. Am. Chem. Soc., 1988, 110, 3948.