

Cycloadditions of *N*-Sulfonyl Nitrones Generated by Lewis Acid Catalyzed Rearrangement of Oxaziridines

Katherine M. Partridge, Mary E. Anzovino, and Tehshik P. Yoon\*

Department of Chemistry, University of Wisconsin–Madison, 1101 University Avenue, Madison, Wisconsin 53706

Received December 21, 2007; E-mail: tyoon@chem.wisc.edu


The weak N–O bond of 1,2-isoxazolidines can undergo facile reductive cleavage to furnish 1,3-aminoalcohols. Due to the ease with which isoxazolidines can be processed into these synthetically valuable structures, considerable effort has been invested toward the development of methods for [3 + 2] nitronc cycloadditions.<sup>1</sup> This enormous body of work, however, has focused primarily upon *N*-alkyl nitrones that produce 1,2-isoxazolidines whose *N*-substituents cannot be removed without concomitant cleavage of the sensitive N–O bond;<sup>2</sup> cycloadditions of nitrones bearing easily hydrolyzed electron-withdrawing *N*-substituents are almost completely unexplored.<sup>3</sup> Recently, 1,2-isoxazolidines have begun to attract significant attention as nucleoside analogues<sup>4</sup> and synthetic transcriptase activators.<sup>5</sup> Exploration of the biological activity of this class of heterocycles would be facilitated by new methods to access the parent *N*-unsubstituted isoxazolidines for diversification into libraries of *N*-modified analogues. Herein, we report a novel method for Lewis acid catalyzed formation and cycloaddition of otherwise inaccessible *N*-nosyl nitrones, which produce 1,2-isoxazolidines that can be deprotected under mild conditions without accompanying ring cleavage.

Our group has been developing new synthetic methods based upon activation of oxaziridines with exogenous catalysts. We recently reported a new aminohydroxylation procedure in which *N*-sulfonyl oxaziridine **1** reacts with styrenes in the presence of Cu(TFA)<sub>2</sub> to provide 1,3-oxazolidines.<sup>6</sup> In an unusual example of divergent, catalyst-controlled reactivity, we have discovered that the isomeric 1,2-isoxazolidine **3** is produced when styrene reacts with **1** in the presence of Sc(OTf)<sub>3</sub> (Scheme 1). This heterocycle presumably arises from rearrangement of the oxaziridine to a transient *N*-sulfonyl nitronc (**2**) that undergoes [3 + 2] cycloaddition with styrene. While the corresponding rearrangement of *N*-alkyl oxaziridines to nitrones is precedented,<sup>7</sup> we are unaware of any previous studies describing the Lewis acid catalyzed rearrangement of *N*-sulfonyl oxaziridines to nitrones.<sup>8</sup>

Careful analysis of the Sc(OTf)<sub>3</sub>-catalyzed reaction revealed that benzaldehyde oxime, *O*-sulfonyl oxime **4**, and polystyrene are significant byproducts. We speculated that these compounds could arise upon heterolytic decomposition of the *N*-sulfonyl nitronc (Scheme 1);<sup>9</sup> the sulfonyl cation could either recombine with the oxime anion to produce **4** or initiate the cationic polymerization of styrene. This mechanistic hypothesis suggests that the efficiency of the desired cycloaddition would be improved by decreasing the propensity of the nitronc to undergo ionic fragmentation. Indeed, optimized reaction conditions for this novel transformation utilized toluene as a nonpolar solvent, which disfavors formation of ionic intermediates, as well as oxaziridines bearing *N*-4-nitrobenzenesulfonyl groups that would further destabilize the problematic sulfonyl cation (*vide infra*).

The effect of various Lewis acids on the cycloadditions of *N*-nosyl oxaziridine **5** in toluene is summarized in Table 1. A wide range of oxophilic Lewis acids afford the desired 1,2-isoxazolidine. The diastereoselectivity of the cycloaddition shows a dependence

## Scheme 1



**Table 1.** Effect of Oxophilic Lewis Acids on the Efficiency and Diastereoselectivity of 1,2-Isoxazolidine Formation<sup>a</sup>

c1ccccc1C=C + NsN1C(=O)C1c2ccccc2
 $\xrightarrow[\text{5 \AA mol sieves, toluene, 23 } ^\circ\text{C}]{\text{Lewis acid}}$ 
c1ccccc1[C@H]2C[C@@H](c3ccccc3)O2N1C(=O)C1

**5** **6**

entry	catalyst	loading mol %	time h	yield <sup>b</sup>	d.r. <sup>c</sup> ( <i>syn/anti</i> )
1	Sc(OTf) <sub>3</sub>	20	6	23%	1.2:1
2	BF <sub>3</sub> ·OEt <sub>2</sub>	20	6	58%	> 10:1
3	SnCl <sub>4</sub>	20	6	80%	6:1
4	TiCl <sub>4</sub>	20	1	93%	> 10:1
5	TiCl <sub>4</sub>	10	4	95%	> 10:1

**Table 2.** Investigation of Olefin Scope in Nitron Cycloadditions

$\text{Ar-CH=CH}_2 + \text{Ns-N-O} \xrightarrow[\text{23 } ^\circ\text{C}]{\text{TiCl}_4, \text{5 \AA mol sieves}} \text{Ph-CH(O-Ns)-CH}_2\text{-Ar}$					
entry <sup>a</sup>	product	mol% TiCl <sub>4</sub> yield, <sup>b</sup> d.r. <sup>c</sup>	entry <sup>a</sup>	product	mol% TiCl <sub>4</sub> yield, <sup>b</sup> d.r. <sup>c</sup>
1		10 mol% 95% yield >10:1 d.r.	7 <sup>e</sup>		10 mol% 87% yield >10:1 d.r.
2		10 mol% 87% yield >10:1 d.r.	8 <sup>e</sup>		10 mol% 65% yield >10:1 d.r.
3 <sup>d</sup>		100 mol% 93% yield >10:1 d.r.	9 <sup>e</sup>		10 mol% 83% yield 3:1 d.r.
4 <sup>d</sup>		100 mol% 69% yield >10:1 d.r.	10 <sup>d</sup>		100 mol% 81% yield >10:1 d.r.
5 <sup>e</sup>		10 mol% 92% yield >10:1 d.r.	11 <sup>e,f</sup>		20 mol% 86% yield 10:1 d.r.
6 <sup>e</sup>		10 mol% 89% yield >10:1 d.r.	12 <sup>d,f</sup>		100 mol% 55% yield

<sup>a</sup> Reactions performed using 1 equiv of oxaziridine and 1.5–2 equiv of olefin in toluene unless otherwise noted. <sup>b</sup> Isolated yields. <sup>c</sup> Diastereomer ratio determined by <sup>1</sup>H NMR. <sup>d</sup> Reaction performed in CH<sub>2</sub>Cl<sub>2</sub>. <sup>e</sup> Alkene added by syringe pump over 2–3 h. <sup>f</sup> Used 3–4 equiv of alkene.

**Table 3.** Investigation of Oxaziridine Scope in Nitron Cycloadditions



$\text{Ph-CH=CH}_2 + \text{Ns-N-O} \xrightarrow[\text{toluene, 23 } ^\circ\text{C}]{\text{TiCl}_4, \text{5 \AA mol sieves}} \text{Ph-CH(O-Ns)-CH}_2\text{-R}$					
entry <sup>a</sup>	product	mol% TiCl <sub>4</sub> yield, <sup>b</sup> d.r. <sup>c</sup>	entry <sup>a</sup>	product	mol% TiCl <sub>4</sub> yield, <sup>b</sup> d.r. <sup>c</sup>
1		10 mol% 95% yield >10:1 d.r.	3		10 mol% 80% yield 10:1 d.r.
2		10 mol% 94% yield >10:1 d.r.	4		10 mol% 91% yield 5:1 d.r.

<sup>a</sup> Reactions performed using 1 equiv of oxaziridine and 1.5 equiv of olefin. <sup>b</sup> Isolated yields. <sup>c</sup> Diastereomer ratio determined by <sup>1</sup>H NMR.

constraints, however, electron-donating and -withdrawing substituents can be accommodated on the nitron precursor (entries 1–3), and other oxaziridines bearing electron-stabilizing moieties such as alkynes undergo facile rearrangement and cycloaddition under our optimized conditions (entry 4).

Synthetic manipulation of the nitron cycloadducts can be accomplished under mild conditions. Treatment of *N*-nosyl isoxazolidine **5** with thiophenol and potassium carbonate<sup>11</sup> reveals the *N*-unsubstituted isoxazolidine **6** in 78% yield (eq 1). Importantly, no products resulting from ring opening of the 1,2-isoxazolidine could be observed upon <sup>1</sup>H NMR analysis of the unpurified reaction

mixture. In contrast, treatment of **5** using sodium naphthalenide results in selective cleavage of the N–O bond of the isoxazolidine without reduction of the nosyl moiety (eq 2). Thus, our new method for generation and cycloaddition of *N*-sulfonyl nitrones enables access to either unprotected 1,2-isoxazolidine heterocycles or *N*-protected 1,3-aminoalcohols under orthogonal conditions.



This new method for the synthesis of *N*-nosyl 1,2-isoxazolidines involves the first Lewis acid catalyzed cycloaddition of *N*-sulfonyl nitrones. Efforts to understand the divergent reactivity of *N*-sulfonyl oxaziridines in the presence of titanium and copper catalysts and to develop enantioselective methods based upon these reactions are the focus of current investigations in our lab.

**Acknowledgment.** We thank the University of Wisconsin–Madison, the NSF (CHE-0645447), and the ACS Petroleum Research Fund (PRF# 44783-G1) for financial support. NMR facilities at UW-Madison are funded by the NSF (CHE-9208463, CHE-9629688) and NIH (RR08389-01). David Michaelis is acknowledged for preliminary studies.

**Supporting Information Available:** Experimental procedures and spectral data for all new compounds are provided. This material is available free of charge via the Internet at <http://pubs.acs.org>.

## References

- (1) For recent reviews, see: (a) Merino, P. In *Science of Synthesis*, Vol. 27; Padwa, A., Ed.; Thieme: Stuttgart, 2004; pp 511–580. (b) Jones, R. C. F.; Martin, J. N. In *Synthetic Applications of 1,3-Dipolar Cycloaddition Chemistry Toward Heterocycles and Natural Products*; Padwa, A., Pearson, W. H., Eds.; John Wiley: New York, 2002; pp 1–81.
- (2) Vasella's carbohydrate-derived auxiliaries are an important exception to this generalization: Vasella, A. *Helv. Chim. Acta* **1977**, *60*, 1273–1295.
- (3) Two isolated examples of *N*-acyl and *N*-carbamoyl nitron cycloadditions have been reported. Like *N*-sulfonyl nitrones, they are not isolable, and their cycloaddition chemistry was further investigated: (a) Hussain, S. A.; Sharma, A. H.; Perkins, M. J.; Griller, D. *J. Chem. Soc., Chem. Commun.* **1979**, 289–291. (b) Guinchard, X.; Vallée, Y.; Denis, J.-N. *Org. Lett.* **2005**, *7*, 5147–5150.
- (4) Pan, S. F.; Amankulor, N. M.; Zhao, K. *Tetrahedron* **1998**, *54*, 6587–6604.
- (5) (a) Rowe, S. P.; Casey, R. J.; Brennan, B. B.; Buhrlage, S. J.; Mapp, A. K. *J. Am. Chem. Soc.* **2007**, *129*, 10654–10655. (b) Buhrlage, S. J.; Brennan, B. B.; Minter, A. R.; Mapp, A. K. *J. Am. Chem. Soc.* **2005**, *127*, 12456–12457. (c) Minter, A. R.; Brennan, B. B.; Mapp, A. K. *J. Am. Chem. Soc.* **2004**, *126*, 10504–10505.
- (6) Michaelis, D. J.; Shaffer, C. J.; Yoon, T. P. *J. Am. Chem. Soc.* **2007**, *129*, 1866–1867.
- (7) (a) Emmons, W. D. *J. Am. Chem. Soc.* **1957**, *79*, 5739–5754. (b) Christensen, D.; Jørgensen, K. A.; Hazell, R. G. *J. Chem. Soc., Perkin Trans. 1* **1990**, 2391–2397.
- (8) *N*-Sulfonyl nitron **2** has been postulated to be an intermediate in the thermal decomposition of **1**: Davis, F. A.; Nadir, U. K.; Kluger, E. W. *J. Chem. Soc., Chem. Commun.* **1977**, 25–26.
- (9) The thermal isomerization of *N*-benzyl nitrones to *O*-benzyl oximes has been observed: (a) Cope, A. C.; Haven, A. C. *J. Am. Chem. Soc.* **1950**, *72*, 4896–4903. (b) Grubbs, E. J.; McCullough, J. D., Jr.; Weber, B. H.; Maley, J. R. *J. Org. Chem.* **1966**, *31*, 1098–1102.
- (10) Davis has commented upon the instability of *N*-sulfonyl oxaziridines bearing electron-rich C-aryl groups: Davis, F. A.; Lamendola, J., Jr.; Nadir, U.; Kluger, E. W.; Sedergran, T. C.; Panunto, T. W.; Billmers, R.; Jenkins, R., Jr.; Turchi, I. J.; Watson, W. H.; Chen, J. S.; Kimura, M. *J. Am. Chem. Soc.* **1980**, *102*, 2000–2005.
- (11) Fukuyama, T.; Jow, C.-K.; Cheung, M. *Tetrahedron Lett.* **1995**, *36*, 6373–6374.

JA711335D