# LETTERS

## Sc(OTf)<sub>3</sub>-Catalyzed Transfer Diazenylation of 1,3-Dicarbonyls with Triazenes via N–N Bond Cleavage

Cong Liu,<sup>†,‡,§</sup> Jian Lv,<sup>†,‡</sup> Sanzhong Luo,<sup>\*,†,‡</sup> and Jin-Pei Cheng<sup>†,‡,§</sup>

<sup>†</sup>Beijing National Laboratory for Molecular Sciences (BNLMS), CAS Key Laboratory for Molecular Recognition and Function, Institute of Chemistry, Chinese Academy of Sciences, Beijing 100190, China

<sup>‡</sup>Collaborative Innovation Center of Chemical Science and Engineering (Tianjin), Tianjin 300071, China

<sup>§</sup>Stake Key Laboratory of Elemento-Organic Chemistry, Department of Chemistry, Nankai University, Tianjin 300071, China

Supporting Information

**ABSTRACT:** A new and efficient method for diazenylation reactions was developed with a  $Sc(OTf)_3$ -catalyzed nitrogennitrogen bond cleavage process with triazenes. The transfer diazenylation reactions accommodate a diverse range of active methylene substrates including simple ketones to give aliphatic azo compounds that are of significant potential as azo prodrugs in high yields under mild conditions.



zo derivatives are commonly utilized in both laboratories **A** and industry as indicators, radical reaction initiators, azo ligands of metal complexes, dyes and pigments, therapeutic agents, modern material sciences, and food additives.<sup>1,2</sup> Reflecting on the development of azo syntheses, classic and general methods in the formation of nitrogen-nitrogen double bonds include electrophilic azo coupling reactions of diazonium salts, Mills coupling of anilines with aromatic nitroso and nitro compounds, and Wallach rearrangements of azoxy derivatives.<sup>3</sup> Recent advances in the synthesis of azo compounds mainly involve catalytic reductive aromatic nitro derivatives,<sup>4</sup> oxidation of anilines,<sup>5</sup> as well as dehydrogenation of arylhydrazines<sup>6</sup> (Scheme 1, A). Development of a chemoselective diazenylation strategy is still in high demand in order to address remaining issues such as excess use of explosive diazonium salts, harsh reaction conditions, and invariably accompanying oxidant or reductant waste. In this regard, the synthesis of aliphatic azo

## Scheme 1. Synthetic Strategies for Azo Derivatives





compounds is particularly challenging due to the lack of functional group tolerance of most of the available synthetic procedures as well as the inherent instability of the aliphatic azo moiety. Recently, aliphatic azo compounds with the general structures ArN = N-alkyl have been developed as prodrugs with promising antibiotic and anti-inflammatory activity,<sup>7</sup> addressing significant demands for economic and selective synthetic protocols. Herein, we present a conceptually distinctive azo-transfer reaction in which an aryl azo moiety can be directly transferred from aryl triazene and installed chemoselectively onto aliphatic carbons under rather mild conditions.

Aryl triazenes (ArN=NNR<sub>1</sub>R<sub>2</sub>), are a class of compounds with intriguing structural and chemical properties and a long history dating back to 1875.<sup>8,9</sup> Recent pursuit of chemoselective methodology has sparked new interest in the chemistry of aryltriazenes. Accordingly, aryltriazenes have been explored as precursors of phenylium cations or radicals in arylation and dehydrogenation reactions.<sup>10,11</sup> The triazene moiety can also be utilized as a directing group in C–H functionalization<sup>12</sup> as well as in oxidative indole synthesis<sup>13</sup> (Scheme 1, B). Though typically regarded as masked diazonium ions, aryl triazenes have seldom been employed in diazenylation reactions. In a single instance, intramolecular coupling of triazene with alkyne has been reported to give cinnolines under either thermal<sup>14</sup> or Pdcatalytic conditions.<sup>15,16</sup> Intermolecular azo-transfer with aryl triazenes has surprisingly not been reported so far.

Initially, we sought to develop an asymmetric  $\alpha$ -arylation reaction using aryl triazenes as phenyl precursors under enamine catalysis; however, no  $\alpha$ -arylation product was detected in our initial experiments with our previously developed chiral primary amines.<sup>17</sup> Instead, we found Lewis acid Sc(OTf)<sub>3</sub> can promote transfer diazenylation with aryl

Received: September 12, 2014 Published: October 8, 2014 triazene via efficient N–N cleavage, which enables chemoselective aryl diazenylation of  $\beta$ -dicarbonyls beyond the reach otherwise<sup>18</sup> (Scheme 1, B), and this diazenylation reaction was hence further explored.

Our investigation began with the phenyl diazenylation of 2methyl acetoacetate 1a, and the representative results are summarized in Table 1. To our gratification, we found that

Table 1. Optimization for Phenyl Diazenylation Reaction of2-Methyl Acetoacetate $^a$ 

		R' N_R'	20 m CH <sub>2</sub> C	ol % Lewis : Cl <sub>2</sub> , 40 °C, 4	$\frac{acid}{8 h} \xrightarrow{O}_{N=1}^{O}$	-0 -N
1a	2					Ph
		<b>2a</b> , R :	= N( <i>i</i> Pr)	2		
	N≈N <sup>-R</sup>	2b, R =			N <sub>2</sub> BF <sub>4</sub>	20
		20, R =	Morpho	oline		20
	Triazene				Diazonium Sal	t
entry	Lewis acid	rea	igent	solvent	time (h)	yield <sup>b</sup> (%)
1	Sc(OTf) <sub>3</sub>	2	2a	$CH_2Cl_2$	48	75
2	Sc(OTf) <sub>3</sub>	2	2b	$CH_2Cl_2$	48	48
3	Sc(OTf) <sub>3</sub>	2	2c	$CH_2Cl_2$	48	66
4	Sc(OTf) <sub>3</sub>	2	2d	$CH_2Cl_2$	48	69
5	Sc(OTf) <sub>3</sub>	2	2e	$CH_2Cl_2$	24	N.D.
6 <sup><i>c</i></sup>	Sc(OTf) <sub>3</sub>	2	2a	EtOH	48	trace
$7^c$	Sc(OTf) <sub>3</sub>	2	2a	DMSO	48	N.D.
8 <sup>c</sup>	Sc(OTf) <sub>3</sub>	2	2a	$CH_3CN$	48	39
$9^d$	Sc(OTf) <sub>3</sub>	2	2a	hexane	48	43
10	Sc(OTf) <sub>3</sub>	2	2a	ether	48	69
$11^e$	$BF_3$ ·Et <sub>2</sub> O (2 equiv)	2	2a	$CH_2Cl_2$	3	31
12	FeCl <sub>3</sub>	2	2a	$CH_2Cl_2$	48	46
13	Bi(OTf) <sub>3</sub>	2	2a	$CH_2Cl_2$	48	32
14	CuBr <sub>2</sub>	2	2a	$CH_2Cl_2$	48	N. D.
15	$Hf(OTf)_4$	2	2a	$CH_2Cl_2$	48	72
16	Sc(OTf) <sub>3</sub> (10 mol 9	6) 2	2a	$CH_2Cl_2$	48	50
17 <sup>f</sup>	Sc(OTf) <sub>3</sub> (10 mol 9	6) 2	2a	$CH_2Cl_2$	24	54
$18^{g,h}$	Sc(OTf) <sub>3</sub> (10 mol 9	6) 2	2a	$CH_2Cl_2$	24	85
19 <sup>g,h</sup>	none	2	2a	$CH_2Cl_2$	24	<10

<sup>*a*</sup>General conditions: **1a** (0.1 mmol), **2** (0.1 mmol), and Lewis acid (20 mol %) at 40 °C in the solvent (0.4 mL) under Ar. <sup>*b*</sup>Isolated yield. <sup>*c*</sup>80 °C. <sup>*d*</sup>60 °C. <sup>*e*</sup>At room temperature. <sup>*f*</sup>CH<sub>3</sub>COCl (0.1 mmol) as the additive. <sup>*g*</sup>TMSCl (0.1 mmol) as the additive. <sup>*h*</sup>**1a** (0.12 mmol). N.D. = no product was observed by TLC.

Lewis acids were viable catalysts for the diazenylation reactions. Among various Lewis acids screened (Table 1, entries 1 and 11–15),  $Sc(OTf)_3$  was eventually identified as an optimal Lewis acid for this reaction (Table 1, entry 1). The expected diazenvlation adduct (3a) can be obtained in 75% yield. The nature of the terminal tertiary amine substituent of triazenes was then tested, with similar results for methyl, isopropyl, benzyl, and cyclic morpholine groups (Table 1, entries 2-4). Surprisingly, the use of the common diazonium salt 2e led to no desired product (Table 1, entry 5), highlighting the chemoselective nature of aryl triazenes. Solvent effects have also been examined, and the reactions were found to work poorly in other solvents such as EtOH, DMSO, CH<sub>3</sub>CN, hexane, and ether (Table 1, entries 6-10). The choice of acidic additives significantly influenced the yield (Table 1, entries 16-18), especially to those inert and bulky substrates, with TMSCl proving to be the optimal reaction additive, presumably due to the capture of the liberating diisopropylamine in the catalytic system (Table 1, entry 18, 85% yield over 24 h). It was noteworthy that, in the absence of  $Sc(OTf)_{3}$ , almost no diazenylation product 3a was obtained (Table 1, entry 19). Therefore, it can be concluded that the optimized reaction should be performed under the catalysis of 10 mol % of  $Sc(OTf)_3$  in the presence of 1.0 equiv of TMSCl in  $CH_2Cl_2$ .

The substrate scope with 1,3-dicarbonyls was next explored (Scheme 2). The diazenylation all occurred at the  $\alpha$ -position as

Scheme 2. Scope of Intermolecular Diazenylation of Active Methylene Compounds and Triazophenyl Reagent $^{a,b}$ 



<sup>*a*</sup>General conditions: **1** (0.12 mmol), **2** (0.1 mmol), TMSCl (0.1 mmol), and  $Sc(OTf)_3$  (10 mol %) at 40 °C in the  $CH_2Cl_2$  (0.4 mL) under Ar. <sup>*b*</sup>Isolated yield. <sup>*c*</sup>2 equiv of **1** and 2 equiv of TMSCl. <sup>*d*</sup>3 equiv of ketone in 80 °C.

expected. The C–N bond formation proceeded in consistently high yield for many acetoacetate derivatives in which the ester moiety ( $R_3$ ) ranges in size from the ethyl group to the steric adamantyl group (3a-e). The reaction also worked on a larger scale (2 mmol) with comparable 79% yield (3a). The influence of  $R_1$  and  $R_2$  groups of 1,3-dicarbonyls was also investigated. With ethyl ketone ( $R_1 = Et$ ) and phenyl ketone ( $R_1 = Ph$ ), the corresponding products **3f** and **3g** were obtained in excellent yield. When unsubstituted ethyl benzoylacetate **1h** was used as the reactant, the product was isolated in 95% yield as a major keto isomer (keto/enol  $\approx$  7:1). The cyclic 1,3-ketoester methyl 2-carboxylate indanone (**1j**) and tetralone (**1k**) also proceeded

well, giving the products in moderate yields. Pleasingly, propargyl (1i) and heterocyclic (1l) substituents were welltolerated to give the desired adducts in 88% and 75% yields, respectively.  $\beta$ -Diketones possess a more acidic C–H bond and a higher enol/keto- ratio than  $\beta$ -keto esters,<sup>19</sup> and this type of substrate worked extremely well to give the diazenvlation adducts in 94% and 91% yield (3m and 3n). Moreover, a variety of active methylene compounds, such as the substituted malonic ester (10), ethyl cyanacetate (1p),  $\beta$ -keto amide (1q),  $\beta$ -keto phosphonate (1r), and  $\alpha$ -(phenylsulfonyl)acetophenone (1s), were well compatible with the reaction conditions. In particular, the reaction also worked for arvl ketones such as propiophenone 1t, furnishing the  $\alpha$ -azo adduct in 40% yield. Other simple ketones such as cyclohexanone and 3-pentanone have also been examined in the reactions, showing unfortunately rather poor reactivity, likely a result of their low  $\alpha$ acidity.

Furthermore, the scope of aromatic triazo reagents has also been examined (Scheme 3). The phenyl ring moiety can be

## Scheme 3. Scope of Intermolecular Diazenylation of $\beta$ -Diketones and Aromatic Triazenes<sup>*a,b*</sup>



<sup>*a*</sup>General conditions: **1m** (0.1 mmol), **2** (0.2 mmol), TMSCl (0.2 mmol), and  $Sc(OTf)_3$  (10 mol %) at 40 °C in the  $CH_2Cl_2$  (0.4 mL) under Ar. <sup>*b*</sup>Isolated yield.

*ortho-, meta-,* or *para-substituted, which can bear an electron-*donating or an electron-withdrawing group, including a halide or ester group. The desired aromatic azo products can be obtained in generally good yields.

In light of the above results and previous reports, a plausible mechanism is described in Scheme 4. The  $\beta$ -ketoester is activated usually through coordination to the Sc(III) center in a bidentate fashion by both of the carbonyl groups. Meanwhile,





the silyl enol ether of an alkenyl  $\beta$ -keto ester might exist in view of the effect to enhance the equilibrium of the enol/keto ratio, since its generation in situ from the reaction of an alkenyl  $\beta$ keto ester and a silyl chloride has been well demonstrated.<sup>20</sup>

The reaction is initialize via the coordination of the pendent triazene reagent to  $Sc(OTf)_3$  on the nitrogen—nitrogen double bond. The marked affinity of the trimethylsilyl group toward terminal nitrogen lone pair and the weak nucleophilic nature of chloride ions restrain the nucleophilic attack to furnish aryl halides in the present of TMSCI.<sup>21</sup> Afterward, conjugate addition of the active  $\beta$ -keto ester and triazene compound via the intermediate III<sup>15,22</sup> occurs to generate the desired product and HN(*i*-Pr)<sub>2</sub>, and the latter might be trapped by TMSCl or HCl to enhance the catalytic turnovers.

Taking the requirement for  $\beta$ -keto ester activation using Lewis acids into consideration, we hypothesized that the unique binary acid system developed by our group<sup>23</sup> might catalyze this new and enantioselective C–N bond formation. Indeed, we discovered that the desired aromatic azo compound was obtained in 57% yield and 24% ee in Sc(OTf)<sub>3</sub>/polyfluoro chiral phosphoric acid without further optimizations (Scheme 5).





 $^a{\rm The}$  reaction was conducted on 0.05 mmol scale in 0.4 mL of  ${\rm CH_2Cl_2}$  at 40  $^{\circ}{\rm C}$  under Ar.

In summary, we have developed a  $Sc(OTf)_3$ -catalyzed diazenylation reaction of active methylene compounds with triazenes via N–N cleavage. This simple transfer diazenylation method was shown to tolerate a range of substrates to afford aliphatic azo compounds that are beyond reach with other methods. Further studies are currently underway to develop a catalytic asymmetric version.

## ASSOCIATED CONTENT

## **Supporting Information**

Experimental procedures, characterization data, and <sup>1</sup>H NMR, <sup>13</sup>C NMR, and HPLC spectra for new products. This material is available free of charge via the Internet at http://pubs.acs.org.

## AUTHOR INFORMATION

## Corresponding Author

\*E-mail: luosz@iccas.ac.cn.

## Notes

The authors declare no competing financial interest.

## ACKNOWLEDGMENTS

We thank the Natural Science Foundation of China (21390400, 21025208, 21202170, and 21472193) and the National Basic Research Program of China (2012CB821600) for financial

#### **Organic Letters**

support. S.L. is supported by the National Program of Topnotch Young Professionals and CAS Youth Innovation Promotion Association.

#### REFERENCES

 For representative applications, see: (a) Hunger, K. In Industrial Dyes: Chemistry, Properties, Applications; Wiley-VCH: Weinheim, 2003.
 (b) Zhao, Y.; Ikeka, T. In Smart Light Responsive Materials: Azobenzene-Containing Polymers and Liquid Crystals; Wiley: Chichester, 2009. (f) Bandyopadhyay, A.; Higuchi, M. Eur. Polym. J. 2013, 49, 1688.

(2) For selected examples, see: (a) Ikeda, T.; Tsutsumi, O. Science
1995, 268, 1873. (b) Natansohn, A.; Rochon, P. Chem. Rev. 2002, 102,
4139. (c) Banerjee, I. A.; Yu, L.; Matsui, H. J. Am. Chem. Soc. 2003,
125, 9542. (d) Yu, Y.; Nakano, M.; Ikeda, T. Nature 2003, 425, 145.
(e) Cisnetti, F.; Ballardini, R.; Credi, A.; Gandolfi, M. T.; Masiero, S.;
Negri, F.; Pieraccini, S.; Spada, G. P. Chem.—Eur. J. 2004, 10, 2011.
(f) Venkataramani, S.; Jana, U.; Dommaschk, M.; Sönnichsen, F. D.;
Tuczek, F.; Herges, R. Science 2011, 331, 445. (g) Szymanski, W.;
Beierle, J. M.; Kistemaker, H. A. V.; Velema, W. A.; Feringa, B. L.
Chem. Rev. 2013, 113, 6114.

(3) (a) Boyer, J. H. In The Chemistry of the Nitro and Nitroso Groups; Interscience: New York, 1969; Part 1. (b) Buncel, E. Acc. Chem. Res. 1975, 8, 132. (c) Szele, I.; Zollinger, H. Top. Curr. Chem. 1983, 112, 1.
(d) Zollinger, H. In Diazo Chemistry I: Aromatic and Heteroaromatic Compounds; Wiley-VCH: Weinheim, 1994. For recent reviews, see:.
(e) Hamon, F.; Djedaini-Pilard, F.; Barbot, F.; Len, C. Tetrahedron 2009, 65, 10105. (f) Merino, E. Chem. Soc. Rev. 2011, 40, 3835.

(4) (a) Grirrane, A.; Corma, A.; García, H. Science 2008, 322, 1661.
(b) Zhu, H.; Ke, X.; Yang, X.; Sarina, S.; Liu, H. Angew. Chem., Int. Ed. 2010, 49, 9657.
(c) Guo, X.; Hao, C.; Jin, G.; Zhu, H.-Y.; Guo, X. Angew. Chem., Int. Ed. 2014, 53, 1973.
(d) Liu, X.; Li, H.-Q.; Ye, S.; Liu, Y.-M.; He, H.-Y.; Cao, Y. Angew. Chem., Int. Ed. 2014, 53, 7624.

(5) (a) Lu, W.; Xi, C. Tetrahedron Lett. 2008, 49, 4011. (b) Grirrane, A.; Corma, A.; Garcia, H. Nat. Protoc. 2010, 5, 429. (c) Zhang, C.; Jiao, N. Angew. Chem., Int. Ed. 2010, 49, 6174. (d) Zhu, Y.; Shi, Y. Org. Lett. 2013, 15, 1942. (e) Jiang, B.; Ning, Y.; Fan, W.; Tu, S.-J.; Li, G. J. Org. Chem. 2014, 79, 4018.

(6) (a) Drug, E.; Gozin, M. J. Am. Chem. Soc. 2007, 129, 13784.
(b) Gao, W.; He, Z.; Qian, Y.; Zhao, J.; Huang, Y. Chem. Sci. 2012, 3, 883.
(c) Bai, L.-S.; Gao, X.-M.; Zhang, X.; Sun, F.-F.; Ma, N. Tetrahedron Lett. 2014, 55, 4545.

(7) (a) Dhaneshwar, S. S.; Gairola, N.; Kandpal, M.; Vadnerkar, G.; Bhatt, L.; Rathi, B.; Kadam, S. S. *Eur. J. Med. Chem.* 2009, 44, 3922.
(b) Cassano, R.; Trombino, S.; Cilea, A.; Ferrarelli, T.; Muzzalupo, R.; Picci, N. *Chem. Pharm. Bull.* 2010, 58, 103. (c) Kennedy, D. A.; Vembu, N.; Fronczek, F. R.; Devocelle, M. *J. Org. Chem.* 2011, 76, 9641.

(8) Baeyer, A. V.; Jaeger, C. Ber. Dtsch. Chem. Ges. 1875, 8, 148.

(9) (a) Nicolaou, K. C.; Boddy, C. N. C. J. Am. Chem. Soc. 2002, 124, 10451. (b) Kimball, D. B.; Haley, M. M. Angew. Chem., Int. Ed. 2002, 41, 3338. (c) Bräse, S. Acc. Chem. Res. 2004, 37, 805. (d) Liu, C.-Y.; Knochel, P. J. Org. Chem. 2007, 72, 7106.

(10) (a) Yang, W.; Zhou, J.; Wang, B.; Ren, H. Chem.—Eur. J. 2011, 17, 13665. (b) Zhao, G.; Wang, B.; Yang, W.; Ren, H. Eur. J. Org. Chem. 2012, 6236. (c) Zhou, J.; Yang, W.; Wang, B.; Ren, H. Angew. Chem, Int. Ed. 2012, 51, 12293.

(11) Voica, A.-F.; Mendoza, A.; Gutekunst, W. R.; Fraga, J. O.; Baran, P. S. Nat. Chem. **2012**, *4*, 629.

(12) (a) Hafner, A.; Bräse, S. Angew. Chem., Int. Ed. 2012, 51, 3713.
(b) Wang, C.; Chen, H.; Wang, Z.; Chen, J.; Huang, Y. Angew. Chem., Int. Ed. 2012, 51, 7242.

(13) Wang, C.; Sun, H.; Fang, Y.; Huang, Y. Angew. Chem., Int. Ed. 2013, 52, 5795.

(14) (a) Kimball, D. B.; Herges, R.; Haley, M. M. J. Am. Chem. Soc.
2002, 124, 1572. (b) Kimball, D. B.; Weakley, T. J.; Herges, R.; Haley,
M. M. J. Org. Chem. 2002, 67, 6395. (c) Kimball, D. B.; Weakley, T. J.;
Herges, R.; Haley, M. M. J. Am. Chem. Soc. 2002, 124, 13463.

(15) Zhu, C.; Yamane, M. Tetrahedron 2011, 67, 4933.

(16) Goeminne, A.; Scammells, P. J.; Devine, S. M.; Flynn, B. L. Tetrahedron Lett. 2010, 51, 6882.

(17) For an account, see: Zhang, L.; Luo, S. Synlett 2012, 23, 1575.
(18) (a) Natansohn, A.; Rochon, P. Chem. Rev. 2002, 102, 4139.
(b) Sourisseau, C. Chem. Rev. 2004, 104, 3851. (c) Japp, F. R.; Klingemann, F. Ber. Dtsch. Chem. Ges. 1887, 20, 2942. (d) Garst, M. E.; Lukton, D. Synth. Commun. 1980, 10, 155. (e) Sakakura, T.; Tanaka, M. J. Chem. Soc., Chem. Commun. 1985, 1309. (f) Sakakura, T.; Hara, M.; Tanaka, M. J. Chem. Soc., Perkin Trans. 1 1994, 289. (g) Nelson, H. M.; Reisberg, S. H.; Shunatona, H. P.; Patel, J. S.; Toste, F. D. Angew. Chem., Int. Ed. 2014, 53, 5600.

(19) Bordwell, F. G. Acc. Chem. Res. 1988, 21, 456.

(20) (a) Langer, P.; Eckardt, T.; Saleh, N. N. R.; Karime, I.; Müller, P. *Eur. J. Org. Chem.* **2001**, 3657. (b) Han, X.; Wang, X.; Pei, T.; Widenhoefer, R. A. *Chem.—Eur. J.* **2004**, *10*, 6333 and references cited therein.

(21) Ku, H.; Barrio, J. R. J. Org. Chem. 1981, 46, 5239.

(22) Fang, Y.; Wang, C.; Su, S.; Yu, H.; Huang, Y. Org. Biomol. Chem. 2014, 12, 1061.

(23) (a) Lv, J.; Zhang, L.; Zhou, Y.; Nie, Z.; Luo, S.; Cheng, J.-P. Angew. Chem., Int. Ed. **2011**, 50, 6610. (b) Lv, J.; Zhang, L.; Luo, S.; Cheng, J.-P. Angew. Chem., Int. Ed. **2013**, 52, 9786.