

Article

Protein-Metal-Ion Interactions Studied by Mass Spectrometry-Based Footprinting with Isotope-Encoded Benzhydrazide

Chunyang guo, Ming Cheng, and Michael L. Gross

Anal. Chem., **Just Accepted Manuscript** • DOI: 10.1021/acs.analchem.8b04088 • Publication Date (Web): 29 Nov 2018

Downloaded from <http://pubs.acs.org> on December 2, 2018

Just Accepted

"Just Accepted" manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides "Just Accepted" as a service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. "Just Accepted" manuscripts appear in full in PDF format accompanied by an HTML abstract. "Just Accepted" manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are citable by the Digital Object Identifier (DOI®). "Just Accepted" is an optional service offered to authors. Therefore, the "Just Accepted" Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the "Just Accepted" Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these "Just Accepted" manuscripts.



ACS Publications

is published by the American Chemical Society, 1155 Sixteenth Street N.W., Washington, DC 20036

Published by American Chemical Society. Copyright © American Chemical Society. However, no copyright claim is made to original U.S. Government works, or works produced by employees of any Commonwealth realm Crown government in the course of their duties.

Protein-Metal-Ion Interactions Studied by Mass Spectrometry-Based Footprinting with Isotope-Encoded Benzhydrazide

Chunyang Guo, Ming Cheng, Michael L. Gross*

Department of Chemistry, Washington University, St. Louis, Missouri 63130, United States

ABSTRACT: Metal ions, usually bound by various amino-acid side chains in proteins, play multiple roles in protein folding, conformational change, cellular communication, and catalysis. Ca(II) and Mg(II), abundant among biologically relevant cations, execute their cellular functions associated with the conformational change of bound proteins. They bind with proteins where carboxylic acid residues are dominant ligands. To develop mass spectrometry for mapping protein-binding sites, we implemented a new carboxyl group footprinter, benzhydrazide, and refined it with isotope encoding. The method uses carbodiimide chemistry to footprint carboxylic residues, whereby 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide activates a carboxyl group followed by nucleophilic attack by benzhydrazide forming a stable labeled product. We tested the effectiveness of isotope-encoded benzhydrazide by studying Ca²⁺ and Mg²⁺ binding of calmodulin, an EF-hand protein. The footprinting results indicate that the four active sites for metal-ion binding (EF hands I, II, III, and IV) and the linker region (peptide 78-86) undergo conformational changes upon Ca(II) and Mg(II) binding, respectively. The outcome is consistent with previously reported results and 3-D structures, thereby validating a new reagent that is more reactive and discriminating for specific amino-acid protein footprinting. This reagent should be important for locating metal-binding sites of other metalloproteins.

Metal ions are essential for many cellular functions.¹ In the range of 25 – 50% of all proteins found within an organism contain metal ions.^{2,3} Metal ions bind via functional groups termed ligands that donate lone pairs to the metal ion. In the bound state, the metal ions in proteins can catalyze a variety of biochemical reactions or stabilize the protein structures.

Based on the properties of coordinated metals, we can classify two types of metals that interact with proteins: 1) borderline/soft ions (e.g., Zn(II), Fe(II) and Cu(II)) that coordinate with nitrogen- and sulfur-containing groups found in His and Cys, and 2) hard cations (e.g., Ca(II) and Mg(II)) that use oxygen ligands as the preferred coordination sites (e.g., Asp and Glu are preferred).⁴ Among all metalloproteins, those containing Mg(II), Ca(II), are relatively abundant.¹⁻⁵ In many cases, Mg(II) and Ca(II) binding stabilizes the structure of folded proteins, and in some cases, locks in a physiologically active conformation. Specifically, Ca(II) plays a role in signal transduction, muscle contraction, nerve-impulse transmission, and calcium homeostasis. To maintain the correct Ca ion concentration in the intra- and extracellular space, the body uses Ca pumps. Most Ca-binding proteins possess a highly conserved Ca-binding motif, the EF hand, that selectively binds Ca in a background of up to 10⁵-fold higher concentrations of Na(I), K(I), and Mg(II).^{6,7} Although divalent Mg has higher charge density and exists at higher concentration in the cytosol than Ca(II), it shares physical and chemical properties with Ca(II) to bind proteins and enable signaling, enzymatic activation and catalysis, and muscle relaxation.^{8,9} A clear knowledge of metal/protein interactions, including the location of the metal binding site and its binding characteristics, is essential to understand the functional mechanisms of metal-binding proteins.

Although nuclear resonance (NMR),¹⁰ X-ray crystallography,³ and Cryo-EM¹¹ are commonly used to determine metal-binding sites, they are limited by size, conformational flexibility, aggregation propensity, or limited sample amount, rendering them frequently inapplicable. Our goal is to develop mass spectrometry (MS)-based methods that complement these high-resolution techniques because MS is general, fast, sensitive, has high-throughput, and provides moderate resolution. One MS-based method, hydrogen deuterium exchange (HDX), has been widely applied in protein structure characterization.¹²⁻¹⁵ HDX can be extended to provide an approach termed protein-ligand interactions by MS, titration, and HD exchange (PLIMSTEX) to monitor the conformational changes of proteins upon Ca²⁺ binding and even measure the binding constants.¹⁶ HDX, however, cannot directly monitor the binding of side chains owing to their fast hydrogen deuterium exchange.

An alternative method for solving this challenge is irreversible covalent labeling whereby a variety of reactions can be invoked to modify amino-acid side chains. In this approach, reactive side chains are stably modified such that the label can be conveniently located by MS in peptides from proteolysis. One approach is to use reactive species such as OH radicals (provided through electron beam or X-ray radiolysis, Fenton chemistry, or photolysis of peroxide),¹⁷ carbenes,^{18,19} iodide radicals,²⁰ and trifluoromethyl radicals.²¹ Slower but still effective footprinting can use carbodiimide²², ICAT,²³ or diethylpyrocarbonate (DEPC)²⁴ chemistries. Because the reactions are slower than those of free radicals, there is always an issue that the labeling itself doesn't perturb the native structures.

Carbodiimide chemistry provides the most versatile method for labeling or crosslinking carboxylic acids. Earlier, we employed 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride (EDC), a crosslinking facilitator, to enable glycine ethyl ester (GEE) to footprint side chains of Asp and Glu. This approach was successfully applied to determine the orientation of the FMO antenna protein,²⁵ to map the binding of Fab-1 and VEGF,²⁶ to locate a membrane-associated tyrosine kinase binding site,²⁷ to map the epitope of human interleukin-6 receptor,²⁸ to determine the binding interface for the oligomeric apolipoprotein E,²⁹ to characterize monoclonal antibody structure,³⁰ and ELISA antibody-antigen interactions.³¹ Nevertheless, GEE footprinting has limitations, including the requirement for a large excess of the GEE reagent (20,000 equivalents with respect to the protein), some hydrolysis of the product during work-up, and a limited opportunity to incorporate isotopes for encoding.

Herein, we explore application of carbodiimide compound (EDC), in the presence of new labeling reagent benzhydrazide (BHD), inspired by Leitner.³² To be specific, carboxylic acid residues are activated by EDC, then BHD, a nucleophilic modifying reagent, attacks the activated carboxyl group. Our goal is to avoid the hydrolysis of the modified product while still characterizing metal-binding sites. We focus on two hard metal ions Ca(II) and Mg(II) that chelate with carboxylic residues because they are ubiquitous in metalloproteins. We chose carbodiimide specifically because it reacts with carboxylic groups whose reactivity will be reduced when chelated with Ca(II) or Mg(II) ions. Furthermore, its size may allow high discrimination for metal-bound vs. unbound carboxylates. To improve the MS quantification, we incorporated five deuteriums in the structure to permit encoding.³³ Results show that the method can accurately measure the binding site of Ca(II) and Mg(II) in a test protein, calmodulin (CaM). By assisting in the elucidation of metal-binding sites in proteins, this method should enrich the output of structural proteomics studies.

EXPERIMENTAL SECTION

Materials. Tris base, HEPES buffer (pH 7.5), urea, water, formic acid, glycine ethyl ester hydrochloride (GEE), calcium chloride, magnesium chloride, ammonium acetate, 1,1'-carbonyldiimidazole, benzoic acid, benzoic acid-2,3,4,5,6-*d*₅, tetrahydrofuran (THF), hydrazine hydrate solution (78-82%), Z-Glu-OMe, and thin-layer chromatography supplies (TLC, silica gel 60 F254) were obtained from Sigma-Aldrich (St. Louis, MO). *N*-Benzyloxycarbonyl-*L*-aspartic acid 1-methyl ester was obtained from Alfa Aesar (Lancashire, UK). Sartorius™ Vivaspin™ 500 μ L centrifugal concentrators were purchased from Sartorius Corporation (Goettingen, Germany). The protein bovine calmodulin (CaM) was from Ocean Biologics (Seattle, WA). The 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride (EDC), trypsin and Zeba™ spin desalting columns (7K MWCO) were from Thermo Scientific (Rockford, IL). Cyclo(Arg-Gly-Asp-D-Phe-Glu) was from Peptides International, Inc. (Louisville, KY).

Synthesis and Characterization of Benzhydrazide and *d*₅-Benzhydrazide. Reagents (no known hazards) were synthesized based on protocol of Li,³⁴ (Figure S1) except for the purification process. In 100 mL reaction flask, 1,1'-carbonyldiimidazole (1.73 g, 10.6 mmol) was added to a solution of benzoic acid (1.00 g, 8.19 mmol) in THF (15 mL). After stirring at 25 °C for 3 h, the mixture

was added dropwise to the hydrazine hydrate solution (78-82%, 1.23 g, 24.6 mmol) in 6 mL THF over 45 min. The reaction was allowed to go to completion overnight at 25 °C, as monitored by TLC. Solvents were evaporated in vacuum to give a crude white product that was isolated and purified by silica gel column chromatography (hexane:ethyl acetate = 1:1) to give the benzhydrazide as a white solid (1.03 g, 93% yield). *d*₅-Benzhydrazide was synthesized by using an identical procedure, giving the *d*₅-benzhydrazide as a white solid (1.00 g, 87% yield). High-resolution mass spectra (HRMS) were recorded with a Bruker MaXis QToF (ESI). HRMS (ESI) *m/z* calcd. for protonated BHD (C₇H₈N₂OH⁺): 137.0715. Found: 137.0714. HRMS (ESI) *m/z* calcd. for *d*₅-BHD (C₇H₃D₅N₂OH⁺): 142.1029. Found: 142.1028. (Figure S2)

Preparation of Protected amino acids/Peptide/Protein Stock Solution. All protected amino acids (Z-Glu-OMe and *N*-Benzyloxycarbonyl-*L*-aspartic acid 1-methyl ester) and cyclic peptide stock solutions of 100 μ M concentration were prepared in 10 mM HEPES buffer. Ca²⁺-free CaM (50 μ M) was prepared by washing three times with HEPES buffer at 15,000 (xg) speed using Vivaspin™ centrifugal concentrators, whereas Ca²⁺-bound CaM was prepared by incubating 50 μ M CaM with 500 μ M CaCl₂ overnight at 25 °C. Similarly, Mg²⁺-bound CaM was prepared by incubating 50 μ M CaM with 500 μ M MgCl₂ overnight at 25 °C. The concentration of CaM in 10 mM HEPES buffer (pH 7.5) was determined by using UV absorbance (molar absorptivity of 2980 Lmol⁻¹ cm⁻¹ at 280 nm) with a NanoDrop spectrophotometer (Thermo Scientific).

Carboxyl Group Modification. The final concentrations of protected amino acids, peptide, or protein for all reactions were adjusted to 10 μ M. For protected amino acids, the labeling was initiated by adding BHD and EDC to give final concentrations of 20 mM and 500 μ M, respectively. The protected amino acid modification was quenched after 20 min by adding 20 μ L ammonium acetate (1 M). For cyclic peptides, the labeling follows the same procedure as protected amino acids, except the labeling was quenched after 24 h. Experiments with CaM were performed in triplicate by using 20 mM BDH and 500 μ M EDC, identical to the concentration used for cyclic peptide labeling. After 5 min of reaction at 37 °C, the solution was quenched by removing the reagents by using a Zeba spin desalting column according to the manufacturer's protocol. (Figure S3)

Trypsin Proteolysis Protocol. After desalting, the filtrate was evaporated to dryness in a vacuum centrifuge. The pellets were dissolved with 50 μ L water to make an equal volume of protein. Aliquots of 10 μ L filtrate of BHD and *d*₅-BHD were combined at 1:1 ratio (v/v) and dried in vacuum (the dry sample can be stored in a freezer at -80 °C for months). Prior to analysis, the protein mixture pellet was denatured with urea (8 M, 5 μ L) at 37 °C for 30 min, followed by adding 45 μ L Tris buffer (pH 8.5). Protein samples were then digested with trypsin at a protease:protein ratio of 1:20 (w/w) at 37 °C overnight. Pure formic acid (1 μ L) was added to stop the digestion.

Solvent Accessible Surface Area (SASA) calculation. We used PDB files of Ca²⁺-free CaM (PDB ID: 1CFC) and Ca²⁺-bound CaM (PDB ID: 1CLL) to GETAREA (<http://curie.utmb.edu/getarea.html>) to calculate individual side-chain SASA.

Mass Spectrometry. All mass spectra for intact proteins were acquired on a Bruker MaXis Q-TOF (Bremen, Germany) in the positive-ion mode, as was reported previously.¹⁸

For proteolytic peptides, digests were analyzed on a QExactive Plus hybrid quadrupole orbitrap mass spectrometer coupled with a Nanospray Flex ion source (Thermo Fisher, Santa Clara, CA). A C18 column was custom fabricated with an integrated emitter (75 $\mu\text{m} \times 150 \text{ mm}$, 1.8 μm , 100 \AA). The gradient was from 2.0% solvent B (80% acetonitrile, 0.1% formic acid) to 20% solvent B over 55 min, then to 50% solvent B over 25 min, and finally to 90% solvent B for 10 min followed by a 15 min re-equilibration step. The 15 most abundant molecular ions were automatically chosen for fragmentation (DDA) by using the mass spectra scanned from m/z 380–2200 at a resolving power (RP) of 70 K (at m/z 200) throughout the chromatography. Precursor ions were isolated in the quadrupole with an isolation window of 2.0 m/z and fragmented with higher-energy collisional dissociation (HCD) with a normalized collision energy (NCE) of 32%. The automatic gain control (AGC) targets were 5×10^5 for MS and 5×10^4 for MS/MS acquisitions. Maximum injection times (maxIT) were 200 ms for MS and 100 ms for MS/MS.

LC-MS/MS Data Analysis The LC-MS/MS data were searched for unmodified and modified CaM tryptic peptides by using ByonicTM Software (Protein Metrics, San Carlos, CA). All BHD and d_5 -BHD modifications on Asp and Glu were added as modifications to the database. The parameters were: 10 ppm precursor mass tolerance, 60 ppm fragment mass tolerance, and CID/HCD fragmentation. Modification-sites were validated by manually checking product-ion spectra and were double-checked with Thermo Xcalibur (a custom program from Thermo Fisher).

Results and Discussion

Evaluation of Isotope Effects. A reliable approach in quantitative methods is to use stable isotope labeling for ratioing. The use of an isotope-encoded footprinter normalizes the MS intensity of analytes to their isotopic analogs and, therefore, effectively compensates for solvent effects, variable ion suppression from other coeluting analytes, and experimental variations accompanying sample preparation, injection, and instrument setup. To develop an isotope-encoded footprinter, certain criteria are required: 1) isotope effects should not significantly affect the kinetics of footprinting (i.e., the reaction rates of a footprinting reaction are nearly constant when one or more atoms of the footprinter are replaced by isotopes. 2) Analytes and their isotopic analogs (nearly) coelute during the LC; in this way, a pair of isotope-labeled analytes would experience the same ionization efficacy during MS analysis, which is important for


accurate quantification. 3) A sufficient mass difference between two isotopically encoded adducts is needed to separate the peaks of the “light and heavy” analogs over a range of from m/z 400 to 2000. Unfortunately, only a limited number of isotopic footprinters are commercially available. Therefore, we prepared both the “light” and “heavy” reagents (see Experimental) that we used in this footprinting.

We first measured the relative labeling efficiency of D and E on protected-aspartic acid and glutamic acid. They show very similar modification extent ($25.1\% \pm 0.4\%$ for protected D v.s. $25.0\% \pm 0.1\%$ for protected E). (Figure. S4)

To test whether the “intrinsic” chemical reactivity is the same for “light” vs. “heavy”, we chose a cyclic peptide cyclo(RGD^{FE}), which has no secondary structure and no complicating N- or C-termini. The EICs (Figure 1a) for unmodified (black), BHD- (light blue), and d_5 -BHD (dark blue) labeled cyclic peptides were integrated, demonstrating that ratio of modified vs. unmodified was nearly the same, $43.2 \pm 1.6\%$ “light” and $44.6 \pm 1.9\%$ “heavy” labeling. Thus, the presence of isotopes in BHD did not affect the reactivity of the carboxylic residues in cyclo(RGD^{FE}).


This cyclic peptide also helps determine whether isotopomer mixture of BHD can label both Asp and Glu if the sibling residues are adjoining. Metal ions (e.g., Ca, Mg) generally have multiple coordinated sites, and usually include several Asp and Glu in a single binding pocket. It is important that the footprinter can modify all carboxylic residues in a pocket-like site rather than react with one and be “silent” with the other. BHD labels both Asp and Glu even when they are adjoining in the cyclic peptide (cRGD^{FE}). To validate the peptide assignments, we manually assigned peaks in the product-ion spectra (MS/MS) for the BHD- and d_5 -BHD modified cyclo(RGD^{FE}), confirming that the modification occurs at both Asp and Glu (Figure 1b and 1c)³⁵. In addition, we did not find di-labeled peptide.

We next determined whether there are any isotope effects in the separation of BHD- and d_5 -BHD modified peptides. If the light and heavy species coelute, matrix effects should not affect the relative efficiency of ionization of the analytes. Because there is a difference in the size of the atoms^{36,37}, substitution of hydrogen with deuterium could lead to differences in the binding interactions (van der Waals interactions) with the stationary phase. Protiated compounds bind to hydrophobic stationary phase more strongly than their deuterated counterparts.³⁶ When comparing the retention time of BHD- and d_5 -BHD modified peptide (Figure 1a), we found a small isotope effect that d_5 -BHD modified peptides eluted approximately 11 s after the BHD-modified counterpart.



We evaluated the BHD reactivity with CaM by fixing the concentration of EDC at 500 μ M, 50 fold excess relative to CaM (10 μ M). Because each CaM molecule contains 37 Asp and Glu, all Asp and Glu on the protein surface have a chance to react. With increasing amounts of BHD from 5-200 mM, the modification extent of CaM increased from 43% to 65% (Figure S5). When the concentration of BHD was increased further, the modification extent did not significantly change, probably because EDC is no longer the limiting reagent. Thus, we chose 200 mM as the optimum concentration of BHD. We also tested the relative reactivity of “light” and “heavy” BHD to CaM. Similar to cyclo(RGDFE), we found that “light” and “heavy” BHD show nearly identical modification extents for the intact CaM protein (Figure 2).

is more reactive when compared to GEE (Figure 2, compare b to c), $67 \pm 3\%$ BHD and $25 \pm 3\%$ GEE labeling.



BHD as a footprinting reagent. Our next goal is to test BHD as a footprinter to establish whether its reactions report the metal-binding site for CaM. Isotope-encoding provided a “light” and “heavy” labeling of the two different states of CaM. In this protocol (Figure S6), Ca-free CaM was labeled by d_5 -BHD in one Eppendorf tube and Ca-bound CaM was




Figure 3. a) EIC for BHD- (light blue) and d_5 -BHD- (dark blue) modified peptide 78-86, b) product-ion (MS/MS) spectrum for peptide 78-86: unique mass shift of b) 123.0845 Da, and c) 118.0531 Da is consistent with d_5 -BHD modified and d_0 -BHD modification on E84.

labeled by d_0 -BHD in another tube. Following the footprinting, the tube contents were combined in a 1:1 ratio prior to trypsin digestion and LC/MS/MS analysis. In this way, we could maintain all experimental conditions (e.g., protein digestion, MS analysis) nearly identical for both light and heavy samples following protein modification.

Data processing showed 100% sequencing coverage of the tryptic peptides. We took advantage of accurate-mass measurements to obtain reliable EICs for integrating the signal intensities representing (Figure 3a) modifications by d_5 -BHD and d_0 -BHD (seen as two distinct masses, +123.0845 ($C_7H_5N_2$) and +118.0531 ($C_7H_6N_2$) on peptide 78-86, respectively (Figure 3b, 3c)). When comparing their retention time, however, we did observe, as before, a small isotope effect (Figure 3a) that d_5 -BHD modified peptide 78-86 eluted 15.6 s earlier than its d_0 -BHD-modified counterpart. This deviation is consistent with the previously reported effect of deuteration.^{36,37} In addition, we found that the labeled peptides are more hydrophobic and eluted approximately 14 min later than their unmodified counterparts (Figure S7). This chromatographic pattern assists the identification of the modified peptides, sometimes facilitating separation of isomeric peptides with modifications on different residues.

Overall, we found that eight regions of the protein undergo large conformational changes with modification extent difference over 50% and p -values less than 0.01.⁴⁰ These regions are represented by peptides 14-21, 22-30, 31-37, 38-74, 76-86, 78-86, 95-106, and 127-148 (Figure 4), consistent with Ca^{2+} -induced regional structural transitions that dramatically alter the molecular surface of CaM. For example, there is only trace amount of BHD-labeled peptides 22-30, and 38-74 for the holo state, whereas modification extents of these two peptides for the apo-state is > 8% (Figure 4, Table S1).

To establish that these MS results are structurally meaningful, we analyzed them with respect to the high-resolution NMR and X-ray

structures. CaM binds Ca^{2+} at four, nonidentical sites that contain the EF-hand structural motif, two in the N-domain (EF-1 and EF-2) and two in the C-domain (EF-3 and EF-4). Each hand contains an acidic, Ca^{2+} -coordinating loop, or "EF-loop" where Ca^{2+} is chelated with seven oxygens from six residues in a pentagonal bipyramidal fashion.^{6,41} EF-1 chelates Ca^{2+} with ligands D20, D22, D24, and E31 from peptide 14-21, 22-30, 31-37. (Figure 5a and 5f). For this binding site, the reagent is extraordinarily discriminating such that the carboxylates of this region undergo little modification when in the holo state (peptide 14-21, 22-30, 31-37).




Figure 4. Comparison of the modification extents of constituent peptides upon footprinting Ca^{2+} -free CaM (dark blue) and Ca^{2+} -bound CaM (light blue).




Figure 5. 3D structure of Ca^{2+} -free CaM from the average NMR structure (PDB ID, 1CFD) and Ca^{2+} -bound CaM from the X-ray crystal structure (PDB ID, 1CLL), Asp and Glu shown as sticks. a) and f) region 20-31 in Ca^{2+} -free CaM and Ca^{2+} -bound CaM, b) and g) region 56-67 in Ca^{2+} -free CaM and Ca^{2+} -bound CaM, c) and h) region 78-86 in Ca^{2+} -free CaM and Ca^{2+} -bound CaM, d) and i) region 93-104 in Ca^{2+} -free CaM and Ca^{2+} -bound CaM, e) and j) region 129-140 in Ca^{2+} -free CaM and Ca^{2+} -bound CaM. (Gray sphere is the metal ion.)

Likewise, EF-2 region, which is represented by peptide 38-74 and chelates Ca^{2+} via D56, D58, D64 and E67 (Figure 5b and 5g), also becomes unreactive in the holo state. Because Asp and Glu lose nearly completely their chemical reactivity, the modification extent difference for EF hand II reaches 97%.

As pointed out earlier, the binding of Ca^{2+} to the EF hand induces a dramatic change of the linker region (peptide 76-86 and 78-86), changing from a flexible loop (Figure 5c) to a rigid α -helix (Figure 5h). Although the conformation of this linker region was under debate that the formation of the α -helical conformation is a crystallization artifact,⁴² many studies⁴³ report that the Ca^{2+} binding does induce domain opening in CaM, indicating the linker region is extended and becomes more rigid than in the apo-CaM. The plasticity of the linker region in Ca^{2+} -bound CaM allows EF-hand domains to take up various orientations with respect to each other.⁴⁴ Our footprinting results also reveal a significant modification change, evidenced by two overlapping peptides that show approximately 70% modification difference (see Table S1). The D78, D80, E82, E83, and E84 on a flexible loop in holo-CaM can easily react with BHD, whereas when this region becomes a rigid α -helix region in the apo form, those residues are less reactive.

EF-3, as represented by peptide 95-106 (Figure 5d), uses D95 and E104 to chelate Ca^{2+} (Figure 5i). Although the modification ratio for this region in apo CaM is only 0.1%, there is negligible modification for the Ca^{2+} -bound CaM, indicating a large conformational change of EF-3 hand. Finally, for EF-4 peptide 127-148 (Figure 5e) shows a 74% modification decrease upon binding. All the Asp and Glu in this region, except a distant E139, are candidates for binding with Ca^{2+} (Figure 5j).




Figure 6. Comparison of the modification extent of constituent peptides from Mg^{2+} -free CaM (dark blue) and Mg^{2+} -bound CaM (light blue).

To investigate whether changes in SASA agree with our result, we compared the residue-level quantification and the calculated SASA of Ca^{2+} -free and Ca^{2+} -bound CaM (Figure S8). The modification extent of all D and E residues at the EF-hands significantly decreases upon Ca^{2+} binding. The results correlate well with the similar trend that most D and E residues of Ca^{2+} -bound CaM at the binding sites dramatically lose their SASAs. Nevertheless, the decreased modification extent of E82/E83/E84 does not correlate well with their theoretical increasing SASAs. This may be due to the reaction in this region not only depends on solvent accessibility but also on stereochemistry. Compared with previous GEE study,⁴⁵ BHD with 10% amount of GEE labels 27 D and E in CaM, showing higher reactivity. The significant differences of labeling efficiency for GEE and BHD are that the alpha effect in BHD increases the

nucleophilicity of the nitrogen owing to the presence of adjacent nitrogen atoms with nonbonding electron pairs, leading BHD to be more reactive.⁴⁶ In addition, BHD is more discriminating given that it successfully reports four binding-sites and that the linker region undergoes large conformational changes.

Mg²⁺ Binding. In addition to Ca²⁺-binding, we extended the approach to study Mg²⁺ binding with CaM. Like Ca²⁺, divalent Mg is also a “hard” ion and prefers “hard” ligands with oxygen being most preferred. In contrast to the well-characterized Ca²⁺-bound structure, the binding of Mg²⁺ to CaM is still unresolved. Conclusions from microcalorimetric studies indicate four identical Mg²⁺-binding sites that do not completely overlap with the Ca²⁺-binding sites in CaM. On the other hand, conclusions from flow dialysis, NMR, and fluorescence indicate that Mg²⁺ shares same binding sites with Ca²⁺.⁴⁷ Only recently, the structure of Mg²⁺-bound CaM⁶ was partially resolved by X-ray crystallography to show the N-terminal Mg²⁺ binding sites are similar to those of Ca²⁺. Nevertheless, the full high-resolution structure of Mg²⁺-bound CaM remains unknown.

Although previous reports disagree on details, they do indicate that Mg²⁺ indeed binds to CaM, likely at the calcium sites. When we compared our footprinting results for Mg²⁺-bound with those of Ca²⁺-bound, we found that eight peptides regions have modification extent difference more than 0.5 and *p*-value less than 0.01 (Figure 6 and Table S2). Those eight peptides regions belong to EF-1 to 4 plus to the linker region. Our results support the conclusion that the binding site of Mg²⁺ resembles that of Ca²⁺.^{5,41,47}

Comparing the footprinting results for EF-1 and the X-ray structure, we found that although D20, D22, and D24 chelate Mg²⁺, E31 is too far from Mg²⁺ to contribute to the coordination. (Figure 7a) In contrast to pentagonal bipyramid coordination for Ca²⁺, octahedrally coordinated Mg²⁺ does not directly bind to Glu12 of the EF-hand. Footprinting results shows that E31 undergoes 0.008% BHD modification in Mg²⁺-bound CaM, but undetectable modification in Ca²⁺-bound CaM, correlating well with the coordination geometry. Furthermore, the modification-extent differences for Mg²⁺ are significant, but not as significant as for Ca²⁺-binding (89% for Mg²⁺- vs. ~100% for Ca²⁺- bound ones), suggesting weaker binding for Mg²⁺. This difference is consistent with the enthalpy changes upon Ca²⁺ binding being exothermic, whereas endothermic, and thus unfavorable, for Mg²⁺ binding to CaM. Consequently, Mg²⁺ does not bind to CaM as tightly as Ca²⁺.⁴⁸

For EF-2 regions, however, Mg²⁺ shows a very different pattern than Ca²⁺ for binding with CaM. From the X-ray structure, Mg²⁺ coordinates with two oxygens from T62 and a backbone carbonyl group. (Figure 7b) The remaining four coordinating positions are occupied instead, by four water molecules that form strong hydrogen bonds with D58 and E67, allowing Asp and Glu residues to gain some protection. In other words, Mg²⁺ directly binds to first-shell water molecules, allowing it to bind in an indirect way to D58 and E67 (the binding is called a second-shell ligand interaction). Our MS results show no modification extent difference in peptide 38-74 with that in Ca²⁺-CaM even though there is an indirect interaction between Mg²⁺ and the nearby carboxylic residues. This identical footprinting, however, also reminds us that a more sensitive reagent is needed if we want to resolve the pattern between first-shell and second-shell ligand binding.




Figure 7. 3D structure of Mg²⁺-bound CaM (PDB ID, 3UCW). a) Region 20-31, b) region 55-67. (Gray sphere is the metal ion; red sphere is water molecule.)

Furthermore, the linker region of Mg²⁺-bound CaM does show significant change, but the changes are not as large as those of Ca²⁺-bound CaM. (Figure 6, peptide 76-86 and 78-86). The modification extent differences of Mg²⁺-bound CaM are on the borderline (57%) whereas the Ca²⁺-bound shows differences at 70%. One expects that Ca²⁺-bound CaM should undergo larger conformation changes in this region on the basis of previous studies.^{5,43,49}

Finally, we can evaluate EF-3 and EF-4 regions even though there is no crystal structure for this part in Mg²⁺-bound CaM. Like for Ca²⁺-bound CaM, we detected trace amount of modification for peptide 95-106 (EF-3). Peptide 127-148, containing EF-4, undergoes relatively less conformational change with Mg²⁺ compared to Ca²⁺-binding, with modification extent differences 58% vs. 74% for Ca²⁺-binding. The ionic radius of Mg²⁺⁵⁰ is too small to allow some carboxylic acid residues to coordinate with Mg²⁺, eliminating any large differences in the reaction of BHD with Mg²⁺ binding.

In summary, peptide 14-21, 22-30, 31-37, 76-86, 78-86, and 127-148 of Mg²⁺-bound CaM experience conformational changes, but they are less significant than those of Ca²⁺-bound CaM. This indicates that Mg²⁺ with smaller ionic radii binds to CaM without chelating with the 12th residues of the EF-hand, consequently causing smaller structural changes.

Conclusion

With this new footprinting reagent, we can readily identify the four Ca binding sites in CaM and correlate them with 3-D structures. In contrast to other irreversible labeling strategies that either monitor the backbone or side chains around the metal-binding site, this reaction directly footprints carboxyl groups at the binding site. For Mg²⁺ binding, which is weaker and more difficult to discriminate, the footprinting still shows that Mg²⁺ binds with four EF hands similar to Ca²⁺.

This approach has several advantages and some disadvantages: 1) BHD coupled with EDC enables improved chemical reactivity and a stable modified product. 2) The isotope-encoded benzylic acid used for *d*₅-BHD synthesis is commercially available and inexpensive; high-yield *d*₅-BHD can be obtained conveniently with a two-step synthesis. 3) The *d*₅-BHD, incorporating five deuterium atoms, improves the resolution and accuracy of quantification, especially for large proteolytic peptides. On the other hand, there is a small isotope effect on LC elution when a long chromatographic gradient was used for our custom-made C18 column. Further, the reagent is unable to

differentiate first- and second-shell ligand interactions, and a more discriminating labeling protocol or reagent are needed. For the future, we wish to extend this new footprinting approach to other hard metal-ion binding and, via a titration protocol, to test its ability to determine metal-binding affinity.

ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge on the ACS Publications website.

Synthetic route and MS spectra for BHD and *d*₅-BHD; modification reaction by BHD coupling reaction; condition screening for BHD; workflow of isotope-encoded covalent labeling; EIC for unmodified peptide 78-86; modification extent and calculated SASA of D and E residues at the EF-hand; summary of carboxylic acid modification percentage for Ca²⁺-free/Ca²⁺-bound CaM and Mg²⁺-free/Mg²⁺-bound CaM.

AUTHOR INFORMATION

Corresponding Author

*E-mail: mgross@wustl.edu. Tel: +1-314-935-4814. Fax: +1-314-935-7484.

Notes

Any additional relevant notes should be placed here.

ACKNOWLEDGMENT

We acknowledge research supported by grants NIH P41RR103422, as well as Dr. Gerstenecker from Washington University in St. Louis for his advice for this project and Protein Metrics for software.

REFERENCES

- (1) Dudev, T.; Lim, C., Competition among metal ions for protein binding sites: Determinants of metal ion selectivity in proteins. *Chem. Rev.* **2014**, 114, 538-556.
- (2) Tainer, J. A.; Roberts, V. A.; Getzoff, E. D., Protein metal-binding sites. *Curr. Opin. Chem. Biol.* **1992**, 3, 378-387.
- (3) Bowman, S. J.; Bridwell-Rabb J. Drennan, C. L. Metalloprotein crystallography: More than a structure., *Acc. Chem. Res.* **2016**, 49, 695-702.
- (4) Gregory, D. S.; Martin, A. C.; Cheetham, J. C.; Rees, A. R., The prediction and characterization of metal binding sites in proteins. *Protein Eng.* **1993**, 6, 29-35.
- (5) Katz, A. K.; Glusker J. P.; Beebe, S. A.; Bock, C. W., Calcium ion coordination: A comparison with that of beryllium, magnesium, and zinc. *J. Am. Chem. Soc.* **1996**, 118, 5752-5763.
- (6) Senguen, F. T.; Grabarek, Z., X-ray structures of magnesium and manganese complexes with the N-terminal domain of calmodulin: Insights into the mechanism and specificity of metal ion binding to an EF-hand. *Biochemistry*, **2012**, 51, 6182-6194.
- (7) Denessiouk, K.; Permyakov, S.; Denesyuk, A.; Permyakov, E.; Johnson, M. S., Two structural motifs within canonical EF-hand calcium-binding domains identify five different classes of calcium buffers and sensors. *PLoS One* **2014**, 9, e109287.
- (8) Dudev, T.; Chang, L.; Lim, C., Factors governing the substitution of La³⁺ for Ca²⁺ and Mg²⁺ in metalloproteins: A DFT/CDM study. *J. Am. Chem. Soc.* **2005**, 127, 4091-4103.
- (9) Payandeh, J.; Pföh, R.; Pai, E. F., The structure and regulation of magnesium selective ion channels. *Biochim. Biophys. Acta* **2013**, 1828, 2778-2792.
- (10) Jesnsen, M.R.; Petersen, G.; Lauritzen, C.; Pedersen, J.; Led, J. J., Metal binding sites in proteins: Identification and characterization by paramagnetic NMR relaxation. *Biochemistry* **2005**, 44, 11014-11023.
- (11) Zhang, Q.; Dai, X.; Cong, Y.; Zhang J.; Chen, D.; Dougherty, M. T.; Wang, J.; Ludtke, S. J.; Schmid, M. F.; Chiu, W., Cryo-EM structure of a molluscan hemocyanin suggests its allosteric mechanism. *Structure*, **2013**, 21, 604-613.
- (12) Kojetin, D. J.; Matta-Camacho, E.; Hughes, T. S.; Srinivasan, S.; Nwachukwu, J. C.; Cavvet, V.; Nowak, J.; Chalmers, M. J.; Marciano, D. P.; Kamenecka, T. M.; Shulman, A. I.; Rance, M.; Griffin P. R.; Bruning, J. B.; Nettles, K. W., Structural mechanism for signal transduction in RXR nuclear receptor heterodimers. *Nat. Commun.* **2015**, 6, 8013.
- (13) Fast, C. S.; Vahidi, S.; Konermann, L., Changes in enzyme structural dynamics studied by hydrogen exchange-mass spectrometry: Ligand binding effects or catalytically relevant motions? *Anal. Chem.* **2017**, 89, 13326-13333.
- (14) Jensen P. F.; Comamala, G.; Trelle, M. B.; Madsen, J. B.; Jørgensen, T. J. D.; Rand, K. D., Removal of N-linked glycosylations at acidic pH by PNGase A facilitates hydrogen/deuterium exchange mass spectrometry analysis of N-linked glycoproteins. *Anal. Chem.* **2016**, 88, 12479-12488.
- (15) Wang, L.; Chance, M. R., Protein footprinting comes of age: Mass spectrometry for biophysical structure assessment. *Mol. Cell. Proteomics* **2017**, 16, 706-716.
- (16) Zhu, M.; Rampel, D. L.; Du, Z.; Gross, M. L., Quantification of protein-ligand interactions by mass spectrometry, titration, and H/D exchange: PLIMSTEX. *J. Am. Chem. Soc.* **2003**, 125, 5252-5253.
- (17) Zhang, B.; Cheng, M.; Rampel, D.; Gross, M. L., Implementing fast photochemical oxidation of proteins (FPOP) as a footprinting approach to solve diverse problems in structural biology. *Methods* **2018**, 144, 94-103.
- (18) Zhang, B.; Rempel, D. L.; Gross, M. L., Protein footprinting by carbenes on a fast photochemical oxidation of proteins (FPOP) platform. *J. Am. Soc. Mass Spectrom.* **2016**, 27, 552-555.
- (19) Jumper, C. C.; Schriemer, D. C., Mass spectrometry of laser-initiated carbene reactions for protein topographic analysis. *Anal. Chem.* **2011**, 83, 2913-2920.
- (20) Chen, J.; Cui, W.; Gross, M. L., New protein footprinting: Fast photochemical iodination combined with top-down and bottom-up mass spectrometry. *J. Am. Soc. Mass Spectrom.* **2012**, 23, 1306-1318.
- (21) Cheng, M.; Zhang, B.; Cui, W.; Gross, M. L., Laser-initiated radical trifluoromethylation of peptides and proteins: Application to

mass-spectrometry-based protein footprinting. *Angew. Chem. Int. Ed.* **2017**, 56, 14007-14010.

(22) Hoare, D. G.; Koshland Jr., D. E., A method for the quantitative modification and estimation of carboxylic acid groups in proteins. *J. Biol. Chem.* **1967**, 242, 2447-2453.

(23) Underbakke, E. S.; Zhu, Y.; Kiessling, L. L., Isotope-coded affinity tags with tunable reactivities for protein footprinting. *Angew. Chem. Int. Ed.* **2008**, 47, 9677-9680.

(24) Borotto, N. B.; Zhou, Y. P.; Hollingsworth, S. R.; Hale, J. E.; Graban, E. M.; Vaughan, R. C.; Vachet, R. W., Investigating therapeutic protein structure with diethylpyrocarbonate labeling and mass spectrometry. *Anal. Chem.* **2015**, 87, 10627-10634.

(25) Wen, J.; Zhang, H.; Gross, M. L.; Blankenship, R. E., Membrane orientation of the FMO antenna protein from *Chlorobaculum tepidum* as determined by mass spectrometry-based footprinting. *Proc. Natl. Acad. Sci. USA* **2009**, 106, 6134-6139.

(26) Weckslar, A. T.; Kalo, M. S.; Deperalta, G., Mapping of Fab-1:VEGF interface using carboxyl group footprinting mass spectrometry. *J. Am. Soc. Mass Spectrom.* **2015**, 26, 2077-2080.

(27) Zhang, H.; Shen, W.; Rempel, D.; Monsey, J.; Vidasky, I.; Gross, M. L.; Bose, R., Carboxyl-group footprinting maps the dimerization interface and phosphorylation-induced conformational changes of a membrane-associated tyrosine kinase. *Mol. Cell. Proteomics.* **2011**, 10, M110.005678.

(28) Li, K.; Chen, G.; Mo, J.; Huang, R.; Deyanova, E. G.; Beno, B. R.; O'Neil, S. R.; Tymiak, A. A.; Gross, M. L., Orthogonal mass spectrometry-based footprinting for epitope mapping and structural characterization: The IL-6 receptor upon binding of protein therapeutics. *Anal. Chem.* **2017**, 89, 7742-7749.

(29) Gau, B.; Garai, K.; Frieden, C.; Gross, M. L., Mass spectrometry-based protein footprinting characterizes the structures of oligomeric apolipoprotein E2, E3, and E4. *Biochemistry* **2011**, 50, 8117-8216.

(30) Kaur, P.; Tomechko S. E.; Kiselar, J.; Shi, W.; Deperalta, G.; Weckslar, A. T.; Gokulrangan, G.; Ling, V.; Chance, M. R., Characterizing monoclonal antibody structure by carboxyl group footprinting. *mAbs* **2015**, 7, 540-552.

(31) Lin, M.; Krawitz, D.; Callahan, M. D.; Deperalta, G.; Weckslar, A. T., Characterization of ELISA antibody-antigen interaction using footprinting-mass spectrometry and negative staining transmission electron microscopy. *J. Am. Soc. Mass Spectrom.* **2018**, 29, 961-971.

(32) Leitner, A.; Joachimiak, L. A.; Unverdorben, P.; Walzthoeni, T.; Frydmam, J.; Förster, F.; Aebersold, R., Chemical cross-linking/mass spectrometry targeting acidic residues in proteins and protein complexes. *Proc. Natl. Acad. Sci. USA* **2014**, 111, 9455-9460.

(33) Kahsai, A. W.; Rajagopal, S.; Sun, J.; Xiao, K., Monitoring protein conformational changes and dynamics using stable-isotope labeling and mass spectrometry. *Nat. Protoc.* **2014**, 9, 1301-1319.

(34) Li, Z.; Bai, X.; Deng, Q.; Zhang, G.; Zhou, L.; Liu, Y.; Wang, J., Wang, Y., Preliminary SAR and biological evaluation of antitubercular triazolothiadiazine derivatives against drug-susceptible and drug-resistant Mtb strains. *Bioorganic Med. Chem.* **2017**, 25, 213-220.

(35) Liu, W.; Ng, J.; Meluzzi, D.; Banderia, N.; Gutierrez, M.; Simmons, T. L.; Schultz, A. W.; Linington, R. G.; Moore, B. G.;

Gerwick, W. H.; Pevzner, P. A.; Dorrestein, P. C., Interpretation of tandem mass spectra obtained from cyclic nonribosomal peptides. *Anal. Chem.* **2009**, 81, 4200-4209.

(36) Turowski, M.; Yamakawa, N.; Meller, J. Kimata, K.; Ikegami, T.; Hosoya, K.; Tanaka, N.; Thornton, E. R., Deuterium isotope effects on hydrophobic interactions: The importance of dispersion interactions in the hydrophobic phase. *J. Am. Chem. Soc.*, **2003**, 125, 13836-13849.

(37) Guo, K.; Ji, C.; Li, L., Stable-isotope dimethylation labeling combined with LC-ESI MS for quantification of amine-containing metabolites in biological samples. *Anal. Chem.*, **2007**, 79, 8631-8638.

(38) Ouyang, H.; Vogel, H. J., Metal ion binding to calmodulin: NMR and fluorescence studies. *Bio. Metals* **1998**, 11, 213-222.

(39) Ngoka, L. C.; Gross, M. L., A nomenclature system for labeling cyclic peptide fragments. *J. Am. Soc. Mass Spectrom.* **1999**, 10, 360-363.

(40) McCarthy, D. J.; Smyth, G. K., Testing significance relative to a fold-change threshold is a TREAT. *Bioinformatics* **2009**, 25, 765-771.

(41) Ohashi, W.; Hirota, H.; Yamazaki, T., Solution structure and fluctuation of the Mg²⁺-bound form of calmodulin C-terminal domain. *Protein Sci.* **2011**, 20, 690-701.

(42) Vetter, S. W.; Leclerc, E., Novel aspects of calmodulin target recognition and activation. *Eur. J. Biochem.* **2003**, 270, 404-414.

(43) Grabarek, Z., Insights into modulation of calcium signaling by magnesium in calmodulin, troponin C and related EF-hand proteins. *Biochim. Biophys. Acta.* **2011**, 1813, 913-921.

(44) Gifford, J. L.; Walsh, M. P.; Vogel, H. J., Structures and metal-ion-binding properties of the Ca²⁺-binding helix-loop-helix EF-hand motifs. *Biochem. J.* **2007**, 405, 199-221.

(45) Zhang, H.; Wen, J.; Huang, R.; Blankenship, R. E.; Gross, M. L., Mass spectrometry-based carboxyl footprinting of proteins: Method evaluation. *Int. J. Mass Spectrom.* **2012**, 312, 78-86.

(46) Singh, N.; Karpichev, Y.; Sharma, R.; Gupta, B.; Sahu, A. K.; Satnami, M.; Ghosh, K. K., From α -nucleophiles to functionalized aggregates: exploring the reactivity of hydroxamate ion towards esterolytic reactions in micelles. *Org. Biomol. Chem.* **2015**, 13, 2827-2848.

(47) Ohki, S.; Ikura, M.; Zhang, M., Identification of Mg²⁺-binding sites and the role of Mg²⁺ on target recognition by calmodulin. *Biochem.* **1997**, 36, 4309-4316.

(48) Gilli, R.; Lafitte, D.; Lopez, C.; Kihoffer, M.; Makarov, A.; Briand, C.; Haiech, J., Thermodynamic analysis of calcium and magnesium binding to calmodulin. *Biochemistry*, **1998**, 37, 5450-5456.

(49) Cates, M. S., Teodoro, M. L., Phillips, G. N., Molecular mechanisms of calcium and magnesium binding to parvalbumin. *Biophys. J.* **2002**, 82, 1133-1146.

(50) Shannon, R. D., Revised effective ionic radii and systematic studies of interatomic distances in halides and chalcogenides. *Acta Crystallogr.* **1976**, A32, 751-767.

