An iodoetherification-dehydroiodination strategy for the synthesis of complex spiroketals from dihydroxyalkene precursors†

K. A. Tony, Xiaohua Li, Darrin Dabideen, Jialiang Li and David R. Mootoo*

Received 13th December 2007, Accepted 13th February 2008 First published as an Advance Article on the web 22nd February 2008 DOI: 10.1039/b719247a

Dihydroxyalkenes or their monoprotected alcohol derivatives are transformed to 5,5- and 5,6-spiroketals through a sequence involving an initial iodocyclization, followed by a silver triflate mediated spiroketalization step on the derived hydroxy-iodoether.

Introduction

Spiroketal subunits comprise the structures of several groups of biologically interesting natural products.1 Their conformational and geometric characteristics also make them attractive as scaffolds in diversity-oriented synthesis.² Accordingly, spiroketal synthesis has attracted considerable attention. Classically, spiroketals have been synthesized through acid-catalyzed cyclization of ketodiol precursors.^{1a,b} Several elegant methodologies have been devised to address structural complexity.³ Methods that employ straightforward segment coupling reactions and mild conditions for spiroketalization are particularly appealing. In this vein, we envisaged an approach in which a dihydroxyalkene or a partially protected derivative 1 is transformed to a spiroketal 4 (Scheme 1). Iodoetherification of 1 could lead to iodinated cyclic ethers like 2.⁴ Dehydroiodination of 2 provides a highly reactive enol ether 3, which would undergo spiroketalization under mildly acidic conditions. In essence, the alkene moiety in 1 is regioselectively functionalized to an acetal, and acts as an alkyne synthon. The strategy may be therefore complementary to the metal-promoted spiroketalization of dihydroxyalkynes (e.g. 5).⁵ Importantly, since precursors like 1 are obtainable via straightforward olefination reactions, this methodology allows for a highly convergent synthesis



of complex spiroketals. We have reported a preliminary application of this strategy to the ABCD bis-spiroketal system of azaspiracid 1.⁶ Herein, we describe a more detailed investigation into the scope of this method.

Results and discussion

Except for **29**, the dihydroxyalkene derivatives were prepared *via* an olefin cross-metathesis (CM) using an excess of one of the olefin partners (Table 1).⁷ The metathesis reactions were performed on various highly oxygenated alkene partners and afforded yields of the CM product ranging from 65–85%.⁸ For yield optimization, easier purification of the CM product or the requirement for a partially protected dihydroxyalkene, it was sometimes more practical to perform the CM with protection of the alcohol groups in one or both of the reaction partners. The diene precursor **29** was assembled through the Wittig olefination of known aldehyde **11**⁹ and phosphonium salt **18**.

Spiroketal precursors 19, 21, 23 and 24 were designed to get a preliminary evaluation of the feasibility with respect to different ring sizes (Table 2). Based on previous investigations¹⁰ on the relative rates of iodoetherification of hydroxyalkenes, we argued that dihydroxyalkenes like 19 and 21 or 23 should favor the 5-exo-trig and the 6-exo-trig pathways, respectively, over other modes of cyclization. Therefore, it should be possible to use substrates 19, and 21 or 23 in which the eventual alcohols of the spiroketal are unprotected, as precursors to 5,5- and 5,6-spiroketal frameworks, respectively. Thus, treatment of 19, 21 and 23 with IDCP (iodonium dicollidine perchlorate) in dichloromethane provided a mixture of cyclization products in 85, 85 and 70% yield respectively. However, the NMR data for these structures did not allow for distinction between the possible regioisomeric cyclization products. These structures were tentatively assigned as 31, 32 and 33 based on the aforementioned regioselectivity considerations. Exposure of the individual product mixtures to silver triflate in dichloromethane in the presence of collidine led to 5,5-spiroketal 43a,b in 70% yield, and the 5,6 frameworks 44a,b and 45a,b in 74 and 55% yield respectively. The mixture 43a,b was chromatographically inseparable and this made stereochemical assignment of the spiroketal configuration in individual components impossible. However, mixtures 44a,b and 45a,b were separable, and their stereochemistry was assigned by 2D COSY and NOESY NMR (see ESI[†]). It should be noted that while the gross structures of spiroketals 43a,b, 44a,b and 45a,b support the structures assigned to 31, 32 and 33 respectively, the possibility that the corresponding regioisomer resulting from the 5-endo-trig pathway could also lead to the identical spiroketal products, makes structures 31, 32 and 33 ambiguous. In the event, the isolation of

Department of Chemistry, Hunter College/CUNY, New York, NY, 10021, USA. E-mail: dmootoo@hunter.cuny.edu; Fax: +1 212-772-5332; Tel: +1 212-772-4356

[†] Electronic supplementary information (ESI) available: Experimental procedures, physical data and NMR spectra for selected new compounds. See DOI: 10.1039/b719247a

Table 1 Synthesis of dihydroxyalkene derivatives and 23



^{*a*} Except for **29**, the dihydroxyalkene derivatives were prepared by CM with 10 mol% Grubbs 2nd generation catalyst in CH_2Cl_2 using an excess of one of the olefin partners. For optimal yields, CM's were in certain cases performed on protected alcohol derivatives and the CM product deprotected to give the precursor for the iodoetherification step. The homodimer derived from the alkene that was used in excess was obtained as a side product. Compound **29** was prepared through a Wittig olefination.

a single spiroketal framework starting from dihydroxyalkenes **19** and **21** in reasonable yields suggests that for 5,5- and 5,6-spiroketal systems, selective protection of the alcohols that constitute the eventual acetal residue may not be necessary.

The extension of the strategy to 6,6-spiroketals was next examined using the hydroxyalkene 24 as a test substrate. In this case, the expectation that 5-*exo*-trig would be favored over 6-*exo*-trig or 6-*endo*-trig pathways meant that protection of one

of the two alcohols of the eventual spiroketal would be necessary. Accordingly, **24** was treated with IDCP under the standard conditions, and the product **34** was desilylated to provide **35**, the precursor for the spiroketalization step. However, treatment of **35** under the standard conditions led to adjacently linked THF-THP **47**, and not the desired spiroketal **46**. This result suggested that the iodoetherification step yielded **34** as expected, but the subsequent AgOTf-mediated reaction favored THF rather than



^{*a*} Iodocyclization reactions were performed by treatment of the dihydroxyalkenes or their mono-protected derivatives with IDCP in anhydrous CH_2Cl_2 . ^{*b*} Except for **50**, spiroketals were obtained by exposure of the hydroxy-iodide precursor to a mixture of AgOTf and collidine in anhydrous CH_2Cl_2 . For **50**, the iodo-acetal **40** was treated with AgOTf in wet THF.

spiroketal formation. This result suggests that application of this strategy to 6,6-spiroketals might in general be problematic,

and consequently we focused on evaluating the synthetic scope for 5,5- and 5,6-frameworks.

The results for carbohydrate-derived substrates **25** and **26** illustrate the compatibility in more highly functionalized settings. The mono-acetates **25** and **26** (as opposed to their deacetylated diol derivatives) were initially screened because these were the direct products from the CM reaction. Thus, iodoetherification of **25** and **26**, followed by acetate hydrolysis of the resulting acetoxy-iodoethers, provided the spiroketalization precursors **37** and **39**. Treatment of the latter with AgOTf in the presence of collidine provided the 5,5- and 5,6-spiroketals **48a,b** and **49a,b** in 76 and 65% yield respectively. When the deacetylated diol derivatives of **25** and **26** were subjected to a two-step iodoetherification–spiroketalization sequence, **48a,b** and **49a,b** were produced in similar overall yield and epimer ratio to the material obtained from the three-step iodoetherification–deacetylation–spiroketalization sequence on **25** and **26**.

In order to determine whether the stereochemistry in the iodoether impacts on the stereoselectivity of the AgOTf-mediated spiroketalization reaction, the latter was performed on individual diastereomers of THP-iodide **33** and THF-iodide **37**. In both cases the ratio of spiroketals produced from the individual iodoether diastereomers was very similar to that obtained from the corresponding mixture (that is, *ca.* 2 : 1 for **45a**:**45b** and 1 : 1 for **48a**:**48b** respectively). These results suggested that the stereochemistry of the iodoether precursor is not directly transferred to the spiroketal product.

The conversion of hydroxy-acetal-alkene substrate **27** to the bisspiroketal **50** was next investigated. However, iodoetherification of **27** produced a complex mixture, in part due to unwanted 5*endo*-trig cyclization involving the OH group of the lactol. The methyl acetal **28** was therefore subjected to the iodocyclization procedure and the crude product exposed to AgOTf in wet THF (without added collidine). This sequence afforded a mixture of bis-spiroketals **50a,b** in 52% overall yield from **28**.

The transformation of **30** to **51a,b** illustrates that the chemoselective elaboration of diene substrates may be possible. Thus, IDCP cyclization on **30** followed by removal of the PMB protecting group provided the hydroxy-iodo-dihydropyran **42**, which led to the spiroketal mixture **51a,b**.

The observation that the stereoselectivity of the spiroketalization step is not affected by the stereochemistry of the iodoether precursor is consistent with a mechanism involving the cyclic oxocarbenium ion 55. However, the pathway leading to 55 is more conjectural (Scheme 2). Intermediate 55 could arise from protonation of an initially formed exocyclic enol ether 54, or *via* iodide activation in 53 and cation formation, followed by hydride transfer. That enol ether 54 (or its endocyclic isomer) was not observed in



Scheme 2 Mechanistic analysis.

the crude reaction mixture, even though spiroketalization occurs in the presence of excess collidine, supports the direct formation of 55 from 53. However, the possibility that 54 is formed then rapidly converted to product under the reaction conditions cannot be excluded.

Conclusion

In conclusion, the transformation of dihydroxyalkenes or their partially protected derivatives to highly substituted 5,5- and 5,6spiroketal frameworks has been explored. The compatibility of this strategy with a wide variety of functional groups and the availability of the dihydroxyalkene precursors through straightforward olefination procedures, of which the olefin metathesis is a prominent example, makes this an attractive methodology for the convergent assembly of complex targets. More extensive mechanistic and synthetic investigations are underway and will be reported in due course.

Acknowledgements

This investigation was supported by grants R01 GM57865 from the National Institute of General Medical Sciences of the National Institutes of Health (NIH). A "Research Centers in Minority Institutions" award RR-03037 from the National Center for Research Resources of the NIH, which supports the infrastructure and instrumentation of the Chemistry Department at Hunter College, and MBRS-RISE award GM60665, are also acknowledged.

Notes and references

- (a) F. Perron and K. F. Albizati, *Chem. Rev.*, 1989, **89**, 1617–1661; (b) J. Pietruszka, *Angew. Chem., Int. Ed*, 1998, **37**, 2629–2636; (c) W. Francke and W. Kitching, *Curr. Org. Chem.*, 2001, **5**, 233–251; (d) M. A. Brimble and D. P. Furkert, *Curr. Org. Chem.*, 2003, **7**, 1461–1484; (e) K. T. Mead and B. N. Brewer, *Curr. Org. Chem.*, 2003, **7**, 227–225; J. E. Aho, P. M. Pihko and T. K. Rissa, *Chem. Rev.*, 2005, **105**, 4406–4440.
- 2 (a) R. Haag, A. G. Leach, S. V. Ley, M. Nettekoven and J. Schnaubelt, Synth. Commun., 2001, 31, 2965–2977; (b) B. A. Kulkarni, G. P. Roth, E. Lobkovsky and J. A. Porco, Jr., J. Comb. Chem., 2002, 4, 56–72; (c) O. Barun, S. Sommer and H. Waldeman, Angew. Chem., Int. Ed., 2004, 43, 3195–3199; (d) G. Zinzalla, L.-G. Milroy and S. V. Ley, Org. Biomol. Chem., 2006, 4, 1977–2002.
- 3 Selected recent synthesis of complex spiroketals: (a) J. Hao and C. J. Forsyth, Tetrahedron Lett., 2002, 43, 1-2; (b) M. T. Crimmins, J. D. Katz, D. G. Washburn, S. P. Allwein and L. F. McAtee, J. Am. Chem. Soc., 2002, 124, 5661-5663; (c) E. B. Holson and W. R. Roush, Org. Lett., 2002, 4, 3719-3723; (d) K. I. Tsang, M. A. Brimble and J. B. Bremner, Org. Lett., 2003, 5, 4425-4427; (e) L. Wang and P. E. Floreancig, Org. Lett., 2004, 6, 569-572; (f) J. Liu and R. P. Hsung, Org. Lett., 2005, 7, 2273-2276; (g) A. V. Statsuk, D. Liu and S. Kozmin, J. Am. Chem. Soc., 2004, 126, 9546-9547; (h) L. R. Takaoka, A. J. Buskmelter, T. E. LaCruz and S. D. Rychnovsky, J. Am. Chem. Soc., 2005, 127, 528-529; (i) J. S. Potuzaz, S. B. Moilanen and D. S. Tan, J. Am. Chem. Soc., 2005, 127, 13796-13797; (j) K. C. Nicolaou, M. O. Frederick, G. Petrovic, K. P. Cole and E. Z. Loizidou, Angew. Chem., Int. Ed., 2006, 45, 2609–2615; (k) X.-T. Zhou and R. G. Carter, Angew. Chem., Int. Ed., 2006, 45, 1787-1790; (1) Y. Li, F. Zhou and C. J. Forsyth, Angew. Chem. Int. Ed., 2007, 46, 279–282; (m) D. A. Evans, T. B. Dunn, L. Kvrnø, A. Beauchemin, B. Raymer, E. J. Olhava, J. A. Mulder, M. Juhl, K. Kagechika and D. A. Favor, Angew. Chem., Int. Ed., 2007, 46, 4698-4703.
- 4 G. Cardilo and M. Orena, Tetrahedron, 1990, 46, 3321-3408.
- 5 For a recent example of Au(I) alkyne spiroketalization, see: B. Liu and J. K. De Brabander, *Org. Lett.*, 2006, **8**, 4907–4910.
- 6 X. Li, J. Li and D. R. Mootoo, Org. Lett., 2007, 4303-4306.

- 7 A. K. Chatterjee, T.-L. Choi, D. P. Sanders and R. M. Grubbs, J. Am. Chem. Soc., 2003, **125**, 11360–11370; S. J. Connon and S. Blechert, Angew. Chem., Int. Ed., 2003, **42**, 1900–1923.
- 8 (a) Alkenes 7, 8, 14, 17 and 18, were obtained through known procedures on (S)-malic acid: R. Zemribo and K. T. Mead, *Tetrahedron Lett.*, 1998, 39, 3895–3898; J. A. Marshall and J. J. Sabatini, Org. Lett., 2005, 7, 4819–4822; (b) Alkenes 15 and 16 were prepared from the homoallylic alcohol derived from the Keck allylation of 5-benzyloxypentanal: G. E. Keck, K. H. Tarbet and L. S. Geraci, J. Am. Chem. Soc., 1993, 115, 8467–8468; (c) Alkene 11: S. J.

Danishefsky, M. P. DeNinno, G. B. Phillips, R. E. Zelle and P. A. Lartey, *Tetrahedron*, 1986, **42**, 2809–2819; (*d*) Alkene **12** was prepared *via* modification of the synthesis of a related sugar derivative: M. Terauchi, H. Abe, A. Matsuda and S. Shuto, *Org. Lett.*, 2004, **6**, 3751–3754.

- 9 C. R. Schmid, J. D. Bryant, M. Dowlatzedan, J. L. Philllips, D. E. Prather, R. D. Schantz, N. L. Sear and C. S. Vianco, *J. Org. Chem.*, 1991, **56**, 4056–4058.
- 10 S. B. Bedford, K. E. Bell, F. Bennet, C. J. Haynes, D. W. Knight and D. E. Shaw, *J. Chem. Soc.*, *Perkin Trans.* 1, 1999, 2142–2153.