


A STEREO CONTROLLED ROUTE TO A KEY INTERMEDIATE FOR THE SYNTHESIS OF MAYTANSINE

M. Samson, P. De Clercq, H. De Wilde and M. Vandewalle^{*}
 State University of Ghent, Department of Organic Chemistry,
 Laboratory of Organic Synthesis,
 Krijgslaan, 271 (S.4), B-9000 Gent (Belgium).


(Received in UK 6 July 1977; accepted for publication 14 July 1977)

Recently a number of approaches have been published¹ towards the synthesis of various parts of maytansine (1), a novel type ansa macrolide, which shows significant antitumor activity².


In this paper we wish to describe the stereocontrolled synthesis of the building blocks 2a and 2b, corresponding to the "eastern zone" of the title molecule and containing carbon atoms 5 to 9; both synthons possess the correct stereochemistry at C-6 and C-7 and allow further elaboration at both ends, as shown by their transformation into respectively 7a and 7b, which are potential acylanion equivalents.




1




2 a; $R_1 = \text{CMe}_2\text{Ph}$; $R_2 = \text{SiMe}_2\text{t.Bu}$
2 b; $R_1 = \text{SiMe}_2\text{t.Bu}$; $R_2 = \text{CH}_2\text{SCH}_3$




3 a; $R_1 = \text{CMe}_2\text{Ph}$
3 b; $R_1 = \text{SiMe}_2\text{t.Bu}$




4 a; $R_1 = \text{CMe}_2\text{Ph}$
4 b; $R_1 = \text{SiMe}_2\text{t.Bu}$



5 a; $R_1 = \text{CMe}_2\text{Ph}$; $R_2 = \text{H}$
5 b; $R_1 = \text{SiMe}_2\text{t.Bu}$; $R_2 = \text{H}$
6 a; $R_1 = \text{CMe}_2\text{Ph}$; $R_2 = \text{SiMe}_2\text{t.Bu}$
6 b; $R_1 = \text{SiMe}_2\text{t.Bu}$; $R_2 = \text{CH}_2\text{SCH}_3$



7a



7b

The efficiency of the present approach is based on the easy synthesis of the five carbon unit by oxidative cleavage of functionalised cyclopentenones such as 4a and 4b. Treatment of the readily available 4-cumyloxy-2-cyclopentenone 3a³ with dimethylcopper lithium (1.5 eq, ether, -78°) followed by addition of chlorotrimethylsilane-pyridine (2 eq, -78° and allowed to warm up to r.t.) gave the silylenolether 4a⁴ [$\nu(\text{neat})$: 1650, 1250, 860, 765 and 705 cm^{-1} ; $\delta(\text{CCl}_4)$: 7.38 (5H, m), 4.40 (1H, q, $^3J = 1.8$ Hz; $^4J = -1.8$ Hz), 3.48 (1H, dt, X of ABX, $\Sigma^3J = 17$ Hz), 2.70 (1H, m), 2.35 (2H, dt, AB of ABX, $^4J = -1.8$ Hz), 1.58 (6H, s), 0.88 (3H, d, $^3J = 6.9$ Hz), 0.20 (9H, s)].

Product 4a was directly treated with ozone (1 eq, CH_2Cl_2 , CH_3OH , -78°); subsequent reduction of the ozonide with sodium borohydride and esterification with diazomethane yielded 5a (49 % from 3a). [$\nu(\text{neat})$: 3500, 1745, 1610 and 1500 cm^{-1}]. Protection of the hydroxyl group with *t*-butyldimethylsilylchloride⁵ (1.05 eq, imidazole, 25°, 3 h) afforded the silylether 6a [$\nu(\text{neat})$: 2870, 1745, 1610, 1500, 1250, 840 and 765 cm^{-1} ; $\delta(\text{CCl}_4)$: 7.42 (5H, m), 4.23 (1H, d of X part of ABX, $^3J = 3.0$ Hz, $|J_{AX} + J_{BX}| = 12.6$ Hz), 3.62 and 3.50 (2H, CD of CDY, $^2J = -10.4$ Hz), 3.58 (3H, s), 2.43 and 2.30 (2H, AB of ABX, $^2J = -15.3$ Hz, $^3J = 7.2$ Hz and 5.4 Hz), 1.69 (1H, m), 1.65 (3H, s), 1.58 (3H, s), 0.99 (9H, s), 0.94 (3H, d, $^3J = 6.6$ Hz), 0.12 (6H, s)]: Treatment with diisobutylaluminium hydride (1 eq, -78°, toluene) yielded the aldehyde 2a [$\nu(\text{neat})$: 1730 cm^{-1} ; $\delta(\text{CCl}_4)$: 9.57 (t, $^3J = 1.1$ Hz)]. The conversion of the aldehyde function to a potential acylanion equivalent was effected by treatment with trimethylorthothioborate under neutral conditions⁶, yielding 7a (80 % yield). [$\delta(\text{CCl}_4)$: 7.45 (5H, m), 4.01 (1H, dt, X of ABX, $\Sigma^3J = 15.0$ Hz), 3.60 and 3.45 (2H, CD of CDY, $^2J = -10.2$ Hz), 3.30 (1H, t, $^3J = 7.2$ Hz), 1.96 (6H, s), 1.65 and 1.58 (ss, 6H), 0.96 (9H, s), 0.91 (3H, d, $^3J = 6.9$ Hz), 0.12 (6H, s)].

In a similar way 4-*t*-butyldimethylsilyloxy-2-cyclopentenone⁷ (3b) led to the alcohol 5b, which was protected as a methylthiomethoxymethyl ether⁸ (DMSO, HOAc, NaOAc, 24 hr, r.t.) yielding (45 % from 3b) the ester 6b [$\nu(\text{neat})$: 1745, 1250, 835 and 770 cm^{-1} ; $\delta(\text{CDCl}_3)$: 4.73 and 4.68 (2H, $^2J = -11.7$ Hz), 4.35 (1H, dt, $^3J = 3.3$ Hz and 6.6 Hz), 3.82 (3H, s), 3.61 and 3.48 (2H, AB of ABX, $^2J = -9$ Hz, $^3J = 6.6$ Hz and 6.6 Hz), 2.57 (2H, d, $^3J = 6.6$ Hz), 2.24 (3H, s), 1.98 (1H, d of sextuplet,

X of ABX, $^3J = 6.6$ Hz and 3.3 Hz, 1.02 (3H, d, $^3J = 6.6$ Hz), 0.97 (9H, s), 0.18 (6H, s)]. Reduction with diisobutylaluminium hydride (1 eq, -78° , toluene) gave quantitatively the aldehyde 2b [$\nu(\text{neat})$: 2720 , 1720 , 1250 , 835 and 770 cm^{-1} ; $\delta(\text{CCl}_4)$: 9.91 (1H, t, $^3J = 2.1$ Hz), 4.70 and 4.65 (2H, $^2J = -11.7$ Hz), 4.40 (1H, dt, X of ABX, $^3J = 15.9$ Hz), 3.53 (2H, d, $^3J = 6$ Hz), 2.66 (2H, AB of ABX), 2.23 (3H, s), 1.97 (1H, m), 1.05 (3H, d, $^3J = 6.6$ Hz), 1.00 (9H, s), 0.18 (6H, s)]. In the present case the formation of another potential acylanion equivalent was investigated; treatment with trimethylsilylcyanide⁹ yielded (95 %) the protected cyanohydrin 7b as a mixture of two diastereoisomers [$\delta(\text{CCl}_4)$: 4.66 (2H, s), 4.54 (1H, m), 4.0 (1H, m), 3.53 (2H, m), 2.22 (3H, s), 1.99 (2H, m), 1.83 (1H, m), 1.06 and 1.03 (3H, d), 1.01 (9H, s), $0.18-0.05$ (15H)].

Both synthons 7a and 7b are synthetically equivalent and allow the introduction of the "southern zone" of the title compound as has been shown by others^{1c}.

In summary we have shown the easy and short construction of the carbon chain 5 to 9 with the correct stereochemistry by oxidative cleavage of an adequately substituted cyclopentene precursor. Since (S)-(+)-4-hydroxy-2-cyclopentenone (3, $R_1 = \text{H}$) has been shown to be readily available from (R,R)-(+)-tartaric acid¹⁰; this approach should prove ideal for a chiral synthesis of the "eastern zone"¹¹. Further work is in progress.

ACKNOWLEDGEMENT

We are indebted to the "Nationaal Fonds voor Wetenschappelijk Onderzoek" for financial help to the laboratory.

REFERENCES

1. a) A.I. Meyers, C.C. Shaw, D. Horne, L.M. Trefonas and R.J. Majeste, *Tetrahedron Lett.*, 1745 (1975); b) A.I. Meyers and R.S. Brinkmeyer, *ibid.*, 1749 (1975); c) E.J. Corey and M.G. Bock, *ibid.*, 2643 (1975); d) W.J. Elliot and J. Fried, *J. Org. Chem.*, 41, 2469 (1976); e) K.M. Kane and A.I. Meyers, *Tetrahedron Lett.*, 771 (1977); f) J.E. Foy and B. Ganem, *ibid.*, 775 (1977); g) E.J. Corey, H.F. Wetter, A.P. Kozikowski and V.R. Rao, *ibid.*, 777 (1977); h) O.E. Edwards and P.-T. Ho, *Can. J. Chem.*, 55, 371 (1977).

2. S. Remillard, L.I. Rebhurn, G.A. Howie, S.M. Kupchan, *Science*, **189**, 1005 (1975) and references cited herein.
3. G. Stork and M. Isobe, *J. Am. Chem. Soc.* **97**, 6260 (1975).
4. There are several examples in prostaglandin chemistry (e.g.; see ref. 3 and ref. cited therein) of this conjugate addition leading exclusively to the trans relation of the 3- and 4-substituents.
5. E.J. Corey and A. Venkateswarlu, *J. Am. Chem. Soc.* **94**, 6190 (1972).
6. J.M. Lalancette and A. Lachance, *Can. J. Chem.*, **47**, 859 (1969); the normal procedures for synthesizing dithianes with the aid of Lewis acids all resulted in cleavage of the protective groups; see e.g. D. Seebach and E.J. Corey, *J. Org. Chem.*, **40**, 231 (1975); although methylthiotrimethylsilane is also known to react under neutral conditions, we were not able to achieve this reaction, even on simple model compounds, see D.A. Evans, K.C. Grimm and L. Truesdale, *J. Am. Chem. Soc.* **97**, 3229 (1975).
7. 4-t.Butyldimethylsilyloxy-2-cyclopentenone was prepared by monosilylation followed by oxidation with Jones reagent of the readily available cis-1,4-dihydroxy-2-cyclopentene : C. Kaneko, A. Sugimoto and S. Tanaka, *Synthesis*, 876 (1975).
8. P.M. Pojer and S.J. Angyal, *Tetrahedron Lett.*, 3067 (1976).
9. D.A. Evans, G.L. Carroll and L.K. Truesdale, *J. Org. Chem.*, **39**, 917 (1974); S. Hunig and G. Wehner, *Synthesis*, 391 (1975).
10. K. Ogura, M. Yamashita and G.-I. Tsuchihashi, *Tetrahedron Lett.*, 759 (1976); it should be noted however that in this paper the absolute configuration of the hydroxycyclopentenone obtained from D-tartaric acid was erroneously assigned.
11. An optically active "eastern zone" has been obtained by resolution; see reference 1d.