

View Article Online View Journal

Dalton Transactions

Accepted Manuscript

This article can be cited before page numbers have been issued, to do this please use: M. Toure, O. Chuzel and J. PARRAIN, *Dalton Trans.*, 2015, DOI: 10.1039/C5DT00814J.



This is an *Accepted Manuscript*, which has been through the Royal Society of Chemistry peer review process and has been accepted for publication.

Accepted Manuscripts are published online shortly after acceptance, before technical editing, formatting and proof reading. Using this free service, authors can make their results available to the community, in citable form, before we publish the edited article. We will replace this Accepted Manuscript with the edited and formatted Advance Article as soon as it is available.

You can find more information about *Accepted Manuscripts* in the **Information for Authors**.

Please note that technical editing may introduce minor changes to the text and/or graphics, which may alter content. The journal's standard <u>Terms & Conditions</u> and the <u>Ethical guidelines</u> still apply. In no event shall the Royal Society of Chemistry be held responsible for any errors or omissions in this *Accepted Manuscript* or any consequences arising from the use of any information it contains.



www.rsc.org/dalton

RSCPublishing

Journal Name

COMMUNICATION

Cite this: DOI: 10.1039/x0xx00000x

Received 00th January 2012, Accepted 00th January 2012

DOI: 10.1039/x0xx00000x

www.rsc.org/

Published on 24 March 2015. Downloaded by Northern Illinois University on 25/03/2015 00:52:45.

Synthesis and structure of Ag(I), Pd(II), Rh(I), Ru(II) and Au(I) NHC-complexes with pendant Lewis acidic boronic ester moiety[†]

Momar Toure,^a Olivier Chuzel^{a,*} and Jean-Luc Parrain^{*,a}

situ (Scheme 1).

Bifunctional Ag(I), Pd(II), Rh(I), Ru(II) and Au(I) complexes containing a NHC ligand and pendant trivalent boron moiety have been synthesized in high yields. Fine-tuned reaction conditions were used to prevent potential ligand selfquenching or polymerization due to the eventual co-existence in situ of free NHC (Lewis base) and boronic ester (Lewis acid) in the same molecule.

Boron-centered ligands have relatively large structural and electronical diversity allowing for different coordination mode to a metal center, as found, in borane, boryl, borylene or boride complexes.¹ In some particular cases of Lewis-basic metals, the Lewis-acidic trivalent boron atom can be coordinated to the metal center by a dative bond, becoming a σ -acceptor Z-type ligand,² mainly with borane derivatives,³ first authenticated by Hill et al. through the isolation of a metallaboratrane in 1999.⁴ Complexes featuring this supported M-B Z-type interactions are derived from the B-H activation of hydrido tris(methimazolyl)borates and related systems, or are obtained directly from ambiphilic phosphine borane ligands⁵ where the boron atom moiety is structurally maintained close to the metal center.

There are rare examples of bidentate or bifunctional ligands containing a boronic ester moiety that are reported in the literature,⁶⁻⁹ and the only observed transition metal-Z interactions with boronic ester moiety disclosed so far are the bridging borane-boryl compounds.¹⁰ Because the structural feature of those bifunctional ligands does not allow the boronic ester moiety to coordinate to the metal center, we designed a more flexible bifunctional ligand allowing the possibility of such an interaction. To increase the interaction, strong σ -donor N-heterocyclic carbene (NHC) ligand was chosen to enhance the Lewis base character of the transition metal. Synthesis of such NHC-boron transition metal complexes was highly challenging due to the necessary generation of coexisting free NHC (Lewis base) and not hindered boron derivative (Lewis acid) in or oligomers



Scheme 1 Possible NHC/boronic ester/metal combined product pathways.

Here, we report the synthesis and the characterization of original stable bifunctional NHC-boronic ester silver, palladium, rhodium, ruthenium and gold complexes. Despite several modifications, the boronic ester moiety remained pendant and no Z-type interaction to the metal center was observed in this series of complexes (Scheme 1).

In the initial attempts, NHC-boron salts were obtained by reacting phenyl-substituted imidazole **1a** with the bromopropyl dioxaborolane 2 that was prepared in one step through hydroboration of allylbromide.¹¹ Attempts to prepare the complex **5a** directly from imidazolium salt 1a and classic bases such as n-BuLi,¹² KHMDS, NaH or t-BuOK did not succeed and only degradation of the starting material was observed. No self-quenching resulting in cyclic NHCboranes, polymerization processes or frustrated Lewis pair was detected. To generate the carbenoid species and avoid decomposition, we decided to use silver oxide as a soft base. Gratifyingly the reaction between Ag₂O and the imidazolium salt **1a** gave the desired cationic bis-NHC silver complex 4a in quantitative yield, demonstrating the importance of the silver oxide mediated approach (Scheme 2). The structure was confirmed by NMR and mass spectrometry analysis (ESI, MeOH, m/z for [M-AgBr₂]⁺

731.4). Subsequent treatment of 4a with 2.5 equiv of PdCl₂(CH₃CN)₂ at room temperature in dichloromethane afforded binuclear dichloride complexes 5a in 84% yield. Transmetallation of the silver complex with other transition



Scheme 2 Synthesis of Ag-NHC boronic ester complexes.

Published on 24 March 2015. Downloaded by Northern Illinois University on 25/03/2015 00:52:45.



Fig.1 Structures of palladium, rhodium, gold and ruthenium NHC-boronic ester complexes.

metals was also investigated. The reactivity of 4a towards [Rh(η^4 cod)Cl]2 or AuCl metallic sources afforded the stable neutral complexes 6a and 7a in 74% and 87% yield respectively (Fig. 1). Negligible difference in chemical shift of the boron atom in ¹¹B NMR (128 MHz, CDCl₃) for imidazolium salts 3a (33.3 ppm) and all complexes prepared (33.3-34.0 ppm) showed that the boron moiety was not ligated to the metal center and remained pendant. This initial evidence was further confirmed for 5a, 6a and 7a in the solid state by X-ray diffraction studies (Fig. 2).

Interestingly, for complexes 5a and 6a the existence of hydrogen bonding or anagostic interaction (three-centers / four-electrons) between the metal and the hydrogen atom of the methylene group in α to the NHC was suspected by ¹H NMR (400 MHz, CDCl₃). Hydrogen atoms of this CH_2 group were equivalent both in 3a (4.57

ppm, 2H, t, ${}^{3}J = 7.3$ Hz) and **4a** (4.13 ppm, 2H, t, ${}^{3}J = 7.3$ Hz), but a downfield shift of these protons in an unresolved broad signal appeared for 5a at room temperature (4.40-4.95 ppm, 2H, m). Upon cooling to 243 K in CD₂Cl₂ solution, this signal split into two distinct sets attributed respectively to the hydrogen atom participating in an interaction with the palladium atom (4.57-4.91 ppm, 1H, m) and to the one that is not (4.27-4.57 ppm, 1H, m) (See ESI). In the case of rhodium complex 6a the ¹H NMR signals of one of these two protons is strongly shifted downfield and these protons appeared as two distinct sets of broad signals at room temperature (4.32 and 4.89 ppm for non-coordinated and coordinated hydrogen atoms respectively) (see ESI). The downfield chemical shift of one of these protons indicated a clearer rhodium-H interaction than those observed for complex 5a, typical of metal-hydrogen bonding or anagostic interaction. The existence of such interactions in 5a and 6a was confirmed in the solid state by X-ray diffraction study (Fig. 2) and their structural parameters (Pd1-H10 2.8203(4) Å, Pd1-C10 3.338(6) Å, Pd1-H10-C10 114.31(3)°; Pd2-H28 2.8580(5) Å, Pd2-C28 3.346(8) Å, Pd2–H28–C28 112.08(4)°; (Rh1–H10) 2.9297(2) Å, Rh1–C10 = 3.449(3) Å and angle (Rh1–H10–C10) = $114.65(2)^{\circ}$) are in good agreement with those of anagostic interaction in transition metal d⁸ complexes (litt.¹⁴ d(M–H) \approx 2.3-2.9 Å, M–H–C \approx 110-170°). The origin of these interactions is still under debate and may involve donation of filled d_{z2} or $d_{xz/yz}$ orbitals of the metal center into the C-H σ^* orbital.¹⁵ In order to obtain a more detailed understanding of the involved orbitals, a Natural Bond Orbital (NBO)¹⁶ analysis was computed (see ESI for details). Two distinct interactions were found, one involving the sp³ C–H orbitals and the antibonding rhodium s-orbital and a second interaction between the antibonding sp³ C-H with a rhodium lone pair with second order perturbation energies, $E_{2P} = 2.94$ and 2.44 kJ mol⁻¹, respectively. Moreover, the hydrogen atom close to the rhodium atom has a slightly higher natural charge of 0.222 than the other hydrogen atom bonded by the same carbon atom (0.204). To conclude, the M-H bond distance and the M–H–C angle values and the very low E_{2P} (<5 kJ mol⁻¹) strongly suggest the establishment of a very weak hydrogen bond,¹⁷ probably led to the alkyl chain conformation. Such interaction was not observed in the corresponding silver or gold complex 4a and 7a respectively.

The presence of privileged conformers having equivalent (4a, 7a) or not equivalent (5a, 6a) protons bearing by the methylene group α to the NHC is of interest. In the last case, this effect is strongly pronounced in complex 6a and to estimate the participation of the rhodium complex geometry and/or the boronic ester moiety in such a)





Published on 24 March 2015. Downloaded by Northern Illinois University on 25/03/2015 00:52:45.

Journal Name



Fig. 2 ORTEP diagram of a) [PdCl₂NHC]₂ complex **5a**; b) [RhCINHC] complex **6a** were depicted with thermal ellipsoids at 50% probability. Most of hydrogen atoms are omitted for clarity.

preferential conformation the analogous rhodium complex **11** (scheme 3) was synthesized. It was observed in **11** that these two protons were downfield shifted and not equivalent, but the effect was not as important as observed in **6a** (see ESI). Clearly, the boronic ester moiety has an effect on the conformation of the alkyl chain, leading in a rhodium complex **6a** more stabilized by an additional C–H hydrogen interaction. This, could generated a difference in the reactivity of **6a** and **11** complexes (see further).

Furthermore, to increase the Lewis basicity at the metallic center and the possibility of Z-interaction to the boron atom, a donor pyrimidine hemilabile ligand was incorporated at the N-position of the NHC instead of the phenyl group (Scheme 2). ¹H NMR and ¹³C NMR spectra of palladium dichloride complex 5b showed clearly three non-equivalent pyrimidine hydrogen and carbon atoms with a downfield shift ($\delta H > 1$) of one aromatic hydrogen atom in the α position to the nitrogen atom. As expected, these observations clearly indicate bidentate coordination of the metal to the NHC and one of the nitrogen atom of the pyrimidine moiety. In the solid state, complex 5b features a pseudo square planar geometry with a Pd1-N1 distance of 2.049(2) Å and a N1-Pd1-C7 and N1-Pd1-Cl1 angles of 80.18(1)° and 93.47(7)°, respectively, that confirms the spectroscopic NMR data (Fig. 3). Unfortunately, attempts to synthesize the rhodium complex analogue from $[Rh(\eta^4-cod)Cl]_2$ gave an undetermined complex mixture.

Alternatively, the treatment of **4b** with $[\text{Ru}(p\text{-cymene})\text{Cl}_2]_2$ gave quantitatively the corresponding 18-electrons ruthenium(II) complex **8b**. The cationic nature of the complex was correlated by high-resolution mass spectrometry (ESI, MeOH, m/z 585.1742 [M-Cl]⁺). Unfortunately, all attempts to crystallize complex **8b** failed, but the NMR spectroscopic data, as seen previously in the complex **5b**, provided evidence for the bidentate nature of the NHC-pyrimidine ligand as well as the pendant nature of the boron moiety (¹¹B NMR, 128 MHz, in CDCl₃, 34.0 ppm). As expected, reaction of **4b** with AuCl gold source gave the linear geometric complex **7b**, where the pyrimidine aromatic ring was twisted to hold the gold metal center far from the nitrogen atom of the pyrimidine (Au1–N3, 3.118(1) Å, Au1– a)



Fig. 3 ORTEP diagram of a) [PdCl₂NHC] complex 5b; b) [AuCINHC] complex 7b were depicted with thermal ellipsoids at 50% probability. Hydrogen atoms are omitted for clarity.

N3–C7–N2 dihedral angle, -34.91(1)°). As previously observed in this series of complexes, NMR data and X-ray characterization revealed again the pendant character of the boron moiety in the complex **7b** (11 B NMR, 128 MHz, in CDCl₃, 33.8 ppm) (Fig. 3).

Despite some evidence in the literature of transition metal-Z interactions with boronic ester moiety,¹⁰ or related reaction intermediate found in borylation reaction mechanism,¹³ such interactions were not observed in this series of flexible complexes, owing to a less buttress structure that do not allow potential recruitment of the boron atom moiety close to the metal center. Interestingly, these stable complexes exhibit a Lewis acidic pendant boronic ester moiety that could serve as an additional binding site in bifunctional catalysis. An initial 1:1 equivalent mixture of complex 5a and Et₃N was analyzed by ¹¹B NMR (128 MHz, CDCl₃) showing a complete upfield chemical shift of the boron atom (from 33.5 to 2.85 ppm) corresponding to a borate species that revealed an unambiguous covalent B-N bond. This information let us confidant to use the boron atom in these complexes as an additional electrophilic activating site. In this context, stability and reactivity of complex **6a** were evaluated in a preliminary hydrosilylation reaction and **6a** was found to be a very efficient catalyst (conv > 99 % by 1 H NMR) (Scheme 3). However, the analogous complex 11 gave the same results in terms of kinetic (see ESI), meaning that the boronic part of the ligand seems to behave only as a spectator in this reaction. Nevertheless, the use of the stable palladium, rhodium, ruthenium and gold complexes described here as potential catalysts for the synergistic activation of reaction partners will be explored soon, as well as extension to structurally analogous more electrophilic NHC-borane ligands.





Conclusions

we have reported the In summary, synthesis and characterization of novel bifunctional NHC-boronic ester ligands and their corresponding silver(I), palladium(II), rhodium(I), ruthenium(II) and gold(I) complexes.¹⁸ It was observed that only the use of Ag₂O as a base allowed us to obtain these complexes in very good yields, and without either a self-quenching or a polymerization process taking place during the generation of the in situ Lewis base (NHC) and Lewis acid (boronic ester). It was confirmed by ¹¹B NMR and X-ray diffraction that the boron moiety is not coordinated at the metal center and remained pendant. Complexes bearing pendant trivalent borane derivatives are of interest¹⁹ and could be an alternative to enhance potential electrophilic activation of a substrate. Reported examples are essentially built on metallocene shapes system,20 but many of them must be stabilized in situ as borate derivatives,²¹ to avoid potential intramolecular rearrangement.²² The stable complexes developed here would let us confidently use them in bifunctional catalysis in the cooperative activation of small molecules.⁹ These hypothesis will be examined in further detail in due course.

Acknowledgments

This work was supported by grant from the French Research Ministry to M.T., Aix-Marseille Université, CNRS UMR 7313. We thanks M. Giorgi (www.spectropole.fr) for the X-ray diffraction studies.

Notes and references

^{*a*} Aix-Marseille Université, Centrale Marseille, CNRS, iSm2 UMR 7313, 13397, Marseille, France. E-mail: olivier.chuzel@univ-amu.fr, jl.parrain@univ-amu.fr

† Electronic Supplementary Information (ESI) available: Experimental procedures, spectroscopic data, detailed calculations and CIF for complexes 5a, 5b, 6a, 7a and 7b. DOI: 10.1039/c000000x/

 H. Braunschweig, C. Kollann and D. Rais, *Angew. Chem. Int. Ed.*, 2006, 45, 5254; H. Braunschweig, R. D. Dewhurst and A. Schneider, *Chem. Rev.*, 2010, 110, 3924.

- H. Braunschweig and R. D. Dewhurst, *Dalton Trans.*, 2011, **40**, 549;
 A. Amgoune and D. Bourissou, *Chem. Commun.*, 2011, **47**, 859;
 H. Kameo and H. Nakazawa, *Chem. Asian J.*, 2013, **8**, 1720;
 G. R. Owen, *Chem. Soc. Rev.*, 2012, **41**, 3535;
 M. Sircoglou, S. Bontemps, M. Mercy, N. Saffon, M. Takahashi, G. Bouhadir, L. Maron and D. Bourissou, *Angew. Chem. Int. Ed.*, 2007, **46**, 8583;
 A. F. Hill, *Organometallics*, 2006, **25**, 4741;
 G. Parkin, *Organometallics*, 2006, **25**, 4744;
 F.-G. Fontaine, J. Boudreau and M.-H. Thibeault, *Eur. J. Inorg. Chem.*, 2008, 5439.
- Selected references: F. Inagaki, C. Matsumoto, Y. Okada, N. 3 Maruyama and C. Mukai, Angew. Chem. Int. Ed., 2015, 54, 818; W. H. Harman, T.-P. Lin and J. C. Peters, Angew. Chem. Int. Ed., 2014, 53, 1081; M.-E. Moret and J. C. Peters, Angew. Chem. Int. Ed., 2011, 50, 2063; H. Fong, M.-E. Moret, Y. Lee and J. C. Peters, Organometallics, 2013, 32, 3053; W. H. Harman and J. C. Peters, J. Am. Chem. Soc., 2012, 134, 5080; S. N. MacMillan, W. H. Harman and J. C. Peters, Chem. Sci., 2014, 5, 590; M. Sircoglou, S. Bontemps, G. Bouhadir, N. Saffon, K. Miqueu, W. Gu, M. Mercy, C.-H. Chen, B. M. Foxman, L. Maron, O. V. Ozerov and D. Bourissou, J. Am. Chem. Soc., 2008, 130, 16729; M.-E. Moret, Y. Lee and J. C. Peters, J. Am. Chem. Soc., 2011, 133, 18118; D. L. M. Suess, C. Tsay and J. C. Peters, J. Am. Chem. Soc., 2012, 134, 14158; H. Kameo, Y. Hashimoto and H. Nakazawa, Organometallics, 2012, 31, 3155; M. Sircoglou, S. Bontemps, M. Mercy, K. Miqueu, S. Ladeira, N. Saffon, L. Maron, G. Bouhadir and D. Bourissou, Inorg. Chem., 2010, 49, 3983; B. E. Cowie and D. J. H. Emslie, Chem. Eur. J., 2014, 20, doi: 10.1002/chem.201404846; W. A. Gunderson, D. L. M. Suess, H. Fong, X. Wang, C. M. Hoffmann, G. E. Cutsail III, J. C. Peters and B. M. Hoffman, J. Am. Chem. Soc., 2014, 136, 14998.
- 4 A. F. Hill, G. R. Owen, A. J. P. White and D. J. Williams, *Angew. Chem. Int. Ed.*, 1999, **38**, 2759.
- 5 G. Bouhadir, A. Amgoune, D Bourissou and I. R. Crossley, In Advances in Organometallic Chemistry; A. F. Hill, M. J. Fink, ed.; 2010, Elsevier: Academic Press, Vol 8, p 1; S. Bontemps, G. Bouhadir, K. Miqueu and D. Bourissou, J. Am. Chem. Soc., 2006, 128, 12056.
- 6 A. Börner, J. Ward, K. Kortus and H. B. Kagan, *Tetrahedron: Asymmetry*, 1993, 4, 2219; L. B. Fields and E. N. Jacobsen, *Tetrahedron: Asymmetry*, 1993, 4, 2229.
- 7 S. Chikkali, S. Magens, D. Gudat, M. Nieger, I. Hartenbach and T. Schleid, *Eur. J. Inorg. Chem.*, 2008, 2207.
- 8 J. Navarro, O. Torres, M. Martín and E. Sola, J. Am. Chem. Soc., 2011, 133, 9738.
- 9 O. Tutusaus, C. Ni and N. K. Szymczak, J. Am. Chem. Soc., 2013, 135, 3403.
- 10 D. Curtis, M. J. G. Lesley, N. C. Norman, A. G. Orpen and J. Starbuck, J. Chem. Soc., Dalton Trans., 1999, 1687; S. A. Westcott, T. B. Marder, R. T. Baker, R. L. Harlow, J. C. Calabrese, K. C. Lam and Z. Lin, Polyhedron, 2004, 23, 2665; R. S. Anju, D. K. Roy, K. Geetharani, B. Mondal, B. Vargheseb and S. Ghosh, Dalton Trans., 2013, 42, 12828.
- 11 G. A. Molander, C. S. Yun, M. Ribagorda and B. Biolatto, J. Org. Chem., 2003, 68, 5534.
- 12 Wang *et al.* have synthesized a NHC-borane platinium complex but the tethered and rigid structure of the free ligand do not allow selfquenching, see: Z. M. Hudson, C. Sun, M. G. Helander, Y.-L. Chang,

Page 5 of 5

Published on 24 March 2015. Downloaded by Northern Illinois University on 25/03/2015 00:52:45.

Z.-H. Lu and S. Wang, J. Am. Chem. Soc., 2012, **134**, 13930. By this way, a more flexible structure led to the corresponding Lewis pair ions, see: Y.-L. Rao, L. D. Chen, N. J. Mosey and S. Wang, J. Am. Chem. Soc., 2012, **134**, 11026.

- 13 I. A. I. Mkhalid, J. H. Barnard, T. B. Marder, J. M. Murphy and J. F. Hartwig, *Chem. Rev.*, 2010, **110**, 890; Q. Li, C. W. Liskey and J. F. Hartwig, *J. Am. Chem. Soc.*, 2014, **136**, 8755.
- D. Braga, F. Grepioni and E. Tedesco, Organometallics, 1997, 16, 1846; M. Brookhart, M. L. H. Green and G. Parkin, Proc. Natl. Acad. Sci., 2007, 104, 6908; A. Mukhopadhyay and S. Pal, Eur. J. Inorg. Chem., 2006, 4879; R. Angamuthu, L. L. Gelauff, M. A. Siegler, A. L. Spek and E.Bouwman, Chem. Commun. 2009, 2700; A. T. Çolak, O. Z. Yesilel and O. Büyükgüngör, J. Mol. Struct. 2011, 991, 68; N. Singh and A. J. Elias, Dalton Trans. 2011, 40, 4882; A. G. Jarvis, P. E. Sehnal, S. E. Bajwa, A. C. Whitwood, X. Zhang, M. S. Cheung, Z. Lin and I. J. S. Fairlamb, Chem. Eur. J. 2013, 19, 6034; M. G. D. Holaday, G. Tarafdar, A Kumar, M. L. P. Reddy and A. Srinivasan, Dalton Trans., 2014, 43, 7699; H. V. Huynh, L. R. Wong and P. S. Ng, Organometallics, 2008, 27, 2231.
- 15 W. Yao, O. Eisenstein and R. H. Crabtree, *Inorg. Chim. Acta*, 1997, 254, 105; Y. Zhang, J. C. Lewis, R. Bergman, J. A. Ellman and E. Oldfield, *Organometallics*, 2006, 25, 3515.
- 16 E. D. Glendening, J. K. Badenhoop, A. E. Reed, J. E. Carpenter, J. A. Bohmann, C. M. Morales, C. R. Landis and F. Weinhold (Theoretical Chemistry Institute, University of Wisconsin, Madison, WI, 2013); http://nbo6.chem.wisc.edu/).
- 17 J. Saßmannshausen, Dalton Trans., 2012, 41, 1919.
- 18 To our knowledge, only one metallic complex featuring bifunctional NHC-boronic ester was reported, see reference 8.
- For bifunctional complex with pendant borane features, see: W.-J. Xu, S.-J. Liu, X.-Y. Zhao, S Sun, S. Cheng, T.-C. Ma, H.-B. Sun, Q. Zhao and W. Huang, *Chem. Eur. J.*, 2010, 16, 7125; J. Vergnaud, M. Grellier, G. Bouhadir, L. Vendier, S. Sabo- Etienne and D. Bourissou, *Organometallics*, 2008, 27, 1140; T. G. Ostapowicz, C. Merkens, M. Hölscher, J. Klankermayer and W. Leitner, *J. Am. Chem. Soc.*, 2013, 135, 2104; A. J. M. Miller, J. A. Labinger and J. E. Bercaw, *J. Am. Chem. Soc.*, 2008, 130, 11874; A. J. M. Miller, J. A. Labinger and J. E. Bercaw, *Organometallics*, 2010, 29, 4499; H. Braunschweig, R. Dirk and B. Ganter, *J. Organomet. Chem.*, 1997, 545-546, 257.
- 20 For bifunctional complexes based on a metallocene-borane pendant shape, see: R. E. v. H. Spence and W. E. Piers, Organometallics, 1995, 14, 4617; R. S. Rojas, B. C. Peoples, A. R. Cabrera, M. Valderrama, R. Fröhlich, G. Kehr, G. Erker, T. Wiegand and H. Eckert, Organometallics, 2011, 30, 6372; P. A. Deck, T. S. Fisher and J. S. Downey, Organometallics, 1997, 16, 1193; A. J. Ashe III, X. Fang and J. W. Kampf, Organometallics, 1999, 18, 2288; H. Braunschweig, R. Dirk, M. Müller, P. Nguyen, R. Resendes, D. P. Gates and I. Manners, Angew. Chem. Int. Ed., 1997, 36, 2338; A. Appel, F. Jäkle, T. Priermeier, R. Schmid and M. Wagner, Organometallics, 1996, 15, 1188; J. Chen, K. Venkatasubbaiah, T. Pakkirisamy, A. Doshi, A. Yusupov, Y. Patel, R. A. Lalancette and F. Jäkle, Chem. Eur. J., 2010, 16, 8861. (e) G. E. Herberich, A. Fisher, D. Wiebelhaus, Organometallics, 1996, 15, 3106; S. A. Larkin, J. T. Golden, P. J. Shapiro, G. P. A. Yap, D. M. J. Foo and A. L. Rheingold, Organometallics, 1996, 15, 2393.

- 21 M. Hill, G. Erker, G. Kehr, R. Fröhlich and O. Kataeva, J. Am. Chem. Soc., 2004, **126**, 11046; C. Herrmann, G. Kehr, R. Fröhlich and G. Erker, Organometallics, 2008, **27**, 2328; M. Emmert, G. Kehr, R. Fröhlich and G. Erker, Chem. Eur. J., 2009, **15**, 8124; S. J. Lancaster, S. Al-Benna, M. Thornton-Pett and M. Bochmann, Organometallics, 2000, **19**, 1599.
- 22 M. Hill, G. Kehr, G. Erker, O. Kataeva and R. Fröhlich, *Chem. Commun.*, 2004, 1020; C. Herrmann, G. Kehr, R. Fröhlich and G. Erker, *Eur. J. Inorg. Chem.*, 2008, 2273.