

Preliminary communication

SOME REACTIONS OF DIALKYL TIN OXIDES WITH ACETYL CHLORIDE AND TRIMETHYLCHLOROSILANE

SHIGEMI KOHAMA

The Osaka Municipal Technical Research Institute, Kitaogimachi 38, Kita-ku, Osaka (Japan)


(Received August 8th, 1975)

Summary

Dialkyltin oxides react with acetyl chloride to give $R_2SnClOAc$ or R_2SnCl_2 , depending on the stoichiometry, and with Me_3SiCl to give $(ClR_2Sn)_2O$.

It has been shown that dimethyltin dichloride reacts with acetic anhydride or glacial acetic acid to give dimethyltin chloride acetate [1] and that the reactions of various *sym*-(dichlorotetraalkyl)distanoxanes with acetic acid also give the corresponding chloride acetates [2].

Such chloride acetates now have been prepared by the reaction of dialkyltin oxide with an equimolar amount of acetyl chloride. Upon further addition of acetyl chloride, the dialkyltin chloride acetates were converted into the corresponding dichlorides. These also were easily formed from the oxides and a large excess of acetyl chloride.



($R = CH_3, C_2H_5, n-C_3H_7, n-C_4H_9, n-C_8H_{17}$)

It was found that the rate of digestion of dialkyltin oxides in acetyl chloride followed the sequence $CH_3 < C_2H_5 < n-C_3H_7 < n-C_4H_9 < n-C_8H_{17}$ for alkyl groups.

Similar treatment of dialkyltin oxides with organic halides such as ethyl bromide, *t*-butyl chloride, allyl chloride and benzyl chloride resulted in recovery of the starting materials.

Unlike the case with acetyl chloride, the reaction of dialkyltin oxides


TABLE 1

REACTIONS BETWEEN DIAKYL TIN OXIDES AND HALIDES

Reactants	Halide (g)	Product	M.p. (°C) Found (Lit.)	Yield (g(%))	Analysis (%) Found (calcd.)				Mol.wt. Found (calcd.)
					Sn	C	H	Cl	
R = Me (4)	AcCl 1.9	Me ₂ SnCl(OAc)	189-190 (189) [1]	4.6 (78)	48.81 (48.79)				
R = Me (4)	Me ₃ SiCl 2.6	(ClMe ₂ Sn) ₂ O	200 (—) [3]	3.3 (71)	61.74 (61.75)				
R = Et (2)	AcCl 0.8	Et ₂ SnCl(OAc)	93-94 (94) [2]	2.2 (80)	43.74 (43.75)				
R = Et (2)	Me ₃ SiCl 1.1	(ClEt ₂ Sn) ₂ O	174 (175.5) [4]	1.8 (79)	53.88 (53.88)				
R = n-Pr (2)	AcCl 0.7	Pr ₂ SnCl(OAc)	74-75 (73) [2]	2.2 (82)	39.72 (39.65)	31.92 (32.09)	5.75 (5.72)		316 (299)
R = n-Pr (2)	Me ₃ SiCl 1.0	(ClPr ₂ Sn) ₂ O	121-122 (122) [4]	2.0 (89)	47.82 (47.79)				
R = n-Bu (3)	AcCl 1.0	Bu ₂ SnCl(OAc)	65-66 (61) [2]	3.3 (84)	36.48 (36.25)	36.39 (36.65)	6.54 (6.49)	10.84 (10.83)	351 (327)
R = n-Bu (3)	Me ₃ SiCl 1.3	(ClBu ₂ Sn) ₂ O	112-113 (111-112) [5]	2.8 (85)	42.90 (42.95)				
R = n-Oct (3)	AcCl 0.7	Oct ₂ SnCl(OAc)	76	3.2 (87)	26.97 (26.99)	48.96 (48.18)	8.60 (8.48)	8.03 (8.06)	443 (439)
R = n-Oct (3)	Me ₃ SiCl 0.9	(ClOct ₂ Sn) ₂ O	88-89	2.7 (84)	30.51 (30.54)	49.27 (49.45)	8.90 (8.82)	8.65 (9.12)	1500 ^a (777)



^a As dimer.

with an equimolar quantity of trimethylchlorosilane did not give the expected trimethylsilyldialkylstannoxanes. Instead, *sym*-(dichlorotetraalkyl)distannoxanes were formed. The resulting distannoxanes were converted to dialkyltin dichlorides by further addition of trimethylchlorosilane. A large excess of trimethylchlorosilane, of course, converted the dialkyltin oxides directly to the dialkyltin dichlorides.



(R = CH₃, C₂H₅, n-C₃H₇, n-C₄H₉, n-C₈H₁₇)

It was confirmed that in the early stages of this reaction the dialkyltin dichlorides existed in a liquid layer of the reaction mixture. A possible explanation of the formation of the distannoxane is the reaction between initially produced dichloride and the unreacted oxide. Evidence for this was provided by our finding that equimolar quantities of dialkyltin dichloride and dialkyltin oxide react quantitatively to yield the distannoxane. Thus the reaction steps for the formation of *sym*-(dichlorotetraalkyl)distannoxane are:



Typical examples are shown in Table 1 (the octyl derivatives are new compounds)

Acetyl chloride and dialkyltin oxides reacted exothermally when they were mixed in chloroform. On the other hand, since trimethylchlorosilane and the oxides did not react smoothly under the same conditions, they were heated at reflux until the oxide dissolved. The solvent then was removed under reduced pressure and the dialkyltin chloride acetates or *sym*-(dichlorotetraalkyl)distannoxanes were recrystallized from petroleum ether. They were characterized by m.p., analysis and, in some cases, by molecular weight determination.

References

- 1 R. Okawara and E.G. Rochow, J. Amer. Chem. Soc., 82 (1960) 3285.
- 2 M. Wada, M. Shindo and R. Okawara, J. Organometal. Chem., 1 (1963) 95.
- 3 R. Okawara, Proc. Chem. Soc., (1961) 383.
- 4 R.K. Ingham, Chem. Rev., 60 (1960) 459.
- 5 A.J. Gibbons, A.K. Sawyer and A. Ross, J. Org. Chem., 26 (1961) 2304.