Formation of Cyclobutanones by the Photolytic Reaction of $(CO)_5Cr=C(OMe)Me$ with Electron-Rich Olefins

Miguel A. Sierra and Louis S. Hegedus*

Department of Chemistry, Colorado State University Fort Collins, Colorado 80523 Received November 30, 1988

Recent research efforts in these laboratories have centered on the development of useful organic synthetic methodology based on the photolytic reactions of chromium "Fischer" carbene complexes, particularly in regards to the development of new β -lactam syntheses.^{1,2} In the course of these studies it became evident that photolysis of chromium-carbene complexes resulted in the reversible production of chromium-ketene complexes, by a photochemically driven CO insertion into the chromium-carbene carbon double bond and that this unstable intermediate was responsible for β -lactam formation (eq 1).³



Chromium-carbene complexes such as 1 undergo thermal reactions with alkenes to produce cyclopropanes, presumably via metallacyclobutane intermediates.⁴ Free ketenes react with alkenes to give cyclobutanones in a formal 2 + 2 cycloaddition process, although unactivated alkenes require forcing conditions.⁵ To probe for the presence of complexed (or free) ketenes in photolytic reactions of chromium-carbene complexes, photolysis (Pyrex, 450-W Hanovia lamp) in the presence of olefins was examined, and the results are summarized in eq 2.

A variety of olefins were converted to cyclobutanones in good to excellent yield by this process. The crude yields, in all cases, were almost quantitative, and the crude material contained little if any byproduct. These cyclobutanones were relatively reactive and decomposed on silica gel, alumina, and flurosil, making purification by chromatography difficult. Pure material was obtained by evaporative distillation. The reaction was highly regio- and stereoselective, producing, in all cases, the single diastereoisomer shown with >95% selectivity. The regioselectivity corresponds to that resulting from attack of the more nucleophilic olefin carbon

(2) For a survey of the literature dealing with the use of chromium-carbene complexes in organic synthesis, see: Imwinkelried, R.; Hegedus, L. S. Organometallics 1988, 7, 702.

(3) (a) Hegedus, L. S.; deWeck, G.; D'Andrea, S. J. Am. Chem. Soc. 1988, 110, 2122. (b) Cyclobutanones have also been produced by intramolecular trapping of chromium-vinyl ketene complexes, produced by the thermal reaction of chromium-carbene complexes with enynes, see: Wulff, W. D.; Kaisler, R. W. Organometallics 1985, 4, 1461. Korkowski, P. F.; Hoye, T. R.; Rydberg, D. B. J. Am. Chem. Soc. 1988, 110, 2676. (c) Reaction of (phenyl)(methoxy)carbene-chromium complexes with enamides under 150 atm of CO pressure also produced cyclobutanones: Dorrer, B.; Fischer, E. O. Chem. Ber. 1974, 107, 2683.

(d) Reviews: (a) Dötz, K. H. Angew. Chem., Int. Ed. Engl. 1984, 23, 587.
(b) Brookhart, M.; Studabaker, W. B. Chem. Rev. 1987, 87, 411. For recent examples with electron poor olefins, see: (c) Wienand, A.; Reissig, H.-U. Tetrahedron Lett. 1988, 29, 2315. (d) Buchert, M.; Reissig, H.-U. Tetrahedron Lett. 1988, 29, 2319.

(5) (a) Ghosez, L.; O'Donnell, M. In *Pericyclic Reactions*; Houk, K., Lehr, R., Eds.; Academic Press: New York, 1977; Vol. 2, pp 79-140. (b) Brady, W. T. In *Chemistry of Ketenes, Allenes, and Related Compounds*; Patai, S., Ed.; Wiley: Chichester, U.K., 1980; pp 279-405.



on the (electrophilic) ketene carbonyl carbon. In all cases, the methyl group of the carbene was syn to the substituent on the olefin, and the methoxy group was anti. With 1,2-disubstituted olefins (2f) the stereochemistry of the olefin was maintained in the cyclobutanone product. Electron poor olefins, such as methyl acrylate and acrylonitrile, did not form cyclobutanones under the same conditions.

The structures and stereochemistry of 2a-f were elucidated by spectroscopic techniques. The constitution was verified by high resolution exact mass measurements. Each compound had a strong infrared absorption at 1775-1780 cm⁻¹, characteristic of a four-membered carbonyl compound. High field proton and carbon NMR spectra confirmed the assigned structures as well as the absence of stereoisomers. The assigned stereochemistry for 2a-f was confirmed by detailed NOE studies on 2b, 2c, and 2f.⁶ The intramolecular version of this reaction⁷ was also quite efficient, producing cyclobutanone 3 in excellent yield, as a single stereoisomer (assigned cis stereochemistry by NOE measurements). Remarkably, 3 was somewhat unstable to the normal oxidative isolation procedure (see below), and up to 30% of lactone 4 was isolated when this procedure was used. Cyclobutanone 3 was the sole product when the reaction was run under carbon monoxide, and the product was purified by chromatography. The regio- and stereochemistry (syn disposition of large groups) strictly corresponds to that observed in the reaction of ethoxyketene (generated from ethoxyacetyl chloride and triethylamine) and olefins.^{7,8} Thus, photolytic reactions of chromium-carbene complexes with olefins result in ketene-derived products, although the question of free versus coordinated ketene remains unresolved. The higher reactivity observed with the carbene-derived systems versus free ketenes⁷ (14 h at 25 °C vs 15 h at 80-100 °C), the higher yields (60-80% vs 40-60%), the stability of the carbene complex to photolytic degradation in the absence of substrate (48 vs 15 h), and the failure to detect ketene dimers either from the reactions described above or from photodegradation in the absense of substrate, all suggest that the metal mediates these reactions in some way.



3 (86%) (3)

The very wide range of chromium-carbene complexes available⁹ coupled with a wide variety of reactions which substituted cyclobutanones undergo¹⁰ make the above process potentially useful

hv, CO, MeCN

^{(1) (}a) Borel, C.; Hegedus, L. S.; Krebs, J.; Satoh, Y. J. Am. Chem. Soc. **1987**, 109, 1101 and references therein. (b) Hegedus, L. S.; D'Andrea, S. J. Org. Chem. **1988**, 53, 3113.

⁽⁶⁾ Physical data (infrared, ¹H, ¹³C NMR spectra, mass spectra, exact mass measurements, and NOE and COSY data) are available as Supplementary Material.

⁽⁷⁾ For a review of intramolecular cycloaddition reactions of ketenes with alkenes, see: Snider, B. B. Chem. Rev. 1988, 88, 793. For intramolecular alkoxyketene/alkene cycloadditions, see: Snider, B. B.; Hui, R. A. H. F. J. Org. Chem. 1985, 50, 5167. Brady, W. T.; Giang, Y. F. J. Org. Chem. 1985, 50, 5177.

⁽⁸⁾ DoMinh, T.; Strausz, O. P. J. Am. Chem. Soc. 1970, 92, 1766.

⁽⁹⁾ Fischer, H. In Transition Metal Carbene Complexes; Verlag Chemie: Deerfield Beach, FL, 1983; pp 1-68.

for organic synthesis. Results of similar studies with aminocarbenes promise even wider application and will be reported in due time.

A typical experimental procedure follows.

Chromium-carbene complex 1 (0.30 g, 1.20 mmol) was placed in a 30-mL Pyrex test tube which was then sealed with a rubber septum and evacuated and purged with argon $(3\times)$. N-Vinylacetamide (0.12 g, 1.39 mmol) in 20 mL of degassed dry acetonitrile was added by syringe, and the resulting solution was irradiated for 15 h (450-W Conrad-Hanovia 7825 medium pressure mercury lamp, Pyrex well). The solvent was then removed under vacuum, and the yellow residue was dissolved in ethyl acetate, filtered through Celite, diluted with one volume of hexane, and permitted to air oxidize overnight in a light box (6-20 W Vitalite fluorescent bulbs). Filtration of the brown suspension, solvent removal, and evaporative distillation (0.2 mm, \sim 120 °C) gave 0.18 g (72%) of 2d as a colorless viscous oil.

Acknowledgment. Support for this research under Grant 2 RO1 GM26178-10 from the National Institutes of General Medical Sciences (Public Health Service) is gratefully acknowledged. M.A.S. thanks the Ministerio de Educacion y Cienca Espana for a postdoctoral fellowship. Mass spectrometry determinations were performed by Midwest Center for Mass Spectrometry, a National Science Foundation Regional Instrumentation facility (Grant No. CHE-8620177).

Supplementary Material Available: Full characterization and spectral data (¹H NMR, ¹³C NMR, and IR) for all new compounds (4 pages). Ordering information is given on any current masthead page.

(10) Bellus, D.; Ernst, B. Angew. Chem., Int. Ed. Engl. 1988, 27, 797.

Catalytic Partial Oxidation of Methane over BaPbO₃: **Possible Involvement of Peroxide Ion**

Karl C. C. Kharas and Jack H. Lunsford*

Department of Chemistry, Texas A&M University College Station, Texas 77843 Received December 7, 1988

Although numerous metal oxides catalyze the oxidative dimerization of methane to higher hydrocarbons,¹ the nature of the catalytically active sites is substantially unknown. Previous speculation that peroxide or superoxide ions are responsible for methane activation over Sm_2O_3^2 and over Na^+/MgO at high temperatures³ is supported by the fact that several simple peroxides, e.g., Na₂O₂, SrO₂, and BaO₂, can stoichiometrically oxidize methane.⁴ Here we show, for the first time, the existence of

Table I. Steady-State Methane Oxidation over BaPbO3^a

space velocity (h ⁻¹)	P _{CH4} (torr)	$P_{\rm CH_4}/P_{\rm O_2}$	$\begin{array}{c} C_2 \\ \text{productivity}^b \\ [\text{mmol} \cdot (\text{g cat})^{-1} \cdot h^{-1}] \end{array}$	C ₂ selectivity ^c (%)	C ₂ yield ^d (%)	
13300	323	4.4	16.0	60.4	14.0	
9100	100	2.4	6.0	51.8	18.0	
10000	106	2.2	7.8	48.1	16.3	
21100	100	2.5	9.4	68.8	13.7	

^aBaPbO₃ (0.35-0.55 g), 875 °C. ^bDefined as mol of CH₄ converted to C_2 products per g catalyst per h. ^cTypically, the C_2H_4/C_2H_6 ratio was 1.1. ^d The C_3^+ yield was <1%.



Figure 1. XPS spectra of unused (bottom) and used (top) BaPbO₃ catalysts. a. O_{1s} region bottom: O^{2-} , 528.4 eV; O_2^{2-} , 530.1 eV; top: O^{2-} , 528.6 eV; $O_2^{2^-}$, 530.3 eV; $CO_3^{2^-}$, 531.6 eV; HCO_3^- , 534.1 eV. b. C_{1s} region bottom: 284.6 eV (by definition) and 286.1, adventitious C; top: 284.6 eV (def.) and 286.1, adventitious C; CO₃²⁻, 288.7 eV; HCO₃²⁻, 292.3 eV.

peroxide ions on a complex metal oxide surface that catalyzes the oxidation of methane to C_2 hydrocarbons. We suggest that these peroxides may exist in a novel, through-bond isomer similar electronically to ozone.5

Barium metaplumbate⁶ was treated in flowing O₂ for several hours before beginning the oxidation reaction. As shown in Table I, moderate yields and high C₂ selectivities were obtained at 875 °C in a conventional 4 mm ID plug-flow reactor when the initial O_2 to CH₄ molar ratio was ~0.5. No methane oxidation was observed in similar experiments wherein quartz chips replaced the BaPbO₃ catalyst. These steady-state results were obtained after 2 h on stream; however, constant C2 productivity was observed initially. This lack of an induction period for C₂ production is consistent with the hypothesis that the BaPbO₃ surface is responsible for the catalytic activity.

The XPS spectrum⁷ of BaPbO₃ after annealing overnight in flowing O₂ at 875 °C, shown in the lowers traces of Figure 1, indicates that this material is not a typical A²⁺B⁴⁺[O²⁻]₃ perovskite.⁸ The binding energies and relative intensities (2.0:1.0)

^{(1) (}a) Keller, G. E.; Bhasin, M. M. J. Catal. 1985, 73, 9-19. (b) Ito, T.; Lunsford, J. H. Nature 1985, 314, 721-722. (c) Campbell, K. D.; Zhang, H.; Lunsford, J. H. J. Phys. Chem. 1988, 92, 750-753. (d) Jones, C. A.; Leonard, J. J.; Sofranko, J. A. Energy Fuels 1987, 1, 12-16. (e) Asami, K.; Hashimoto, S.; Shikada, T. Chem. Lett. 1986, 1233-1236. (f) Aika, K.; Moriyama, T.; Takazaki, N. J. Chem. Soc. Chem. Commun. 1986, 210-211. (g) DeBoy, J. M.; Hicks, R. F. J. Chem. Soc., Chem. Commun. 1988, 982-984. (h) Otsuka, K.; Jinno, K.; Morikawa, A. Chem. Lett. 1988, (h) Otsuka, K.; Jinno, K.; Morikawa, A. Chem. Lett. 1988, (i) Inai, H.; Tagawa, T. J. Chem. Soc., Chem. Commun. 1986, 52–53, 1986.
 (j) Machida, K.; Enyo, M. J. Chem. Soc., Chem. Commun. 1987, 1639–1640.
 (k) Nagamoto, H.; Amanuma, K.; Nobutomo, H.; Inoue, H. Chem. Lett. 1988, 237–240.
 (l) Lee, J. S.; Oyama, S. T. Catal. Rev. Sci. Eng. 1988, 30, 249-280.

⁽²⁾ Otsuka, K.; Jinno, K. Inorg. Chim. Acta 1986, 121, 237-241.

^{(3) (}a) Lunsford, J. H.; Lin, C.-H.; Wang, J.-X.; Campbell, K. D. Mat. Res. Soc. Symp. Proc. 1988, 111, 305-313. (b) Lunsford, J. H. In Oxygen Complexes and Oxygen Activation by Transition Metals; Martell, A. E.; Sawyer, D. T., Eds.; Plenum Press: New York, 1988; pp 265-272. (4) Otsuka, K.; Said, A. A.; Jinno, K.; Komatsu, T. Chem. Lett. 1987,

^{77-80.}

⁽⁵⁾ Goddard, W. A. III; Dunning, T. H., Jr.; Hunt, W. J.; Hay, P. J. Acc. Chem. Res. 1973, 6, 368-376.

⁽⁶⁾ Vidyasagar, K.; Gopalakrishnan, J.; Rao, C. N. R. J. Solid State Chem. 1985, 58, 29-37.

⁽⁷⁾ X-ray photoelectron spectra were acquired with an H.P. 5950A spectrometer equipped with an Al K α source, a quartz monochromator, and a flood trometer equipped with an Al K α source, a quartz monochromator, and a flood gun that was set at 1.0 V, 0.3 mA. Analyzer chamber pressures were less than 5×10^{-9} Torr. Binding energies were determined with respect to adventitious carbon whose binding energy was defined as 284.6 eV.¹⁷ (8) (a) Rao, C. N. R.; Ganguly, P.; Sarma, D. D. J. Am. Chem. Soc. 1987, 109, 6893-6895. (b) Ganguly, P.; Hegde, M. S. Phys. Rev. B 1988, 37, 5107-5112. (c) Hegde, M. S.; Ganguly, P. Phys. Rev. B 1988, 38, 4557-4561.