


This article was downloaded by: [Queensland University of Technology]
On: 20 November 2014, At: 23:09
Publisher: Taylor & Francis
Informa Ltd Registered in England and Wales Registered Number: 1072954
Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH,
UK



Phosphorus, Sulfur, and Silicon and the Related Elements

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/gpss20>

Regioselective Cycloadditions of β -Substituted Vinylphosphonate with Nitrile Oxides

Yong Ye ^{a b}, Li-bo Ma ^{a b}, Yong Luo ^b, Wen-hu Wang ^b, Lun-zu Liu ^b & Yu-fen Zhao ^a

^a Department of Chemistry, Key Laboratory of Chemical Biology and Organic Chemistry, Zhengzhou University, Zhengzhou, China

^b State Key Laboratory of Elemento-Organic Chemistry, Institute of Elemento-Organic Chemistry, Nankai University, Tianjin, China

Published online: 19 Dec 2008.

To cite this article: Yong Ye, Li-bo Ma, Yong Luo, Wen-hu Wang, Lun-zu Liu & Yu-fen Zhao (2008) Regioselective Cycloadditions of β -Substituted Vinylphosphonate with Nitrile Oxides, *Phosphorus, Sulfur, and Silicon and the Related Elements*, 184:1, 135-140, DOI: [10.1080/10426500802080717](https://doi.org/10.1080/10426500802080717)

To link to this article: <http://dx.doi.org/10.1080/10426500802080717>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and

are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

Regioselective Cycloadditions of β -Substituted Vinylphosphonate with Nitrile Oxides

Yong Ye,^{1,2} Li-bo Ma,^{1,2} Yong Luo,² Wen-hu Wang,²
Lun-zu Liu,² and Yu-fen Zhao¹

¹Department of Chemistry, Key Laboratory of Chemical Biology and Organic Chemistry, Zhengzhou University, Zhengzhou, China

²State Key Laboratory of Elemento-Organic Chemistry, Institute of Elemento-Organic Chemistry, Nankai University, Tianjin, China

Vinylphosphonates have been widely utilized in organic synthesis during the last two decades. In this article, a novel, highly regioselective cycloaddition utilizing nitrile oxides and vinylphosphonate for the synthesis of isoxazoline and fused isoxazoline rings was developed. The method has the advantages in that it tolerates a wide variety of functional groups.

Keywords 1,3-Dipolar cycloadditions; isoxazoline; vinylphosphonates

INTRODUCTION

Vinylphosphonates have been widely utilized in organic synthesis during the last two decades and became very useful for the construction of functionalized organophosphorus compounds.^{1–2} In particular, cycloaddition of nitrile oxides to vinylphosphonates is of considerable interest, as the resulting isoxazolines are versatile intermediates in the synthesis of a variety of natural products.^{3–4} Furthermore, the phosphonate functional group has been thought to be usually stable under the various transformations of isoxazolines including their reductive N–O bond cleavage.⁵

Recently, we have discovered that 3-diethylphosphonoethylene and vinylphosphonate substituted at the α -position by phosphonyl or carbonyl groups react with nitrile oxides to afford only one regioselective

Received 29 January 2008; 19 February 2008.

We thank the National Natural Science Foundation of China (No. 20602032), Education Department of Henan Province, and Zhengzhou University.


Address correspondence to Yong Ye, Department of Chemistry, Key Laboratory of Chemical Biology and Organic Chemistry, Zhengzhou University, Zhengzhou, 450052, China. E-mail: yeyong03@mail.nankai.edu.cn

product, which is a valuable intermediate of the synthesis of variously functionalized phosphonates.^{6,7} As a continuation of this study, we report here the behavior of β -substituted vinylphosphonate with nitrile oxides. The reaction has been explored with various substituted nitrile oxide to achieve information on the influence of the substituent at the β -position upon the regioselectivity and reactivity of the cycloaddition.

RESULTS AND DISCUSSION

The reaction of β -substituted vinylphosphonate with benzonitrile oxide generated in situ from benzohydroximoyl chloride and triethylamine in THF occurred smoothly at room temperature to afford the products **3** in good yield (shown in Scheme 1). The cycloadducts **3** characterized by elemental analysis and ^1H NMR spectra, in part also by ^{31}P NMR. Their related data are shown in Table I. Using the optimized conditions, cycloaddition of vinylphosphonate with phenyl substituted was faster than that of alkyl substituted. Reaction time was 12–18 h with **3e–g** and 24–48 h with **3a–d**.

β -Substituted vinylphosphonate undergoes a highly regioselective cycloaddition with nitrile oxide. The regioisomer was not detected by TLC and isolated by column chromatography. The structures of the adducts **3** rely upon the data of NMR. The regiochemistry follows from the comparison of the spacings between the isoxazolinic hydrogens. Previous NMR studies proved this technique to be useful for structural assignment of isoxazoline derivatives,^{8,9} especially in the case of trans-isomers. A proton in the 5-position absorbs lower field than a 4-proton, owing to the paramagnetic shift caused by the adjacent oxygen atom. In the case of the isoxazoline, $\Delta\delta_{5,4}$ ($\delta_5 - \delta_4$) is always notably larger for the 4-isoxazoline (1.4–2 ppm) than for the 5-isomers (1 ppm).⁶ Because of the higher deshielding effect of a relative to a phosphonyl substituent,



SCHEME 1 Cycloaddition of β -substituted vinylphosphonate with aryl nitrile oxides.

CONCLUSION

In conclusion, we have developed a novel 1,3-cyclic addition reaction utilizing nitrile oxides and vinylphosphonate for the synthesis of 4-phosphonyl isoxazoline and fused isoxazoline rings. The method has advantages in that it tolerates a wide variety of functional groups, is straightforward, and provides good yields of products from readily accessible starting materials. Further study is underway to expand the scope of this methodology, as well as to ascertain mechanistic details of the cycloaddition process.

EXPERIMENTAL

General

All melting points are uncorrected. Elemental analyses were carried on a Yanaco CHN Corder MT-3 apparatus. ^1H and ^{31}P NMR spectra were measured by using a Bruker 300 spectrometer with TMS and 85% H_3PO_4 as the internal and external reference respectively and with CDCl_3 as the solvent. Solvents used were purified and dried by standard procedures. Compounds **1** were synthesized according to Kiddle and Babler¹¹ and Monique *et al.*¹² Benzonitrile oxides were prepared efficiently from the corresponding hydroximoyl chlorides via base induced dehydrochlorination.¹³

General Procedure for the Synthesis of 3a–g

To a stirred solution of compounds **1** (2.0 mmol) and hydroxamic chlorides (2.2 mmol) in dry THF (15 mL) under N_2 , a solution of Et_3N (0.22 g, 2.2 mmol) in dry THF (15 mL) was added dropwise at -10° . The mixture was stirred at room temperature until the consumption of vinylphosphonates, monitored by TLC. Then, the reaction mixture was filtered to remove triethylamine hydrochloride, and the solvent was evaporated in vacuum. The residue was chromatographed on a silica gel column with petroleum ether/ethyl acetate 3:1(V:V) to give pure **3a–g** as a solid or oil.

3a: ^1H NMR: δ 0.85 (t, 3H); 1.14–1.80 (m, 16H); **3.60** (dd, 1H, $J = 4.16, 19.8$ Hz); 3.87–4.17 (m, 4H); **5.04** (m, 1H); 7.15 (m, 2H); 7.81 (m, 2H); ^{31}P NMR: 19.81; anal. calcd. for $\text{C}_{19}\text{H}_{29}\text{FNO}_4\text{P}$: C, 59.21; H, 7.58; N, 3.63. Found C, 58.91; H, 7.68, N, 3.73.

3b: ^1H NMR: δ 0.84 (t, 3H); 1.13–1.79 (m, 16H); **3.70** (dd, 1H, $J = 4.06, 19.6$ Hz); 3.9–4.2 (m, 4H); **5.1** (m, 1H); 7.33 (d, 2H, $J = 8.84$ Hz); 7.75 (d, 2H, $J = 8.84$ Hz); anal. calcd. for $\text{C}_{19}\text{H}_{29}\text{ClNO}_4\text{P}$: C, 56.79; H, 7.27; N, 3.49. Found C, 56.75; H, 7.17, N, 3.52.

3c: $^1\text{H NMR}$: δ 0.86 (t, 3H); 1.12–1.82 (m, 16H); 2.33 (s, 3H); **3.61** (dd, 1H, $J = 4.12, 19.7$ Hz); 3.86–4.16 (m, 4H); **5.06** (m, 1H); 7.20 (d, 2H, $J = 8.12$ Hz); 7.68 (d, 2H, $J = 8.12$ Hz); anal. calcd. for $\text{C}_{20}\text{H}_{32}\text{NO}_4\text{P}$: C, 62.98; H, 8.45; N, 3.67. Found C, 63.02; H, 8.20, N, 3.76.

3d: $^1\text{H NMR}$: δ 0.85 (t, 3H); 1.12–1.79 (m, 16H); **3.71** (dd, 1H, $J = 4.14, 19.82$ Hz); 3.89–4.2 (m, 4H); **5.1** (m, 1H); 7.37 (m, 3H); 7.79 (m, 2H); anal. calcd. for $\text{C}_{19}\text{H}_{30}\text{NO}_4\text{P}$: C, 62.11; H, 8.23; N, 3.81. Found C, 62.03; H, 8.17, N, 3.92.

3e: $^1\text{H NMR}$: δ 1.17–1.31 (m, 6H); 3.92–4.15 (m, 5H); **6.09** (dd, 1H, $J = 5.22, 20.84$ Hz); 7.23–7.38 (m, 6H); 7.77–7.80 (m, 4H); anal. calcd. for $\text{C}_{19}\text{H}_{22}\text{NO}_4\text{P}$: C, 63.50; H, 6.17; N, 3.90. Found C, 63.31; H, 6.04, N, 3.94.

3f: $^1\text{H NMR}$: δ 1.13–1.24 (m, 6H); 2.25 (s, 3H); 3.91–4.14 (m, 5H); **5.98** (dd, 1H, $J = 4.18, 20.86$ Hz); 7.08 (d, 2H, $J = 8.34$ Hz); 7.60 (d, 2H, $J = 8.34$ Hz); anal. calcd. for $\text{C}_{20}\text{H}_{24}\text{NO}_4\text{P}$: C, 64.33; H, 6.48; N, 3.75. Found C, 64.36; H, 6.25, N, 3.53.

3g: yellow solid, mp 144–146°. $^1\text{H NMR}$: δ 1.21–1.37 (m, 6H); **3.93** (dd, 1H, $J = 5.22, 19.82$ Hz); 4.03–4.18 (m, 4H); **6.23** (dd, 1H, $J = 5.22, 21.9$ Hz); 7.53 (d, 2H, $J = 8.83$ Hz); 7.98 (d, 2H, $J = 9.04$ Hz); 8.23 (d, 4H, $J = 8.8$ Hz); anal. calcd. for $\text{C}_{19}\text{H}_{20}\text{N}_3\text{O}_8\text{P}$: C, 50.79; H, 4.49; N, 9.35. Found C, 50.94; H, 4.30, N, 9.24. HRMS (ESI): m/z calcd. for $\text{C}_{19}\text{H}_{20}\text{N}_3\text{O}_8\text{P}$: 449.0988; found: 449.0995.

General Procedure for the Synthesis of 5

To A stirred solution of phosphonyl hydroximoyl bromides (0.432 g, 1.5 mmol) and β -substituted vinylphosphonate (15 mmol) in dry ether (10 mL), a solution of Et_3N (0.2 g, 2 mmol) in dry ether (10 mL) was added dropwise at -10° . After half an hour, the solution ascends to ambient temperature. The reaction mixture was stirred continue for three days. After filtered, the solvent was evaporated in vacuum. The residue was chromatographed on a silica gel column with petroleum ether/ethyl acetate 1:3 to give pure **5a–c** as an oil.

5a: yield 19.6%, $^1\text{H NMR}$: δ 7.32 (m, 5H), 5.89–6.03 (dd, 1H), 4.69(m,4H), 4.12–4.22 (m, 5H), 1.20–1.39 (m, 30H), $^{31}\text{P NMR}$: δ 0.89 ($^\alpha\text{P}$), -3.25 ($^\beta\text{P}$), 18.53 ($^\gamma\text{P}$); anal. calcd. for it, found(calcd): C47.72 (47.71), H6.79 (6.93), N 4.29(4.28).

5b: yield 31.4%, $^1\text{H NMR}$: δ 4.65–4.75 (m, 5H), 3.99–4.10 (m, 5H), 1.08–1.41 (m, 40H), 0.73–0.76 (m, 3H), $^{31}\text{P NMR}$: δ 1.08 ($^\alpha\text{P}$), -3.14 ($^\beta\text{P}$), 19.2 ($^\gamma\text{P}$); anal. calcd. for it, found(calcd): C46.98 (47.13) 8.13(8.06) 4.28(4.23).

5c: yield 56.1%, $^1\text{H NMR}$: δ 8.19–8.23 (d, 2H), 7.51–7.55 (d, 2H), 6.06 (dd, 1H), 4.83 (m, 4H), 4.21 (m, 5H), 1.23–1.37 (m, 30H), $^{31}\text{P NMR}$: δ 0.53

(α P), -3.32 (β P), 17.68 (ν P); anal. calcd. for it, found (calcd): $44.59(44.64)$
 $6.35(6.34)$ $5.98(6.01)$.

REFERENCES

- [1] For review, see: (a) T. Ageno, T. Okauchi, T. Minami, and M. Ishida, *Organic & Biomolecular Chemistry*, **3**(5), 924 (2005); (b) T. Minami, T. Okauchi, and R. Kouno, *Synthesis*, **3**, 349 (2001).
- [2] For applications: (a) Z. A. Doddridge, R. D. Bertram, and C. J. Hayes, *Biochemistry*, **42**(11), 3239 (2003); (b) W. Schwab, R. Bartlett, U. Gebert, H. U. Schorlemmer, G. Dickneite, and H. H. Sedlacek, *Ger. Offen.*, DE 3736113 (1989); (c) C. Yuan and C. Li, *Phosphorus, Sulfur, and Silicon*, **103**(1–4), 133, (1995).
- [3] For review: (a) A. P. Kozikowski, *Acc. Chem. Res.*, **17**, 410 (1984); (b) C. Grundmann and P. Grunanger, *The nitrile oxides* (Springer, Berlin, 1971). (c) A. Hassner, K. S. K. Murthy, R. Maurya, W. Dehaen, and O. Friedman, *J. Heterocycl. Chem.*, **31**, 687 (1994).
- [4] For applications to natural product synthesis: (a) G. Brieger and J. N. Bennet, *Chem. Rev.*, **80**, 63 (1980); (b) G. Broggin and G. Zecchi, *Synthesis*, 905 (1999). (c) A. I. Kotyatkina, V. N. Zhabinsky, and V. A. Khripach, *Russ. Chem. Rev.*, **70**, 641 (2001).
- [5] H.-J. Gi, Y. Xiang, R. F. Schinazi, and K. Zhao, *J. Org. Chem.*, **62**, 88 (1997).
- [6] Y. Ye, Y. Zheng, G. Xu, and L. Liu, *Heteroatom Chemistry*, **14**(3), 254 (2003).
- [7] Y. Ye, G. Xu, Y. Zheng, and L. Liu, *Heteroatom Chemistry*, **14**(4), 309 (2003).
- [8] G. Bianchi, R. Gandolfi, P. Grunanger, and A. Perotti, *J. Chem. Soc. (C)*, 1598 (1967).
- [9] R. Sustmann, R. Huisgen, and H. Huber, *Chem. Ber.*, **100**, 1802 (1967).
- [10] (a) K. N. Anisimov and A. N. Nesmeyanov, *Izvestiya Akademii Nauk SSSR, Seriya Khimicheskaya*, 610 (1954); (b) K. N. Anisimov, N. E. Kolobova, and A. N. Nesmeyanov, *Izvestiya Akademii Nauk SSSR, Seriya Khimicheskaya*, 240 (1955); (c) R. R. Shagidullin, V. A. Pavlov, B. I. Buzykin, N. V. Aristova, and L. F. Chertanova, *Zh. Obshch. Khim.*, **61**, 1590 (1991).
- [11] J. J. Kiddle and J. H. Babler, *J. Org. Chem.*, **58**, 3572 (1993).
- [12] R. Monique, D. V. Anthony, and V. Jean, *Synth. Commun.*, **14**, 833 (1984).
- [13] G. W. Perold, A. P. Steyn, and F. V. K. Reiche, *J. Am. Chem. Soc.*, **79**, 462 (1957).