

Reaction between Conjugated Azoalkenes and Pyrazolinones: A Precious Entry to New Conjugated Azodiene and Asymmetric 4,4'-Bipyrazole Derivatives

Orazio A. Attanasi, Paolino Filippone,* Chiara Fiorucci, and Fabio Mantellini
Istituto di Chimica Organica, Università di Urbino, Piazza della Repubblica 13, I-61029 Urbino, Italy

(Received May 8, 2000; CL-000444)


The first direct nucleophilic addition under mild reaction conditions of heterocyclic compounds to conjugated azoalkenes is reported. This reaction readily gives rise to new conjugated azodiene or asymmetric 4,4'-bipyrazole derivatives by means of two independent pathways.

In the wide scenarios of the azo-ene systems,¹⁻⁵ conjugated azodienes are relatively little investigated compounds, in spite of their potential usefulness in organic synthesis.¹⁻⁶ Furthermore, asymmetric 4,4'-bipyrazole derivatives are difficult to prepare by current synthetic methods reported in the literature.⁷

We now report the synthesis of new conjugated azodienes **4a-d** or asymmetric 4,4'-bipyrazol-4,4'-in-3,3'-ones **5a-b** by unprecedented¹⁻⁵ direct nucleophilic 1,4-addition of 1-methyl-3-*n*-propyl-2-pyrazolin-5-one **2a** or 1-phenyl-3-methyl-2-pyrazolin-5-one **2b** to the azo-ene system of conjugated azoalkenes **1a-c** in the presence of strong base anion exchanger resin Duolite® A102 (Rohm and Haas). This reaction affords firstly the intermediate CH-hydrazone adducts and then NH-hydrazone forms, as shown by the disappearance in ¹H-NMR of the CH (4.37) signal and the appearance of a new NH (10.32) signal during few hours in the NMR tube.^{4d,8}

Subsequently, the above-mentioned products were obtained by two independent pathways. In fact, conjugated azodienes **4** having one of the two conjugated olefinic double bonds in the heterocycle ring arise by mild oxidation of the hydrazone derivatives **3** with selenium dioxide at room temperature. Asymmetric 4,4'-bipyrazol-4,4'-in-3,3'-ones derive by simple treatment of the same hydrazone adducts in methanol under reflux by means of the intramolecular cyclization of the hydrazone side-chain in position 4 of the pyrazolinone ring. The solvolytic cleavage of N-COR¹ bond was also observed (see Scheme 1 and Table 1).⁸

In conclusion we report here the first direct nucleophilic addition of heterocyclic compounds to the heterodiene system of conjugated azoalkenes. While the reaction of conjugated azoalkenes with activated methylene and methine compounds was extensively studied showing they are powerful tools in organic synthesis,²⁻⁵ the present findings open a broad field of



Scheme 1. Reagents and conditions: i) Duolite® A102, THF, room temp.; ii) SeO₂, THF, room temp.; iii) MeOH, reflux.

Table 1. Yields of products **3a-d**, **4a-d** and **5a,b**

1	2	3	Yield/% ^a	4	Yield/% ^a	5	Yield/% ^a
1a	2a	3a	75	4a	90	5a	81
1a	2b	3b	68	4b	81		
1b	2b	3c	71	4c	88		
1c	2b	3d	78	4d	92	5b	84

^a Yields of isolated products.

further innovative investigations on the chemistry of these versatile intermediates. Indeed, this reaction represents an unknown, mild and useful entry to a new class of conjugated azodienes in which one of two ene functions is located in the pyrazole ring. Moreover, the preparation of asymmetric 4,4'-bipyrazole derivatives difficult to obtain by other methods is also reported.

This work was supported by financial assistance from the Università degli Studi di Urbino and the Consiglio Nazionale delle Ricerche (C.N.R.-Roma).

References and Notes

- 1 O. A. Attanasi and L. Caglioti, *Org. Prep. Proced. Int.*, **18**, 299 (1986).
- 2 J. G. Schantl, "1-Azo-1-alkene" in "Houben-Weyl", 4th ed, ed. by H. Kropf and E. Schaumann, Thieme, Stuttgart (1990), Vol. E15, p. 909; J. G. Schantl, *Farmaco*, **50**, 379 (1995); J. G. Schantl, *Molecules*, **1**, 212 (1996).
- 3 K. Banert, "Synthesis of five-membered heterocycles from novel functionalized allenes" in "Targets in Heterocyclic Systems – Chemistry and Properties", ed. by O. A. Attanasi and D. Spinelli, Società Chimica Italiana, Roma (2000), Vol. 3, Chap. 1, p. 1; S. Polanc, "Recent applications of hydrazides and related compounds for the synthesis of heterocycles" in "Targets in Heterocyclic Systems – Chemistry and Properties", ed. by O. A. Attanasi and D. Spinelli, Società Chimica Italiana, Roma (2000), Vol. 3, Chap. 2, p. 33.
- 4 a) O. A. Attanasi and P. Filippone *Synlett*, **1997**, 1128; b) A. Arcadi, O. A. Attanasi, L. De Crescentini, and E. Rossi, *Tetrahedron Lett.*, **38**, 2329 (1997); c) O. A. Attanasi, P. Filippone, and C. Fiorucci, *Tetrahedron*, **53**, 1467 (1997); d) O. A. Attanasi, L. De Crescentini, P. Filippone, E. Foresti, R. Galeazzi, I. Ghiviriga, and A. R. Katritzky, *Tetrahedron*, **53**, 5617 (1997); e) O. A. Attanasi, L. De Crescentini, E. Foresti, G. Gatti, R. Giorgi, F. R. Perrulli, and S. Santeusano, *J. Chem. Soc., Perkin Trans. 1*, **1997**, 1829; f) O. A. Attanasi, P. Filippone, C. Fiorucci, and F. Mantellini, *Synlett*, **1997**, 1361; g) O. A. Attanasi, P. Filippone, C. Fiorucci, E. Foresti, and F. Mantellini, *J. Org. Chem.*, **63**, 9880 (1998); h) A. Arcadi, O. A. Attanasi, B. Guidi, E. Rossi, and S. Santeusano, *Chem. Lett.*, **1999**, 59; i) O. A. Attanasi, L. De Crescentini, P. Filippone, F. R. Perrulli, and S. Santeusano, *Synlett*, **1999**, 339; j) O. A. Attanasi, P. Filippone, C. Fiorucci, and F. Mantellini, *Tetrahedron Lett.*, **40**, 3891 (1999); k) O. A. Attanasi, L. De Crescentini, P. Filippone, B. Guidi, F. R. Perrulli, and S. Santeusano, *Synlett*, **1999**, 1367; l) A. Arcadi, O. A. Attanasi, B. Guidi, E. Rossi, and S. Santeusano, *Eur. J. Org. Chem.*, **1999**, 3117; m) O. A. Attanasi, P. Filippone, E. Foresti, B. Guidi, and S. Santeusano, *Tetrahedron*, **55**, 13423 (1999); n) O. A. Attanasi, P. Filippone, B. Guidi, F. R. Perrulli, and S. Santeusano, *Heterocycles*, **51**, 2423 (1999). o) O. A. Attanasi, P. Filippone, B. Guidi, T. Hippe, F. Mantellini, and L. F. Tietze, *Tetrahedron Lett.*, **40**, 9277 (1999); p) O. A. Attanasi, R. Ballini, L. De Crescentini, P. Filippone, and F. Mantellini, *J. Org. Chem.*, **64**, 9653 (1999); q) O. A. Attanasi, L. De Crescentini, P. Filippone, E. Foresti, and F. Mantellini, *J. Org. Chem.*, **65**, 2820 (2000).
- 5 G. Ferguson, A. J. Lough, D. Mackay, and G. Weeratunga, *J. Chem. Soc., Perkin Trans. 1*, **1991**, 3361; M. Avalos, R. Babiano, P. Cintas, J. L. Jiménez, M. M. Molina, J. C. Palacios, and J. B. Sánchez, *Tetrahedron Lett.*, **32**, 2513 (1991); T. L. Gilchrist and A. Lemos, *J. Chem. Soc., Perkin Trans. 1*, **1993**, 1391; M. Avalos, R. Babiano, P. Cintas, J. L. Jiménez, J. C. Palacios, and J. B. Sánchez, *Tetrahedron: Asymm.*, **6**, 945 (1995); M. S. South, T. L. Jakuboski, M. D. Westmeyer, and D. R. Dukeshner, *J. Org. Chem.*, **61**, 8921 (1996); V. Buß, and L. Eggers, *Tetrahedron: Asymm.*, **8**, 1531 (1997); Z. V. Todres and G. Ts. Hovsepyan, *J. Chem. Soc., Perkin Trans. 1*, **1997**, 747; C. Joergensen and C. Pedersen, *Carbohydr. Res.*, **299**, 307 (1997); L. J. O. Figueiredo and C. Kasheres, *J. Org. Chem.*, **62**, 1164 (1997); N. Caposciali, C. Dell'Erba, M. Novi, G. Petrillo and C. Tavani, *Tetrahedron*, **54**, 5315 (1998); F. Palacios, D. Aparicio, and J. M. de los Santos, *Tetrahedron*, **55**, 13767 (1999).
- 6 W. Barbieri, L. Caglioti, F. Gasparrini, and G. Rosini, *Chim. Ind. (Milan)*, **52**, 267 (1970).
- 7 J. Elguero, "Pyrazoles" in "Comprehensive Heterocyclic Chemistry II", ed. by A. R. Katritzky, C. W. Rees, and E. F. V. Scriven, Pergamon-Elsevier Science, Amsterdam (1996), Vol. 3, Chap. 1; "Heterocyclic Chemistry", ed. by H. Suschitzky and O. Meth-Cohn, Royal Society of Chemistry, London (1978-1985), Vol. 1-4; "Progress in Heterocyclic Chemistry", ed. by H. Suschitzky and E. F. V. Scriven, Pergamon Press, Oxford (1989-1995), Vol. 1-7; "Advances in Heterocyclic Chemistry", ed. by A. R. Katritzky, Academic Press, New York (1963-1994), Vol. 1-7; "Progress in Heterocyclic Chemistry", ed. by H. Suschitzky and G. W. Gribble, Pergamon Press, Oxford (1996), Vol. 8; "Progress in Heterocyclic Chemistry", ed. by G. W. Gribble and T. L. Gilchrist, Pergamon Press, Oxford (1997), Vol. 9; "Progress in Heterocyclic Chemistry", ed. by G. W. Gribble and T. L. Gilchrist, Pergamon Press, Oxford (1998), Vol. 10.
- 8 Typical experimental procedure for the synthesis of 1,4-adducts **3a-d**: To a stirred solution of conjugated azoalkenes **1a-c** (1 mmol) in THF (20 ml) was added 1-methyl-3-*n*-propyl-2-pyrazolin-5-one **2a** or 1-phenyl-3-methyl-2-pyrazolin-5-one **2b** (1 mmol) and Duolite® A102 (400 mg). The suspension was allowed to react at room temperature for 1 hr under magnetic stirring. The product was subsequently isolated by column chromatography on silica gel with cyclohexane-ethyl acetate mixtures to give 1,4-adducts **3a-d**. **3d**: mp 118-120 °C; ¹H-NMR (CDCl₃): 1.51 (s, 9 H, CMe₃), 2.06 (s, 3 H, Me), 2.24 (s, 3 H, Me), 3.72 (s, 3 H, OMe), 4.37 (s, 1 H, CH), 7.10-7.81 (m, 5 H, Ph), 8.21 (bs, 1 H, NH, D₂O exch.), 10.96 (bs, 1 H, NH, D₂O exch.); ¹³C-NMR (CDCl₃): 12.6, 17.2, 26.1, 52.9, 82.8, 93.5, 119.3, 121.7, 126.5, 130.7, 138.6, 147.3, 150.1, 151.5, 153.0, 170.7; MS *m/z* (relative intensity) 402 (M⁺, 14), 346 (74), 329 (9), 302 (15), 287 (22), 270 (100). Anal. Calcd for C₂₀H₂₆O₅N₄: C, 59.69; H, 6.51; N, 13.92; Found: C, 59.71; H, 6.50; N, 13.91. Typical experimental procedure for the synthesis of conjugated azodienes **4a-d**: To a stirred solution of 1,4-adducts **3a-d** (1 mmol) in THF (20 ml) was added selenium dioxide (1 mmol). The suspension was allowed to react at room temperature for 15 min under magnetic stirring. The products were subsequently isolated by column chromatography on silica gel with cyclohexane-ethyl acetate mixtures to give conjugated azodienes **4a-d**. **4d**: mp 152-154 °C; ¹H-NMR (CDCl₃): 1.54 (s, 9 H, CMe₃), 2.03 (s, 3 H, Me), 2.26 (s, 3 H, Me), 3.96 (s, 3 H, OMe), 7.09-7.92 (m, 5 H, Ph), 8.67 (b s, 1 H, NH, D₂O exch.); ¹³C-NMR (CDCl₃): 13.2, 17.9, 28.1, 53.3, 82.6, 116.7, 125.0, 127.0, 128.8, 137.6, 139.8, 144.7, 146.7, 152.2, 161.9, 166.4; MS *m/z* (relative intensity) 400 (M⁺, 12), 344 (13), 300 (11), 295 (14), 272 (39), 259 (82), 240 (100). Anal. Calcd for C₂₀H₂₄O₅N₄: C, 59.99; H, 6.04; N, 13.99; Found: C, 59.97; H, 6.05; N, 14.00. Typical experimental procedure for the synthesis of 4,4'-bipyrazol-4,4'-in-3,3'-ones **5a,b**: 1,4-Adduct **3a-d** (1 mmol) was dissolved in methanol (25 ml) and heated under reflux for 3 hr. The product was subsequently isolated by column chromatography on silica gel with ethyl acetate to give 4,4'-bipyrazol-5,4'-in-5,3'-ones **5a,b**. **5b**: mp 284-286 °C; ¹H-NMR (DMSO): 2.09 (s, 6 H, 2 Me), 7.10-7.81 (m, 5 H, Ph), 10.92 (b s, 3 H, 3 NH, D₂O exch.); ¹³C-NMR (DMSO): 11.3, 13.0, 93.9, 119.6, 124.7, 128.9, 138.2, 138.5, 147.3, 159.8; MS *m/z* (relative intensity) 270 (M⁺, 100). Anal. Calcd for C₁₄H₁₄O₂N₄: C, 62.21; H, 5.22; N, 11.84; Found: C, 62.19; H, 5.23, N, 11.85.