

# SnX<sub>2</sub>/Al-promoted formylation of per(poly)fluoroalkyl halides with *N,N*-dimethylformamide

Chang-Ming Hu\* and Jian Chen

Shanghai Institute of Organic Chemistry, Academia Sinica, 345 Lingling Lu, Shanghai 200032 (China)

(Received April 20, 1993; accepted June 15, 1993)

## Abstract

In the presence of catalytic amounts of stannous salts (20 mol%) and aluminium powder, per(poly)fluoroalkyl iodides or bromides react smoothly with DMF under mild conditions to give the corresponding per(poly)fluoroalkyl aldehydes in good to excellent yield. A tin-containing intermediate was obtained. A possible mechanism is proposed.

## Introduction


Per(poly)fluoroalkyl aldehydes and their monohydrates are useful intermediates [1]. Several classical methods have been used to synthesize such compounds [2–4] but all suffer from poor yields or tedious procedures. Recently, it was reported that some initiators could effectively promote the formylation of per(poly)fluoroalkyl halides with DMF to give the corresponding per(poly)fluoroalkyl aldehydes [5].

Stannous salts have been used extensively as Barbier-type allylation reagents for carbonyl compounds [6]. In the presence of a stoichiometric amount of SnCl<sub>2</sub>, Kitazume and Ishikawa reported the per(poly)fluoroalkylation of aldehydes and ketones with R<sub>f</sub>I [7]. In our continuing studies on redox couple-promoted reactions in organofluorine chemistry, we found that a bimetal redox couple composed of a catalytic amount of an anhydrous stannous salt and aluminium powder was sufficiently active to promote the formylation of R<sub>f</sub>I or R<sub>f</sub>Br with DMF. A possible mechanism involving an adduct intermediate is proposed.

## Results and discussion

In the presence of catalytic amounts of anhydrous stannous chloride and aluminium powder, per(poly)fluoroalkyl halides (iodides **1** or bromides **2**) react readily with DMF to give the formylation products R<sub>f</sub>CH(OH)<sub>2</sub> **4** in good yield. These per(poly)fluoroalkylaldehyde monohydrates (**4**) were dehydrated

with phosphorus pentoxide at 60 °C to give the corresponding anhydrous aldehydes **5** in nearly quantitative yield. The yields and reaction conditions are listed in Table 1.



(1a) ClC<sub>4</sub>F<sub>8</sub>I; (1b) ClC<sub>6</sub>F<sub>12</sub>I; (1c) ClC<sub>8</sub>F<sub>16</sub>I; (1d) C<sub>4</sub>F<sub>9</sub>I;  
 (2a) C<sub>6</sub>F<sub>13</sub>Br; (2b) C<sub>8</sub>F<sub>17</sub>Br; (3) CH<sub>2</sub>=CHC<sub>6</sub>F<sub>12</sub>Cl

a. SnCl<sub>2</sub> (20 mol%), 1 equiv. Al, r.t., 10 h; b. 2.5 equiv. P<sub>2</sub>O<sub>5</sub>, 60 °C.

Scheme 1.

Under such conditions, the reaction proceeded smoothly at room temperature within 10 h. The corresponding per(poly)fluoroalkyl aldehyde monohydrates **4** were obtained in good yield. Perfluoroalkyl bromides gave similar results, except that a longer reaction time was necessary. The main by-product was R<sub>f</sub>H which amounted to less than 10%. However, perfluoroalkyl chlorides **3** remained intact under these or even more drastic conditions (Run 8). The reactivity of per(poly)fluoroalkyl halides decreased in the order: R<sub>f</sub>I > R<sub>f</sub>Br > R<sub>f</sub>Cl.

Polyfluoroalkyl halides such as 2-chlorotetrafluoroethyl iodide containing vicinal dihalogens were dehalogenated to give tetrafluoroethene (Run 5). In fact, SnCl<sub>2</sub> on its own was sufficiently active to eliminate halogens from vicinal dihalogen compounds.

\*Author to whom correspondence should be addressed.

TABLE 1. Tin(II)/Al-promoted formylation of R<sub>f</sub>X with DMF<sup>a</sup>


Run No.	R <sub>f</sub> X	SnY <sub>2</sub>	Temp./time (°C/h)	Product	Yield <sup>b</sup> (%)
1	Cl(CF <sub>2</sub> ) <sub>4</sub> I	SnCl <sub>2</sub>	16/6	Cl(CF <sub>2</sub> ) <sub>4</sub> CHO	80
2	Cl(CF <sub>2</sub> ) <sub>6</sub> I	SnCl <sub>2</sub>	15/8	Cl(CF <sub>2</sub> ) <sub>6</sub> CHO	97
3	Cl(CF <sub>2</sub> ) <sub>8</sub> I	SnCl <sub>2</sub>	15/10	Cl(CF <sub>2</sub> ) <sub>8</sub> CHO	88
4	F(CF <sub>2</sub> ) <sub>4</sub> I	SnCl <sub>2</sub>	15/8	F(CF <sub>2</sub> ) <sub>4</sub> CHO	90
5	Cl(CF <sub>2</sub> ) <sub>2</sub> I	SnCl <sub>2</sub>	15	no product	<sup>c</sup>
6	F(CF <sub>2</sub> ) <sub>8</sub> Br	SnCl <sub>2</sub>	15/12	F(CF <sub>2</sub> ) <sub>8</sub> CHO	89
7	F(CF <sub>2</sub> ) <sub>6</sub> Br	SnCl <sub>2</sub>	15/12	F(CF <sub>2</sub> ) <sub>6</sub> CHO	84
8	CH <sub>2</sub> =CH(CF <sub>2</sub> ) <sub>6</sub> Cl	SnCl <sub>2</sub>	60/10	no reaction	<sup>d</sup>
9	Cl(CF <sub>2</sub> ) <sub>4</sub> I	SnBr <sub>2</sub>	15/6	Cl(CF <sub>2</sub> ) <sub>4</sub> CHO	83
10	Cl(CF <sub>2</sub> ) <sub>6</sub> I	SnBr <sub>2</sub>	15/9	Cl(CF <sub>2</sub> ) <sub>6</sub> CHO	75
11	Cl(CF <sub>2</sub> ) <sub>8</sub> I	SnBr <sub>2</sub>	15/10	Cl(CF <sub>2</sub> ) <sub>8</sub> CHO	79
12	F(CF <sub>2</sub> ) <sub>8</sub> Br	SnBr <sub>2</sub>	15/15	F(CF <sub>2</sub> ) <sub>8</sub> CHO	74
13	Cl(CF <sub>2</sub> ) <sub>6</sub> I	SnF <sub>2</sub>	15/5	Cl(CF <sub>2</sub> ) <sub>6</sub> CHO	85
14	F(CF <sub>2</sub> ) <sub>8</sub> Br	SnF <sub>2</sub>	15/6	F(CF <sub>2</sub> ) <sub>8</sub> CHO	90
15	Cl(CF <sub>2</sub> ) <sub>6</sub> I	SnI <sub>2</sub>	15/10	no reaction	<sup>d</sup>

<sup>a</sup>All compounds had similar boiling or melting points, and <sup>1</sup>H NMR, <sup>19</sup>F NMR, MS and IR spectra identical to those described in the literature [5].

<sup>b</sup>Isolated yields based on R<sub>f</sub>X.


<sup>c</sup>CF<sub>2</sub>=CF<sub>2</sub> was obtained.

<sup>d</sup>Starting material was completely recovered.




The solvent seemed to play quite an important role. DMF could be used both as a reactant and a solvent. DMSO could also be used as a solvent without significant difference. However, in acetonitrile, THF or protic solvents, only R<sub>f</sub>H was obtained.

It was shown that catalytic amounts of both stannous chloride and aluminium powder were necessary because stoichiometric amounts of SnCl<sub>2</sub> or of aluminium on their own were unable to promote the reaction. On the other hand, stannous chloride could be replaced by other stannous salts. For example, SnF<sub>2</sub> was more and SnBr<sub>2</sub> less effective than SnCl<sub>2</sub>, but SnI<sub>2</sub> would not promote the reaction at all. The catalytic activity of these stannous salts decreased in the order: SnF<sub>2</sub> > SnCl<sub>2</sub> > SnBr<sub>2</sub> > SnI<sub>2</sub>. Aluminium could not be replaced by other metals such as zinc, magnesium or iron.



Although stoichiometric amounts of SnCl<sub>2</sub> were incapable of promoting the formylation reaction, SnCl<sub>2</sub> did react with a per(poly)fluoroalkyl iodide to form an adduct. Thus, in dimethyl sulfoxide (DMSO), ClC<sub>4</sub>F<sub>8</sub>I **1a** reacted with SnCl<sub>2</sub> to give an adduct **5a** in c. 50% conversion. Although compound **5a** could not be isolated pure, its presence in DMSO was evident [7] since the <sup>19</sup>F NMR signal of CF<sub>2</sub>I was shifted upfield by c. 40 ppm and became a broad singlet. DMF was then added to this solution. It was found that in the absence of

aluminium powder the 4-chloroperfluorobutyltin(IV) halide **5a** remained intact. In contrast, when aluminium powder was added, the desired product and ClC<sub>4</sub>F<sub>8</sub>H were obtained.



Compound **5a** was sensitive to moisture giving a colourless solid on exposure in air, whose experimental formula corresponded to the stannonic acid ClC<sub>4</sub>F<sub>8</sub>SnO<sub>2</sub>H (**6**) (formula weight, 387) as determined by <sup>19</sup>F NMR, <sup>1</sup>H NMR and IR spectroscopy, and by microanalysis (see data listed in Table 2). Mass spectrometry showed isotope peaks of tin, but there were also peaks beyond *m/z* = 400, even up to *m/z* = 900. This indicates that stannonic acid **6** is trimerized under these conditions. In fact, once converted into stannonic acids stannic alkyl trihalides polymerized readily [8]. Solid **6** remained intact both in the presence and absence of aluminium powder.


All these results seem to imply that the reaction may proceed through the following pathway. Compound **1** reacts with SnCl<sub>2</sub> to form adduct **5**, which is further reduced by aluminium into **7**. Compound **7** reacts with DMF to form adduct **8** which gives product **4** on hydrolysis.

TABLE 2. IR, <sup>1</sup>H NMR, <sup>19</sup>F NMR and MS spectra of some per(poly)fluoroalkyl aldehyde monohydrates

Compounds	<sup>19</sup> F (ppm)	<sup>1</sup> H (ppm)	IR (cm <sup>-1</sup> )	MS (m/e %)
Cl(CF <sub>2</sub> ) <sub>4</sub> CH(OH) <sub>2</sub>	-8.5(s, 2F)	5.9(m, 1H)	3300(s)	265(0.84)
	43(s, 2F)	6.1(s, 2H)	1180(s)	131(100)
	45(s, 2F)		1160(s)	100(36.6)
	54(s, 2F)			85(31.6)
Cl(CF <sub>2</sub> ) <sub>6</sub> CH(OH) <sub>2</sub> <sup>a</sup>	-8(s, 2F)	5.8(m, 1H)	3300(s)	365(4.0)
	44(s, 2F)	6.4(s, 2H)	1180(s)	131(100)
	46(m, 6F)		1130(s)	
	52(s, 2F)			
	52(s, 2F)			
Cl(CF <sub>2</sub> ) <sub>4</sub> SnO <sub>2</sub> H <sup>b</sup>	-9(s, 2F)	3.8(m)	3400(vs)	389(M <sup>+</sup> + 1)
	39(m, 4F)		1610(s)	121,
	43(s, 2F)		1290(m)	120,
			1250(vs)	118,
				116(Sn <sup>+</sup> )

<sup>a</sup>Elemental analysis: Calc. for C<sub>7</sub>H<sub>3</sub>ClF<sub>12</sub>O<sub>2</sub>: C, 23.05; H, 0.30; F, 62.55%. Found: C, 23.23; H, 0.33; F, 62.46%.

<sup>b</sup>Elemental analysis: Calc. for C<sub>4</sub>HClF<sub>8</sub>O<sub>2</sub>Sn: C, 12.38; H, 0.26; Cl, 9.16%. Found: C, 12.02; H, 0.26; Cl, 8.69%.



The present work thus furnishes a readily available redox couple useful in the formylation of R<sub>f</sub>I or R<sub>f</sub>Br with DMF under mild conditions forming products in good to excellent yield. These aldehydes are useful starting materials in the synthesis of acids, alcohols or halides or relevant compounds containing an R<sub>f</sub> group.

## Experimental

All melting or boiling points are uncorrected. IR spectra were recorded on a Shimadzu IR-440 spectrometer. <sup>19</sup>F NMR spectra were recorded on a Varian EM-360L spectrometer in CCl<sub>4</sub> or CDCl<sub>3</sub> at 56.4 MHz using CF<sub>3</sub>CO<sub>2</sub>H as external standard. Chemical shifts in ppm were positive for upfield shifts. <sup>1</sup>H NMR spectra were recorded on a Varian EM-360A spectrometer at 60 MHz in CCl<sub>4</sub> or CDCl<sub>3</sub> using SiMe<sub>4</sub> as external standard. MS spectra were measured on a Finnigan MS-4021 spectrometer.

A typical procedure was as follows. To a vigorously stirred suspension consisting of 10–12 mmol of aluminium powder and 1 mmol (0.2 g) of SnCl<sub>2</sub> in 10 ml

of dry *N,N*-dimethylformamide (DMF) was added dropwise a solution consisting of 10 mmol of R<sub>f</sub>X (X = I, Br) in 1 ml of DMF. The mixture was stirred for several hours at room temperature. Then dilute aq. HCl was added and the residue filtered off. The filtrate was extracted three times with ethyl acetate. The combined organic extract was washed with water, brine and dried over Na<sub>2</sub>SO<sub>4</sub>. Removal of the solvent gave a pale yellow solid which was identified as per(poly)fluoroalkyl-aldehyde monohydrate. Dehydration with P<sub>2</sub>O<sub>5</sub> at 60–70 °C gave the corresponding aldehyde.

## Acknowledgement

We thank the National Science Foundation of China for financial support.

## References

- (a) D.J. Cook, O.R. Piece and E.T. McBee, *J. Am. Chem. Soc.*, **76** (1954) 83; (b) N. Ono, H. Kawamura and K. Maruyama, *Bull. Chem. Soc. Jpn.*, **62** (1980) 3386; (c) K.I. Pashkevich, R.R. Latypov and V.I. Filyakova, *Bull. Acad. Sci. USSR (Engl. Trans.)*, **35** (1986) 2361; (d) K. Tanaka, Y. Sugimoto, Y. Okafuji, M. Tachikawa and K. Mitsunashi, *J. Heterocycl. Chem.*, **26** (1989) 381.
- (a) D.R. Husted and A.H. Ahlbrecht, *J. Am. Chem. Soc.*, **74** (1952) 5422; (b) M. Braid, H. Iserson and F.E. Lawlor, *J. Am. Chem. Soc.*, **76** (1954) 4027; (c) F. Brown and W.K.R. Musgrave, *J. Chem. Soc.*, (1952) 5039; (d) A.L. Henne, R.L. Pelley and R.M. Alm, *J. Am. Chem. Soc.*, **72** (1950) 3371; (e) N.O. Brace, *J. Org. Chem.*, **26** (1961) 4005; (f) R.B. Greenwald and D.H. Evans, *J. Org. Chem.*, **41** (1976) 1470.

- 3 R.W. Lang, *Helv. Chim. Acta*, 71 (1988) 369.
- 4 S. Benefice-Malout, H. Blancou and A. Commeyras, *J. Fluorine Chem.*, 45 (1989) 87.
- 5 C.-M. Hu and X.-Q. Tang, *J. Fluorine Chem.*, 61 (1993) 217.
- 6 T. Mukaiyama, T. Hada and S. Shoda, *Chem. Lett.*, (1981) 723.
- 7 T. Kitazume and N. Ishikawa, *Chem. Lett.*, (1981) 1337.
- 8 E.W. Ebel, in J.C. Bailar (ed.), *Comprehensive Inorganic Chemistry*, Pergamon, Oxford, 1973, Vol. 2, p. 43.