### Accepted Manuscript

Iron-catalyzed oxidative dehydrogenative coupling of ethers with aryl tetrazoles

Kai-qiang Zhu, Liang Wang, Qun Chen, Ming-yang He

 PII:
 S0040-4039(15)01110-7

 DOI:
 http://dx.doi.org/10.1016/j.tetlet.2015.06.091

 Reference:
 TETL 46483

To appear in: Tetrahedron Letters

Received Date:25 May 2015Revised Date:23 June 2015Accepted Date:29 June 2015



Please cite this article as: Zhu, K-q., Wang, L., Chen, Q., He, M-y., Iron-catalyzed oxidative dehydrogenative coupling of ethers with aryl tetrazoles, *Tetrahedron Letters* (2015), doi: http://dx.doi.org/10.1016/j.tetlet. 2015.06.091

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

# **ACCEPTED MANUSCRIPT**

### **Graphical Abstract**





#### Tetrahedron Letters journal homepage: www.elsevier.com

### Iron-catalyzed oxidative dehydrogenative coupling of ethers with aryl tetrazoles

#### Kai-qiang Zhu, Liang Wang,\* Qun Chen and Ming-yang He\*

School of Petrochemical Engineering, Changzhou University, Changzhou 213164, P. R. China.

#### ARTICLE INFO

ABSTRACT

Article history: Received Received in revised form Accepted Available online An iron-catalyzed oxidative dehydrogenative coupling of ethers with aryl tetrazoles has been developed. A wide variety of tetrazoles and ethers survived the reaction conditions to deliver the corresponding hemiaminal derivatives in moderate to good yields.

2009 Elsevier Ltd. All rights reserved.

eywords: Dehydrogenative coupling Iron catalysis Tetrazoles Ethers

Tetreazoles as well as their derivatives are very important *N*-containing heterocycles that showed wide applications in organic chemistry,<sup>1</sup> medicinal chemistry,<sup>2</sup> and material science.<sup>3</sup> They were also employed as directing groups in C–H activation reactions.<sup>4</sup> Thus, development of efficient methods for functionalization of tetrazoles is of great importance.

The N-alkylation and arylation of tetrazoles are the most straightforward methods for the synthesis of tetrazole derivatives. In recent years, transition-metal-catalyzed N-arylation of tetrazoles has been well-established.<sup>5</sup> The N-alkylation of tetrazoles, however, has a limited scope. Generally, they were prepared via nucleophilic substitution reaction between a tetrazole and an alkyl halide.<sup>6</sup> The major drawbacks associated with this procedure were the utilization of alkyl halides, extra bases, and the formation of isomeric dialkylated mixtures. Just recently, our group report a metal-free alkylation of aryl tetrazoles with benzylic C-H substrates by using n-Bu<sub>4</sub>NI as catalyst and t-BuOOH as oxidant.<sup>7</sup> Later, we reported a direct amination of ethers with aryl tetrazoles and triazoles under the same reaction conditions.<sup>8</sup> However, high catalyst loading and a large excess of alkyl ethers were used, and moderate yields were obtained in most cases. We hypothesized that the low solubility of tetrazoles in alkyl ethers contributed to the low conversion. Therefore, an efficient method for this reaction is still highly desirable.

As a cheap, readily available and nontoxic metal, iron catalysts have received wide attention in organic synthesis.<sup>9</sup> Particularly, iron-catalyzed the cross-dehydrogenative coupling (CDC) reactions are highly attractive from both environmental and economic points of view;<sup>10</sup> which provides a facile approach to construct various *N*-containing heterocycles. In coujunction with our current interest in the functionalization of tetrazoles,<sup>4c,7,8</sup> we herein report an iron-catalyzed oxidative dehydrogenative coupling of ethers with aryl tetrazoles (Scheme 1).



Scheme 1. Iron-catalyzed amination of alkyl ethers with aryl tetrazoles.

Initially, reaction of 5-phenyl-2*H*-tetrazole **1a** and tetrahydrofuran (THF) **2a** was investigated to optimize the reaction conditions (Table 1). Interestingly, the desired product **3a** was obtained in 17% yield in the presence of 5 equivalents of *tert*-butyl hydroperoxide (TBHP) using THF as solvent (Table 1, entry 1). Addition of FeCl<sub>3</sub>.6H<sub>2</sub>O (5 mol%) greatly promoted the reaction, providing **3a** in 66% yield (Table 1, entry 2). The reaction did not occur in the absence of TBHP, indicating that both iron catalyst and TBHP were essential for this transformation. Considering the low solubilities of tetrazoles in other alkyl ethers, various solvents were examined (Table 1,

<sup>\*</sup>Corresponding author. Tel.: +86-519-86330263; fax: +86-519-86330251. E-mail address: lwcczu@126.com; hemingyangjpu@yahoo.com

## ACCEPTED MANUSCRIPT

#### Tetrahedron Letters

entries 4-8). Ethyl acetate and 1,2-dichloroethane turned out to be the best solvents, giving the desired product **3a** in 85% and 83% yield respectively. Moderate yields were obtained by using dichloromethane or acetonitrile as a solvent. Reaction in *N*,*N*dimethylformamide, however, afforded trace amount of **3a**. Other iron catalysts were also tested and relatively lower yields were obtained (Table 1, entries 9-13). Further studies showed that reducing the amount of FeCl<sub>3</sub>.6H<sub>2</sub>O, TBHP as well as THF led to lower yield (48, 65 and 67% yields, respectively). Oxidants were also crucial for this reaction, and other oxidants including H<sub>2</sub>O<sub>2</sub> and di-*tert*-butyl peroxide (DTBP) resulted in poor yields (Table 1, entries 18 and 19). Reaction temperature was finally evaluated. The results clearly showed that 90 °C was the optimum.

Table 1 Optimized the reaction conditions.<sup>a</sup>

(	N=N NH +		Fe salts	N=N N	5
	1a	2a		3a	
Entry	Catalyst (mol%)	2a (equiv)	Oxidant	Solvent	$\text{Yield}(\%)^{b}$
			(equiv)		
1	-	-	TBHP (5)	THF	17 <sup>c</sup>
2	$FeCl_{3.}6H_{2}O(5)$	-	TBHP (5)	THF	66 <sup>c</sup>
3	FeCl <sub>3</sub> .6H <sub>2</sub> O (5)	-	-	THF	NR <sup>c,d</sup>
4	$FeCl_{3.6}H_{2}O(5)$	10	TBHP (5)	EtOAc	85
5	$FeCl_{3.6}H_{2}O(5)$	10	TBHP (5)	DCM	70
6	$FeCl_{3.6}H_{2}O(5)$	10	TBHP (5)	DCE	83
7	$FeCl_{3.6}H_{2}O(5)$	10	TBHP (5)	CH <sub>3</sub> CN	63
8	$FeCl_{3.6}H_{2}O(5)$	10	TBHP (5)	DMF	trace
9	$FeCl_3(5)$	10	TBHP(5)	EtOAc	71
10	$FeBr_3(5)$	10	TBHP (5)	EtOAc	72
11	$Fe(acac)_3(5)$	10	TBHP (5)	EtOAc	52
12	$FeCl_2(5)$	10	TBHP (5)	EtOAc	28
13	$Fe(OAc)_2(5)$	10	TBHP (5)	EtOAc	21
14	FeCl <sub>3</sub> .6H <sub>2</sub> O (2.5)	10	TBHP (5)	EtOAc	48
15	FeCl <sub>3</sub> .6H <sub>2</sub> O (10)	10	TBHP (5)	EtOAc	74
16	$FeCl_{3.6}H_{2}O(5)$	10	TBHP (3)	EtOAc	65
17	FeCl <sub>3</sub> .6H <sub>2</sub> O (5)	5	TBHP(5)	EtOAc	67
18	$FeCl_{3.}6H_{2}O(5)$	10	$H_2O_2(5)$	EtOAc	$NR^{d}$
19	$FeCl_{3.}6H_{2}O(5)$	10	DTBP (5)	EtOAc	23
20	FeCl <sub>3</sub> .6H <sub>2</sub> O (5)	10	TBHP (5)	EtOAc	75 <sup>e</sup> , 82 <sup>f</sup>

<sup>*a*</sup> Reaction conditions: **1a** (0.3 mmol), **2a**, catalyst, oxidant, and solvent (1 mL) were heated at 90 °C under nitrogen for 12 h. <sup>*b*</sup> Isolated yields. <sup>*c*</sup> THF as substrate and solvent. <sup>*d*</sup> No reaction. <sup>*e*</sup> Reaction at 80 °C. <sup>*f*</sup> Reaction at 100 °C.

With the optimized yields in hand, a series of substituted tetrazoles were employed to explore the generality of this protocol using THF 2a as standard substrate (Table 2).<sup>11</sup> The results are summarized in Table 2. As shown from the table, most of aryl tetrazoles reacted with THF to give the desired products 3 in moderate to good yields (45~85%). Functional groups such as methyl, methoxyl, chloro, fluoro, and trifluoromethyl survived the reaction conditions. When 5-(3nitrophenyl)-2H-tetrazole was applied under the standard conditions, trace amount of product was detected. Para-hydroxyl substituted tetrazole, however, did not survived. Moreover, alkyl tetrazole such as 5-benzyl-2H-tetrazole was also checked (although it is not shown in Table 2); however, rather poor yield was obtained. It was also worth noting that coupling site took place mainly at the N2-position, with only trace amount of N1alkylated isomers.<sup>6</sup>

Various alkyl ethers were then evaluated and the results are summarized in Table 3. Tetrahydropyran also gave the desired product 3k in 59% yield. For substrate 1,4-dioxane and 1,3dioxolane, different results were obtained. The former reaction gave product 3l in 64% yield by using DCE as solvent; however, no amination product 3m was detected (3m was obtained as the product in TBAI/TBHP system<sup>8</sup>). Instead, this reaction provided two other compounds **3n** and **3o**.<sup>12</sup> Similarly, only compound **3n** (43%) was detected when diethyl ether was used as the substrate. Other straight chain ethers also showed good compatibilities and provided the corresponding products **3q** and **3r** in 51% and 80% yields, respectively. In addition, unsaturated alkyl ethers such as 3,4-dihydro-2*H*-pyran was employed to check the regioselectivity. Interestingly, neither allylated tetrazole nor hemiaminal **3s** was obtained. Instead, a double bond addition reaction occurred between tetrazole and 3,4-dihydro-2*H*-pyran, giving product **3k** in 82% yield.

Table 2 Reaction scope for aryl tetrazoles.<sup>a</sup>



<sup>*a*</sup> Reaction conditions: **1** (0.3 mmol), **2a** (3 mmol), FeCl<sub>3.6</sub>H<sub>2</sub>O (0.015 mmol, 5 mol%), TBHP (1.5 mmol, 5 equiv), and EtOAc (1 mL), 90 °C, N<sub>2</sub>, 12 h. <sup>*b*</sup> Isolated yields. <sup>o</sup> No reaction.

Table 3. Reaction scope for alkyl ethers.<sup>a</sup>



<sup>*a*</sup> Reaction conditions: **1a** (0.3 mmol), **2** (3 mmol), FeCl<sub>3.6</sub>H<sub>2</sub>O (0.015 mmol, 5 mol%), TBHP (1.5 mmol, 5 equiv), and EtOAc (1 mL), 90 °C, N<sub>2</sub>, 12 h, isolated yields. <sup>*b*</sup> DCE as solvent. <sup>*c*</sup> Product not detected.

Control experiments were performed to gain insight into the mechanism (Scheme 2). Initially, when 2,2,6,6-tetramethylpiperidine *N*-oxide (TEMPO) was added, the reaction was completely suppressed. Meanwhile, the TEMPO-adduct **4** was obtained (eq 1). This result indicated that a radical pathway

## ACCEPTED MANUSCRIPT

may be involved in this reaction. Moreover, kinetic isotopic effect (KIE) experiment was carried out under the standard reaction conditions (eq 2). The result shows a significant isotopic effect ( $k_H/k_D = 2.7$ ), indicating that the C-H bond cleavage is the rate-determinating step of this transformation.



Scheme 2. Control experiments.

Based on the above results and the similar study that previously reported, <sup>10h,13,14</sup> a possible mechanism was proposed (Scheme 3). Barton and co-workers confirmed that Fe(III) salt reacted with TBHP to generate *tert*-butylperoxy radical and Fe(II) in their study (step a).<sup>12</sup> TBHP can decompose into *tert*-butoxyl radical and hydroxyl anion in the presence of the ferrous catalyst (step b).<sup>10h</sup> Thus, we believe that one hydrogen atom of ether is abstracted by either *tert*-butylperoxy radical or *tert*-butoxyl radical to form radical **A**, followed by oxidation by Fe<sup>3+</sup> to generate oxonium ion **B** (step c). Meanwhile, deprotonation of tetrazole gives **C** (step d). Finally, nucleophilic addition of **C** to **B** provides the desired product (step e).





In summary, an iron-catalyzed oxidative dehydrogenative coupling of ethers with aryl tetrazoles has been developed. This protocol provides a simple and straightforward approach for the preparation of pharmaceutically and synthetically useful tetrazole derivatives. Substrates with a series of functional groups proceed smoothly to provide the corresponding products in moderate to good yields.

#### Acknowledgments

This project was financially supported by the National Natural Science Foundation of China (NO. 21302014), the Natural Science Foundation for Colleges and Universities of Jiangsu Province (13KJB150002), and ordinary university graduate student research innovation projects of Jiangsu province (No. KYZZ\_0303).

#### References and notes

- Larock, R. C. Comprehensive Organic Transformations: A Guide to Functional Group Preparations; VCH Publishers: New York, 1989.
- 2 (a) Fernandez-Bachiller, M. I.; Perez, C.; Monjas, L.; Rademann, J.; Rodríguez-Franco, M. I. *J. Med. Chem.* **2012**, *55*, 1303; (b) Das, S. G.; Srinivasan, B.; Hermanson, D. L.; Bleeker, N. P.; Doshi, J. M.; Tang, R.; Beck, W. T.; Xing, C. *J. Med. Chem.* **2011**, *54*, 5937.
- 3 (a) Aromí, G.; Barrios, L. A.; Roubeau, O.; Gamez, P. Coord. Chem. Rev. 2011, 255, 485; (b) Singh, R. P.; Verma, R. D.; Meshri, D. T.; Shreeve, J. M. Angew. Chem., Int. Ed. 2006, 45, 3584; (c) Tao, G. H.; Guo, Y.; Joo, Y. H.; Twamley, B.; Shreeve, J. M. J. Mater. Chem. 2008, 18, 5524.
- 4 (a) Seki, M. ACS Catal. 2011, *1*, 607; (b) Diers, E.; Kumar, N. Y. P.; Mejuch, T.; Marek, I.; Ackermann, L. *Tetrahedron* 2013, 69, 4445; (c) Wang, L.; Wu, W. T.; Chen, Q.; He, M. Y. Org. Biomol. Chem. 2014, *12*, 7923.
- 5 (a) Lam, P. Y. S.; Clark, C. G.; Saubern, S.; Adams, J.; Winters, M. P.; Chan, D. M. T.; Combs, A. *Tetrahedron Lett.* **1998**, *39*, 2941; (b) Davydov, D. V.; Beletskaya, I. P.; Semenov, B. B.; Smushkevich, Y. I. *Tetrahedron Lett.* **2002**, *43*, 6217; (c) Fedorov, A. Y.; Finet, J.-P. *Tetrahedron Lett.* **1999**, *40*, 2747; (d) Beletskaya, I. P.; Davydov, D. V.; Gorovoy, M. S. *Tetrahedron Lett.* **2002**, *43*, 6221; (e) J. Li, Y.; Gao, L.-X.; Han, F.-S. *Chem. Commun.* **2012**, *48*, 2719; (f) Liu, C.-Y.; Li, Y.; Ding, J.-Y.; Dong, D.-W.; Han, F.-S. *Chem. Eur. J.* **2014**, *20*, 2373; (g) Onaka, T.; Umemoto, H.; Miki, Y.; Nakamura, A.; Maegawa, T. J. Org. *Chem.* **2014**, *79*, 6703.
- 6 Aridoss, G.; Laali, K. K. Eur. J. Org. Chem. 2011, 6343.
- 7 Wang, L.; Zhu, K. Q.; Chen, Q.; He, M. Y. J. Org. Chem. 2014, 79, 11780.
- 8 Wang, L.; Zhu, K. Q.; Wu, W. T.; Chen, Q.; He, M. Y. Catal. Sci. Technol. 2015, 5, 2891.
- 9 For representative reviews on iron chemistry, see: (a) Fürstner, A. Angew.Chem., Int. Ed. 2009, 48, 1364; (b) Czaplik, W. M.; Mayer, M.; Cvengros, J.; Jacobi von Wangelin, A. ChemSusChem 2009, 2, 396; (c) Sherry, B. D.; Fü'rstner, A. Acc. Chem. Res. 2008, 41, 1500; (d) Correa, A.; Mancheňo, O. G.; Bolm, C. Chem. Soc. Rev. 2008, 37, 1108; (e) Enthaler, S.; Junge, K.; Beller, M. Angew. Chem., Int. Ed. 2008, 47, 3317; (f) Bolm, C.; Legros, J.; Le Paih, J.; Zani, L. Chem. Rev. 2004, 104, 6217.
- (a) Jia, F.; Li, Z. Org. Chem. Front. 2014, 1, 194; (b) Gaster, E.; Vainer, Y.; Regev, A.; Narute, S.; Sudheendran, K.; Werbeloff, A.; Shalit, H.; Pappo, D. Angew. Chem., Int. Ed. 2015, 54, 4198; (c) Guo, S.; Li, Y.; Wang, Y.; Guo, X.; Meng, X.; Chen, B. Adv. Synth. Catal. 2015, 357, 950; (d) Zhao, J.; Fang, H.; Zhou, W.; Han, J.; Pan, Y. J. Org. Chem. 2014, 79, 3847; (e) Cheng, Y.; Dong, W.; Wang, L.; Parthasarathy, K.; Bolm, C. Org. Lett. 2014, 16, 2000; (f) Parnes, R.; Kshirsagar, U.; Werbeloff, A.; Regev, C.; Pappo, D. Org. Lett. 2012, 14, 3324; (g) Liu, H.; Cao, L.; Sun, J.; Fossey, J.; Deng, W. Chem. Commun. 2012, 48, 2674; (h) Pan, S.; Liu, J.; Li, H.; Wang, Z.; Guo, X.; Li, Z. Org. Lett. 2010, 12,1932; (i) Li, Y.; Li, B.; Lu, X.; Lin, S.; Shi, Z. Angew. Chem., Int. Ed. 2009, 48, 3817.
- 11 Typical procedure for the synthesis of 3: To a mixture of tetrazole 1 (0.3 mmol), alkyl ethers 2 (3 mmol), and FeCl<sub>3</sub>.6H<sub>2</sub>O (4.1 mg, 0.015 mmol, 5 mol%), ethyl acetate (1 mL) was added under nitrogen. Then *t*-BuOOH (70% aqueous, 1.5 mmol, 5 equiv) was dropped into the mixture under nitrogen. The reaction mixture was stirred at 90 °C for 12 h. After reaction, the mixture was allowed to cool to room temperature. The solvent was then removed under vacuum, and the residue was purified by silica gel chromatography using a mixture of PE/EA to afford the desired product 3.
- 12 The plausible mechanism for generation of compounds **3n** and **3o** is unclear at current stage. Further studies are undergoing in our laboratory.
- 13 Barton, D. H. R.; Le Gloahec, V. N. Tetrahedron 1998, 54, 15457.
- 14 Porcheddu, A.; De Luca, L. Adv. Synth. Catal. 2012, 354, 2949.