

with the p44 reductase gave highly optically pure (*R*)-2-alkyl-cyclohexanones, **6b–8b**.¹⁸ These results indicate that the stereospecificity of the hydrogenation of the C-C double bond was opposite between the p90 and p44 reductases.

Thus, it was demonstrated that the enone reductases from *N. tabacum* were able to reduce enantiotropically the C-C double bond of enones to afford optically active 2-alkylated ketones. It is worth noting that each enantiomer of 2-alkylated ketones can be synthesized by selective use of these enone reductases from *N. tabacum*.

The authors thank the Instrument Center for Chemical Analysis of Hiroshima University for the measurements of ¹H NMR, GC-MS and CD spectra.

References and Notes

- 1 T. Hirata, H. Hamada, T. Aoki, and T. Suga, *Phytochemistry*, **21**, 2209 (1982).
- 2 T. Suga, H. Hamada, and T. Hirata, *Chem. Lett.*, **1987**, 471.
- 3 T. Suga, H. Hirata, H. Hamada, and S. Murakami, *Phytochemistry*, **27**, 1041 (1988).
- 4 T. Hirata, Y. X. Tang, K. Okano, and T. Suga, *Phytochemistry*, **28**, 3331 (1989).
- 5 T. Hirata, S. Izumi, K. Shimoda, and M. Hayashi, *J. Chem. Soc., Chem. Commun.*, **1993**, 1426.
- 6 K. Shimoda, D. I. Ito, S. Izumi, and T. Hirata, *J. Chem. Soc., Perkin Trans. 1*, **1996**, 355.
- 7 The suspension cells of *N. tabacum*¹ were cultivated at 25 °C for 3 weeks in Murashige and Skoog's medium⁸ on a rotary shaker (75 rpm). The cells (200 g) were homogenized in 0.1 M Na-Pi buffer (pH 6.8) containing 10 mM 2-mercaptoethanol and 5 mM dithiothreitol. After centrifugation at 10000 *g* for 15 min, the supernatant was fractionated by treatment with (NH₄)₂SO₄ (40 to 60% satn.). The crude enzyme soln obtained was desalted and then applied to a DEAE-Toyopearl column with a 0–0.5 M linear gradient of NaCl in 50 mM Tris-HCl buffer (pH 8.0) containing 1.0 mM 2-mercaptoethanol and 1.0 mM dithiothreitol (buffer A) to give three enone reductase fractions, which contained the 44, 74 and 90 kDa proteins, respectively. Each of the fractions was further purified on a hydroxylapatite column with buffer A and a Red-Toyopearl column with buffer A containing a 0–1.0 M linear gradient of NaCl.
- 8 T. Murashige and F. Skoog, *Physiol. Plant.*, **15**, 473 (1962).
- 9 The mixture (2 mL) of enone (100 μmol), NADPH (200 μmol), Triton X-100 (0.1%), and enzyme preparation (5 μg) in 50 mM Tris-HCl buffer (pH 8.0) containing 1.0 mM 2-mercaptoethanol and 1.0 mM dithiothreitol was incubated at 35 °C for 8 h. After incubation, the reaction mixture was subjected to GLC and GC-MS analyses, and then the products were purified by column chromatography on silica gel. The absolute configurations and enantiomeric purities of the products were determined by circular dichroism (CD) spectra and the peak area of the corresponding enantiomers in the GLC analyses on CP cyclodextrin β 236M-19 column.
- 10 2-Alkylidenecyclohexanones (**1** and **2**) were prepared from the α-alkylidene acetals of cyclohexanone according to the reported procedure.^{11,12} **1**: MS (EI) *m/z* 110 (*M*⁺); ¹H NMR (500 MHz; CDCl₃) δ 6.01 (1H, t, *J* = 1.9 Hz, >C=CH₂), 6.12 (1H, t, *J* = 1.9 Hz, >C=CH₂); **2**: MS (EI) *m/z* 138 (*M*⁺), ¹H NMR (CDCl₃) δ 1.05 (3H, t, *J* = 7.6 Hz, Me), 2.11 (2H, q, *J* = 7.6 Hz, –CH₂–Me), 6.61 (1H, tt, *J* = 7.4 and 2.1 Hz, >C=CH–). 2-Alkyl-2-cyclohexen-1-ones (**3–5**) were prepared from *O*-methoxybenzoic acid by reductive alkylation according to the reported procedure.¹³ **3**: MS (EI) *m/z* 110 (*M*⁺), ¹H NMR (CDCl₃) δ 1.77 (3H, q, *J* = 1.5 Hz, Me), 6.74 (1H, tq, *J* = 4.3 and 1.5 Hz, >C=CH–); **4**: MS (EI) *m/z* 124 (*M*⁺), ¹H NMR (CDCl₃) δ 1.00 (3H, t, *J* = 7.5 Hz, Me), 2.20 (2H, qq, *J* = 7.5 and 1.5 Hz, –CH₂–Me), 6.69 (1H, tt, *J* = 4.1 and 1.5 Hz, >C=CH–); **5**: MS (EI) *m/z* 138 (*M*⁺), ¹H NMR (CDCl₃) δ 0.89 (3H, t, *J* = 7.4 Hz, Me), 1.41 (2H, se, *J* = 7.5 Hz, –CH₂–Me), 2.15 (2H, tq, *J* = 7.4 and 1.2 Hz, –CH₂–CH₂–Me), 6.69 (1H, t, *J* = 4.3 Hz, >C=CH–).
- 11 F. Huet, M. Pellet, and J. M. Conia, *Tetrahedron Lett.*, **1977**, 3505.
- 12 K. Matsumoto, Y. Kawabata, J. Takahashi, Y. Fujita, and M. Hatanaka, *Chem. Lett.*, **1998**, 283.
- 13 D. F. Taber, *J. Org. Chem.*, **41**, 2649 (1976).
- 14 **6a**: [θ]₂₈₈ +960 (*c* 0.25, MeOH) {lit.¹⁵ [θ]₂₈₈ –987 for *R* enantiomer}; **8a**: [θ]₂₈₈ +1860 (*c* 0.25, MeOH) {lit.¹⁶ [θ]₂₈₈ +2480}.
- 15 C. J. Cheer and C. Djerassi, *Tetrahedron Lett.*, **1976**, 3877.
- 16 A. I. Meyers, D. R. Williams, G. W. Erickson, S. White, and M. Druehlinger, *J. Am. Chem. Soc.*, **193**, 3081 (1981).
- 17 **6a**: [θ]₂₈₈ +993 (*c* 0.3, MeOH); **7a**: [θ]₂₈₈ +2293 (*c* 0.1, MeOH); **8a**: [θ]₂₈₈ +2351 (*c* 0.12, MeOH).
- 18 **6b**: [θ]₂₈₈ –989 (*c* 0.2, MeOH); **7b**: [θ]₂₈₈ –2341 (*c* 0.1, MeOH) {lit.¹⁵ [θ]₂₈₈ +2200 for *S* enantiomer}; **8b**: [θ]₂₈₈ –2483 (*c* 0.1, MeOH).