Tetrahedron 65 (2009) 3139-3148

Contents lists available at ScienceDirect

# Tetrahedron

journal homepage: www.elsevier.com/locate/tet

# Catalytic enantioselective conjugate additions with $\alpha$ , $\beta$ -unsaturated sulfones

# Hongming Li, Jun Song, Li Deng\*

Department of Chemistry, Brandeis University, Waltham, MA 02454-9110, USA

## ARTICLE INFO

Article history: Received 13 October 2008 Received in revised form 16 November 2008 Accepted 19 November 2008 Available online 25 November 2008

Keywords: Organocatalyst Hydrogen bonding Cinchona alkaloids Chiral sulfones Conjugate addition

# 1. Introduction

Sulfones are versatile intermediates in organic synthesis.<sup>1</sup> Enantioselective C–C bond forming reactions generating chiral sulfones are of significant interest to asymmetric synthesis.<sup>2</sup> In particular asymmetric conjugate additions with trisubstituted carbon nucleophiles to  $\alpha,\beta$ -unsaturated sulfones **4** (Scheme 1) provide a straightforward strategy to form optically active building blocks bearing the sulfone functionality and a quaternary stereocenter.<sup>3,4</sup> In this full article, we document in detail our efforts on the development of such reactions catalyzed by practically accessible, cinchona alkaloid-based organic catalysts.<sup>3</sup>

In 2004 we demonstrated that 6'-OH cinchona alkaloid derivatives such as **5**, bearing a hydrogen bond acceptor in quinuclidine and a hydrogen bond donor in 6'-OH, could catalyze the conjugate additions of a broad range of enolizable carbon Michael donors **2** to nitroalkenes **1** with remarkably high enantioselectivity and diastereoselectivity (Scheme 1).<sup>5,6</sup> These studies provided the first examples of cinchona alkaloids with rotational freedom around C8–C9 and C4'–C9 bonds serving as highly efficient acidbase bifunctional catalysts for asymmetric reactions. Mechanistic and conformational studies unambiguously established that **5** adopted a *gauche*-open active conformer (Scheme 1) in the transition state.<sup>5b</sup> With this key information in hand, we are able to construct a transition state model (**6**, Scheme 1) that is consistent with the stereochemistry outcome of the **5**-catalyzed conjugate

# ABSTRACT

We describe the development of a highly efficient catalytic asymmetric conjugate addition to  $\alpha$ , $\beta$ -unsaturated sulfones. Utilizing practical bifunctional organic catalysts and involving air- and moisturetolerant conditions, conjugate additions of a wide range of Michael donors to  $\alpha$ , $\beta$ -unsaturated sulfones proceeded in excellent enantioselectivity/diastereoselectivity and high yield. This efficient and operationally simple new catalytic asymmetric reaction should provide a versatile approach for the asymmetric synthesis of chiral sulfones bearing all-carbon quaternary stereocenters.

© 2008 Elsevier Ltd. All rights reserved.

additions to nitroalkenes **1**. In this model the C6′–OH cinchona alkaloids **5**, adopting the *gauche*-open conformation, simultaneously activated and oriented the Michael donors **3** (generated by tautomerization of ketones **2**) and the nitroalkenes **1** via a network of hydrogen bonding interactions.<sup>5b</sup> We reasoned that if the replacement of the nitro with the sulfone functionality in the Michael donor did not disrupt or alter the pattern of hydrogen bonding network between the catalyst and the reacting substrates, 6′-OH cinchona alkaloid **5** might be able to catalyze conjugate additions of enolizable Michael donors **2** to  $\alpha$ , $\beta$ -unsaturated sulfones with high diastereoselectivity and enantioselectivity. Thus, we initiated an investigation of these conjugate additions to  $\alpha$ , $\beta$ -unsaturated sulfones with cinchona alkaloids **5**.

# 2. Results and discussion

# 2.1. Catalyst discovery and reaction optimizations

Our experiments began with a screening of a variety of cinchona alkaloid derivatives for their ability to promote enantioselective addition of  $\alpha$ -phenyl  $\alpha$ -cyanoacetate **10A** to the commercially available phenyl vinyl sulfone (**4a**). As summarized in Table 1, the conjugate addition proceeded smoothly in toluene at room temperature in the presence of various natural and modified cinchona alkaloids. The enantioselectivity of the reaction was, however, found to be critically dependent on the structure of the cinchona alkaloid derivatives. Reactions with natural cinchona alkaloids such as quinidine and quinine generated the 1,4-adduct **11A** as a nearly racemic mixture (entries 1 and 2). The enantioselectivity was in general improved with cinchona alkaloids bearing a C9-substituent





<sup>\*</sup> Corresponding author. Tel.: +1 781 736 2529; fax +1 781 736 2516. *E-mail address:* deng@brandeis.edu (L. Deng).



Scheme 1. Rationale for the development of a catalytic asymmetric conjugate addition to α,β-unsaturated sulfones (4) with 6'-OH cinchona alkaloids 5.

(entries 3–7, 9–12). Importantly, in line with our hypothesis, the 6'-OH cinchona alkaloids **5** were found to be much more effective than those bearing C6'–OMe. The pronounced effect of the C9-substituent on the enantioselectivity of the C6'–OH cinchona alkaloids

**5** is also noteworthy. While modest enantioselectivity was afforded by either the C6'–OH cinchona alkaloid bearing a C9–OH (entry 8) or a rigid C6'–OH cinchona alkaloid derivative (entry 7), significantly higher enantioselectivity could be attained with C6'–OH

# Table 1

Cinchona alkaloid-catalyzed addition of  $\alpha$ -phenyl  $\alpha$ -cyanoacetate **10A** to vinyl phenyl sulfone (**4a**)<sup>a</sup>



Ph	SO <sub>2</sub> Ph	toluene (0.5M)	Ph * SO <sub>2</sub> Ph	
10A	4a	11, 1711	11A	

Entry	Catalyst	Conversion <sup>b</sup> /%	ee <sup>c</sup> /%
1	Q	>98	0
2	QD	>98	3
3	DHQD-PHN	79	24
4	(DHQD) <sub>2</sub> PYR	>98	33
5	(DHQR) <sub>2</sub> PHAL	80	9
6	(DHQR) <sub>2</sub> AQN	>98	4
7	β-ICD	98	42
8	QD-5a	90	45
9	QD-5b	>98	63
10	QD- <b>5c</b>	91	74
11	Q-5b	65	74
12	Q- <b>5c</b>	>98	84

<sup>a</sup> Reactions were run with **10A** (0.3 mmol), **4a** (0.1 mmol), and the catalyst (0.02 mmol) in toluene (0.2 mL) at rt for 17 h.

<sup>b</sup> Determined by <sup>1</sup>H NMR analysis.

<sup>c</sup> Determined by HPLC analysis.

#### Table 2

Solvent effect on the addition of α-phenyl α-cyanoacetate 10A to vinyl phenyl sulfone (4a) with Q- $5c^{\rm a}$ 

Entry	Time/h	Solvent	Conversion <sup>b</sup> /%	ee <sup>c</sup> /%
1	17	TBME	9	58
2	17	CHCl <sub>3</sub>	17	79
3	17	$CH_2Cl_2$	17	74
4	17	DCE	22	81
5	17	Toluene	46	95
6	72	Toluene	95	95

<sup>a</sup> Reactions were run with **10A** (0.3 mmol), **4a** (0.1 mmol), and Q-**5c** (0.02 mmol) in toluene (0.2 mL) at -25 °C.

<sup>b</sup> Determined by <sup>1</sup>H NMR analysis.

<sup>c</sup> Determined by HPLC analysis.

cinchona alkaloids bearing either an aryl or alkyl ether at C9 (entries 9 and 10). Thus, the tunable group at C9 provided a crucial handle for the optimization of the catalytic enantioselectivity of **5**. Our screening studies identified **5c** as the best catalyst (entries 10 and 12).

Next, we examined the reaction in various solvents. The reaction was found to proceed at the fastest rate and in the highest enantioselectivity in toluene. It is noteworthy that such a non-polar solvent is known to favor the formation and stabilization of hydrogen bonding complexes between solvates. An excellent enantioselectivity could be achieved with Q-**5c** in toluene at -25 °C (entries 5 and 6, Table 2).

# 2.2. Conjugate additions of $\alpha$ -substituted $\alpha$ -cyanoacetate to vinyl sulfones

After establishing the optimal reaction conditions we began to investigate the substrate scope of the **5c**-catalyzed conjugate addition to vinyl sulfone (**4a**). The  $\alpha$ -cyanoacetates **10** bearing a range of  $\alpha$ -aryl and  $\alpha$ -heteroaryl groups of varying electronic and steric properties underwent efficient enantioselective additions to phenyl vinyl sulfone (**4a**), providing the corresponding 1,4-adducts **11** bearing the all-carbon quaternary stereocenter in excellent enantioselectivity and good to excellent yield (Table 3).

Having established a general scope with respect to  $\alpha$ -aryl  $\alpha$ -cyanoacetates, we attempted to extend the scope to  $\alpha$ -alkyl  $\alpha$ -cyanoacetates. We found that, compared to their aryl congeners,  $\alpha$ -alkyl  $\alpha$ -cyanoacetates were significantly less active as Michael donor for the cinchona alkaloid-catalyzed conjugate addition to

#### Table 3

Additions of  $\alpha$ -aryl  $\alpha$ -cyanoacetates **10** to vinyl phenyl sulfone (**4a**) with **5c**<sup>a,b,c</sup>

EtO <sub>2</sub> C <b>10</b>	CN R	+ SO <sub>2</sub> Ph 4a		Q-5c (QD-5	Et	0 <sub>2</sub> C CN R * 11	`SO₂Ph
Entry		R	T/°C	Time/h	Product	Yield <sup>d</sup> /%	ee <sup>e</sup> /%
1	10A	Ph-	-25	72 (72)	11A	89 (80)	95 (91)
2	10B	4-Me-Ph-	0	48	11B	96	93
3	10C	4-MeO-Ph-	0	70	11C	92	93
4	10D	4-F-Ph-	-25	72	11D	90	94
5	10E	4-Cl-Ph-	-25	69 (72)	11E	95 (94)	94 (89)
6	10F	4-Br-Ph-	-25	69 (72)	11F	95 (95)	94 (88)
7	10G	3-Cl-Ph-	-25	60	11G	96	93
8	10H	2-Naphthyl–	-25	60 (60)	11H	96 (95)	97 (90)
9	10I	2-Thienyl	-25	48 (48)	11I	95 (91)	93 (88)

<sup>a</sup> Unless noted, reactions were run with **10** (0.5–0.6 mmol), **4a** (0.2 mmol), and **5c** (0.04 mmol).

<sup>b</sup> The catalyst was recovered in greater than 95% yield after the reaction.

<sup>c</sup> The results in parentheses were obtained with QD-**5c** to give the opposite enantiomer.

<sup>d</sup> Isolated yield.

<sup>e</sup> Determined by HPLC analysis.

#### Table 4

Enantioselective conjugate addition of  $\alpha\text{-alkyl}$   $\alpha\text{-cyanoacetates}$  10 to vinyl sulfone with  $5c^{\text{a.b.c}}$ 



Entry	10	4	T/°C	Time/h	Product	Conversion/%	Yield <sup>d</sup> /%	ee <sup>e</sup> /%
1	10J	4a	23	17	11J	17	N.D.	87
2	10J	4b	23	17	12J	88	N.D.	86
3	10J	4b	0	96	12J	100	76	94 <sup>f</sup>
4	10K	4b	0	96 (96)	12K	100 (100)	85 (83)	92 (88)

 $^{\rm a}$  Unless noted, reactions were run with  $10\,(0.5\text{--}0.6\text{ mmol})$  and  $4\,(0.2\text{ mmol})$  with  $5c\,(0.04\text{ mmol}).$ 

<sup>b</sup> The catalyst was recovered in greater than 95% yield after the reaction.

 $^{\rm c}$  The results in parentheses were obtained with QD-5c to give the opposite enantiomer.

<sup>d</sup> Isolated yield

<sup>e</sup> Determined by HPLC analysis.

<sup>f</sup> The absolute configuration of adduct (+)-**12J** was determined to be 'S' by X-ray structure analysis, see Section 4.5 for details.

vinyl sulfones. While addition of  $\alpha$ -phenyl  $\alpha$ -cyanoacetate **10A** to phenyl vinyl sulfone **4a** with Q-**5c** proceeded to completion at room temperature after 17 h (entry 12, Table 1), the addition of  $\alpha$ -allyl- $\alpha$ -cyanoacetate **10J** to **4a** proceeded to only 17% conversion during the same period (entry 1, Table 4). Importantly, both reactions proceeded in comparable enantioselectivity (87% ee vs 84% ee).

We reasoned that the cinchona alkaloid-catalyzed conjugate addition with  $\alpha$ -alkyl  $\alpha$ -cyanoacetate might be accelerated considerably by enhancing the electrophilicity of the vinyl sulfone, which could be implemented via the introduction of electron-

Table 5

Enantioselective conjugate addition of  $\alpha$ -cyanoketones **13** and  $\beta$ -ketoesters **14** to vinyl sulfone **4a** catalyzed by **5c**<sup>a,b</sup>



Entry	Michael Donor	Time/h	Product	Yield <sup>c</sup> /%	ee <sup>d</sup> /%
1	13A	24	15A	96 (90)	95 (97)
2	13B	24	15B	96 (98)	95 (95)
3	13C	24	15C	91 (91)	96 <sup>e</sup> (97)
4	13D	24	15D	0	N.D.
5	14A	48	16A	93 (96)	96 (96)
6	14B	48	16B	96 (100)	96 (97)
7	14C	24	16C	0	N.D.

<sup>a</sup> Unless noted, reactions were run with **13** or **14** (0.40 mmol), **4a** (0.20 mmol) in toluene (0.40 mL) with Q-**5c** (20 mol %).

<sup>o</sup> The results in parentheses were obtained with QD-5c.

<sup>c</sup> Isolated yield.

<sup>d</sup> Determined by HPLC analysis.

<sup>e</sup> The absolute configuration of (–)-**15C** was determined to be 'S' by X-ray structure analysis, see Section 4.5 for details.

#### Table 6

Additions of  $\beta$ -ketoester **14D** to  $\beta$ -substituted vinyl triflone **17** with Q-**5c**<sup>a</sup>



Entry	Acceptor	Solvent	Conversion	dr <sup>b</sup>	ee of major isomer <sup>c</sup> /%	ee of minor isomer <sup>c</sup> /%
1	17a	Toluene	0	ND	ND	ND
2	17b	Toluene	100	5.5:1	96	56
3	17b	THF	100	3.7:1	81	6
4	17b	Et <sub>2</sub> O	100	4.3:1	92	12
5	17b	EtOAc	100	4.6:1	87	6
6	17b	ClCH <sub>2</sub> Cl <sub>2</sub> Cl	100	7.0:1	94	23
7	17b	CHCl <sub>3</sub>	100	5.7:1	96	26
8	17b	CH <sub>2</sub> Cl <sub>2</sub>	100	8.0:1	95	20

<sup>a</sup> Unless noted, reactions were run with 14D (0.025 mmol), 17 (0.0275 mmol) in a solvent (0.10 mL) with QD-5c (10 mol%).

<sup>b</sup> Determined by <sup>1</sup>H NMR analysis.

<sup>c</sup> Determined by HPLC analysis.

withdrawing substituents on the aromatic ring of the phenyl vinyl sulfone **4**. Based on our proposed transition state model outlined in Scheme 1, steric variations of the phenyl group of **4** should not disrupt how catalyst **5c** interacts with the reacting substrates. With this consideration in mind, we investigated the conjugate addition of **10J** to 3,5-bis(triflouromethyl)phenyl vinyl sulfone (**4b** $)^7$  with Q-**5c** at room temperature. We were pleased to find that the reaction was indeed significantly accelerated, proceeding to 88% conversion after 17 h. Importantly, the corresponding 1,4-adduct **12** was obtained in 86% ee (entry 2, Table 4). Thus, the ability of 6'-OH cinchona alkaloid catalysts to tolerate the structural change of the vinyl sulfone **4** allowed us to significantly improve the rate of the

conjugate additions of an  $\alpha$ -alkyl  $\alpha$ -cyanoacetate by tuning the electronic property of the vinyl sulfone acceptor. Importantly, the high enantioselectivity could be readily extended to the conjugate addition of  $\alpha$ -methyl  $\alpha$ -cyanoacetate **10K** to **4b** (entry 4, Table 4).

# 2.3. Conjugate additions of $\alpha$ -substituted $\alpha$ -cyanoketones and $\beta$ -ketoesters to vinyl sulfones

We also investigated the **5c**-catalyzed additions of cyclic  $\alpha$ -cyanoketones (**13A**–**C**) and cyclic  $\beta$ -ketoesters (**14A** and **B**) to vinyl sulfone **4a**. As illustrated in Table 5, reactions with these Michael donors proceeded in high yield and excellent enantioselectivity.

#### Table 7

Diastereoselective and enantioselective conjugate additions to  $\beta$ -alkyl vinyl triflones **17** catalyzed by **5** $c^{a}$ 



Entry	Donor	17	T/°C	Time/h	Product	dr	ee <sup>b</sup> /%	yield <sup>c</sup> /%
1	10A	17b	-50	96	18Ab	5.6:1 (4.6:1)	93 (87)	87 <sup>d</sup> (76) <sup>d</sup>
2	10D	17b	-50	96	18Db	26:1 (12:1)	92 (84)	94 <sup>d</sup> (91) <sup>d</sup>
3	10F	17b	-50	96	18Fb	49:1 (26:1)	81 (86)	92 <sup>d</sup> (92) <sup>d</sup>
4	13E	17b	-50	24	19Eb	>49:1 (>49:1)	99 (94)	95 <sup>d</sup> (94)
5	14D	17b	-20	96	20Db	32:1 (9:1)	98 (97)	$90^{d}(93)^{d}$
6	14D	17c	-20	72	20Dc	24:1 (8:1)	98 (95)	100 (95)
7	14E	17d	-20	48	20Ed	>49:1 (>49:1)	95 (89)	91 <sup>d</sup> (80) <sup>d</sup>

<sup>a</sup> Unless noted, reactions were run with **10** or **13** or **14** (0.10 or 0.20 mmol), **17** (0.10 mmol), and Q-**5c** (0.02 mmol) in methylene chloride (0.20 mL), and the results in parentheses were obtained with QD-**5c** to give the opposite enantiomer.

<sup>b</sup> Determined by HPLC analysis.

<sup>c</sup> Isolated yield.

<sup>d</sup> Yield of pure major diastereomer.

However, with either acyclic  $\alpha$ -cyanoketone (**13D**) or acyclic  $\beta$ -ketoester (**14C**), no conjugate addition product was obtained. Only starting materials were recovered.

# 2.4. Conjugate additions to $\beta\text{-substituted }\alpha,\beta\text{-unsaturated sulfones}$

Adjacent quaternary and tertiary stereocenters are common structural motifs in natural products. Highly diastereoselective and enantioselective conjugate additions of prochiral trisubstituted carbon nucleophiles to prochiral  $\beta$ -substituted Michael acceptors with a chiral catalyst could provide a one-step and stereoselective construction of such complex motifs from simple precursors. Thus, we began to investigate the conjugate additions of various nucleophiles to  $\beta$ -substituted  $\alpha$ , $\beta$ -unsaturated sulfones with Q-**5c**.

We first examined the Q-5c-catalyzed conjugate additions of  $\beta$ -ketoester **14D** to  $\beta$ -methyl vinyl sulfone **17a** and  $\beta$ -phenyl vinyl sulfone 17a' in toluene at room temperature. Unfortunately, no addition product was found, and only starting materials were recovered. Trifluoromethanesulfonyl group (CF<sub>3</sub>SO<sub>2</sub>-) is one of the strongest neutral electron-withdrawing groups known. If SO<sub>2</sub>Ph in 17 was replaced by SO<sub>2</sub>CF<sub>3</sub>, the resulting vinyl sulfones should be much more active toward conjugate additions. Following this thought, we began to investigate the QD-5c-catalyzed conjugate addition of  $\beta$ -ketoester **14D** to  $\beta$ -substituted vinyl triflone **17b**.<sup>8</sup> We were pleased to find that, in contrast to reactions with **17a**, the reaction proceeded smoothly to completion after 12 h at room temperature in toluene to afford the desired 1.4-adduct in excellent ee and promising dr (entry 2, Table 6). Further optimization revealed that CH<sub>2</sub>Cl<sub>2</sub>, ClCH<sub>2</sub>CH<sub>2</sub>Cl, and CHCl<sub>3</sub> were better solvents, and the conjugate addition proceeded in the highest enantioselectivity and diastereoselectivity in CH<sub>2</sub>Cl<sub>2</sub> (entry 8, Table 6).

Encouraged by these results, a variety of trisubstituted carbon Michael donors were subjected to the **5c**-catalyzed conjugate additions to various  $\beta$ -substituted vinyl triflones **17b–d**.<sup>8</sup> Excellent diastereoselectivity and enantioselectivity were obtained with various  $\alpha$ -aryl  $\alpha$ -cyanoacetates **10**,  $\alpha$ -cyanoketones **13**, and cyclic  $\beta$ -ketoesters **14** (Table 7).

# 3. Conclusions

In summary, we have developed a highly efficient catalytic asymmetric conjugate addition to  $\alpha,\beta$ -unsaturated sulfones. This reaction utilizes readily accessible and recyclable chiral organic catalysts and involves an air- and moisture-tolerant simple experimental protocol that avoids low temperature and low volumetric productivity. The general scopes with respect to both the donors and  $\alpha,\beta$ -unsaturated sulfones attained by the bifunctional cinchona alkaloid catalysts are particularly noteworthy. A wide range of  $\alpha$ -substituted  $\beta$ -carbonyl Michael donors, such as  $\alpha$ -cyanoacetate.  $\alpha$ -cyanoketone, and  $\beta$ -ketoesters, could be employed in the reaction. The reaction also tolerates variations in the  $\alpha$ , $\beta$ -unsaturated sulfones. Specifically, vinyl sulfones bearing either no or a β-substituent could be applied in this reaction. Consequently, the reaction provides a highly versatile catalytic entry for the asymmetric synthesis of chiral sulfones containing all-carbon quaternary stereocenters.

## 4. Experimental

# 4.1. General

<sup>1</sup>H and <sup>13</sup>C NMR spectra were recorded on a Varian instrument (400 MHz and 100 MHz, respectively) and internally referenced to SiMe<sub>4</sub> signal. Low resolution mass spectra for all the new compounds were recorded on a Hewlett-Packard 5989A GC/MS, and

exact mass spectra on a VG 7070 high resolution mass spectrometer. Infrared spectra were recorded on a Perkin Elmer FT-IR Spectrometer. Specific rotations were measured on a Jasco Digital Polarimeter. Liquid chromatography was performed using forced flow (flash chromatography) of the indicated solvent system on EM Science silica gel 60 (SiO<sub>2</sub>, 230-400 mesh). Analytical gas-liquid chromatography (GLC) was performed on a Hewlett-Packard 6890 Series instrument equipped with a split mode capillary injection system, a flame ionization detector, using a HP-5 GC column or a GC column with chiral stationary phase [Gamma cyclodextrin Trifluoroacetyl (30 m×0.25 mm) or HP Chiral (20% Permethylated B-Cyclodextrin, 30 m×0.25 mm)]. All the GC analyses of chiral cyanohydrin carbonates were carried out with both a crude reaction sample and a sample purified by silica gel chromatography. High performance liquid chromatography (HPLC) analyses were performed on a Hewlett-Packard 1100 Series instrument equipped with an isostatic pump, using a Daicel Chiralpak AS  $(250 \times 4.6 \text{ mm})$ , AD (250×4.6 mm) or Hypersil SI (200×4.6 mm) Column. UV detection was monitored at 254 nm or 220 nm.

# 4.2. Typical procedure for enantioselective addition of $\alpha$ -aryl (10A–I) or alkyl $\alpha$ -cyanoacetates (10J–K) to vinyl sulfone 4a or 4b with 5c

At  $-25 \circ$ C or 0 °C, to a solution of  $\alpha$ -cyanoacetate **10** (for **10A–C**, using 0.6 mmol; for **10D–I**, using 0.5 mmol, for **10J–K**, using 0.2 mmol) in toluene (0.4 mL) was added either Q-**5c** or QD-**5c** (20 mol%) and vinyl phenyl sulfone **4a** or **4b** (0.2 mmol) as specified in Tables 3 and 4. The resulting mixture was kept at the indicated temperature until sulfone is completely consumed. The reaction mixture was directly subjected to silica gel flash chromatography using the eluent specified below to afford the 1,4-adduct in the yields and enantiomeric excess summarized above. Finally, the column was washed with MeOH and the catalyst was recovered in greater than 95% yield. The recovered catalyst was shown to be identical to that before the reaction by NMR analysis.

#### 4.2.1. Compound (+)-11A (Table 3, entry 1)

This product was obtained as a colorless oil in 89% yield after flash chromatography (elution gradient: ethyl acetate/hexane=1:8 to 6:1) and 95% ee as determined by HPLC analysis [Daicel chiralcel OD, hexanes/IPA=90:10, 1.0 mL/min,  $\lambda$ =220 nm,  $t_{\rm R}$  (major)=16.48 min,  $t_{\rm R}$  (minor)=14.93 min] from a reaction catalyzed by Q-**5c** (20 mol %) at -25 °C for 72 h.  $[\alpha]_D^{25}$  +31.6 (*c* 1.0, CHCl<sub>3</sub>); <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$  7.89 (d, *J*=8.0 Hz, 2H), 7.69 (t, *J*=7.2 Hz, 1H), 7.59 (t, *J*=8.0 Hz, 2H), 7.45-7.39 (m, 5H), 4.28-4.15 (m, 2H), 3.27 (dt, *J*=4.4, 12.8 Hz, 1H), 3.04 (dt, *J*=4.4, 12.8 Hz, 1H), 2.75-2.59 (m, 2H), 1.21 (t, *J*=7.2 Hz, 3H); <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>)  $\delta$  166.2, 138.3, 134.1, 132.6, 129.54, 129.48, 128.0, 125.8, 117.0, 63.7, 52.3, 52.1, 30.8, 13.7; HRMS (CI) *m*/z calcd for (C<sub>19</sub>H<sub>19</sub>NO<sub>4</sub>S+H<sup>+</sup>): 358.1113, found: 358.1108; IR (CHCl<sub>3</sub>)  $\nu$  3064, 2984, 2939, 1745, 1448, 1321, 1237, 1087 cm<sup>-1</sup>.

# 4.2.2. Compound (-)-11A

This product was obtained as a colorless oil in 80% yield and 91% ee from a reaction catalyzed by QD-5c (20 mol %) at -25 °C for 72 h.

#### 4.2.3. Compound (+)-11B (Table 3, entry 2)

This product was obtained as a colorless oil in 96% yield after flash chromatography (ethyl acetate/hexane=1:7) and 93% ee as determined by HPLC analysis [Daicel Chiralpak AD, hexanes/ IPA=85:15, 1.0 mL/min,  $\lambda$ =220 nm,  $t_R$  (major)=13.33 min,  $t_R$ (minor)=22.06 min] from a reaction catalyzed by Q-**4c** (20 mol%) at 0 °C for 48 h. [ $\alpha$ ] $_{D}^{55}$  +31.4 (*c* 1.02, CHCl<sub>3</sub>); <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$  7.89 (d, *J*=8.0 Hz, 2H), 7.69 (t, *J*=7.2 Hz, 1H), 7.58 (t, *J*=8.0 Hz, 2H), 7.30 (d, *J*=8.4 Hz, 2H), 7.20 (d, *J*=8.4 Hz, 2H), 4.27– 4.11 (m, 2H), 3.25 (dt, *J*=5.6, 12.8 Hz, 1H), 3.03 (dt, *J*=4.4, 12.8 Hz, 1H), 2.72–2.58 (m, 2H), 2.35 (s, 3H), 1.21 (t, *J*=7.2 Hz, 3H); <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>)  $\delta$  166.3, 139.5, 138.3, 134.0, 130.1, 129.5, 129.4, 127.9, 125.6, 117.2, 63.6, 52.06, 51.98, 30.7, 20.9, 13.6; HRMS (CI) *m/z* calcd for (C<sub>20</sub>H<sub>21</sub>NO<sub>4</sub>S+H<sup>+</sup>): 372.1270, found: 372.1276; IR (CHCl<sub>3</sub>)  $\nu$  2983, 1744, 1447, 1321, 1235, 1148, 1087 cm<sup>-1</sup>.

### 4.2.4. Compound (+)-11C (Table 3, entry 3)

This product was obtained as a colorless oil in 92% yield after flash chromatography (ethyl acetate/hexane=1:6) and 93% ee as determined by HPLC analysis [Daicel Chiralpak AD, hexanes/ IPA=85:15, 0.9 mL/min,  $\lambda$ =220 nm,  $t_R$  (major)=28.93 min,  $t_R$ (minor)=33.20 min] from a reaction catalyzed by Q-**5c** (20 mol%) at 0 °C for 70 h.  $[\alpha]_D^{55}$  +31.0 (*c* 1.10, CHCl<sub>3</sub>); <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$  7.89 (d, *J*=7.2 Hz, 2H), 7.69 (t, *J*=7.2 Hz, 1H), 7.58 (t, 7.2 Hz, 2H), 7.34 (td, *J*=2.4, 9.2 Hz, 2H), 6.90 (td, *J*=2.4, 9.2 Hz, 2H), 4.27– 4.14 (m, 2H), 3.81 (s, 3H), 3.25 (dt, *J*=5.2, 12.0 Hz, 1H), 3.04 (dt, *J*=4.0, 12.0 Hz, 1H), 2.72–2.57 (m, 2H), 1.21 (t, *J*=6.8 Hz, 3H); <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>)  $\delta$  166.4, 160.2, 138.3, 134.1, 129.4, 128.0, 127.1, 124.2, 117.2, 114.8, 63.6, 55.3, 52.1, 51.6, 30.7, 13.7; HRMS (CI) *m/z* calcd for (C<sub>20</sub>H<sub>21</sub>NO<sub>5</sub>S+H<sup>+</sup>): 388.1219, found: 388.1210; IR (CHCl<sub>3</sub>) *v* 2982, 1743, 1512, 1307, 1257, 1236, 1147 cm<sup>-1</sup>.

# 4.2.5. Compound (+)-11D (Table 3, entry 4)

This product was obtained as a colorless oil in 90% yield after flash chromatography (ethyl acetate/hexane=1:5) and 94% ee as determined by HPLC analysis [Daicel chiralcel OD, hexane/ IPA=90:10, 1.0 mL/min,  $\lambda$ =220 nm,  $t_{\rm R}$  (major)=21.42 min,  $t_{\rm R}$ (minor)=17.98 min] from a reaction catalyzed by Q-5c (20 mol %) at -25 °C for 72 h.  $[\alpha]_D^{25}$  +27.2 (c 1.13, CHCl<sub>3</sub>); <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$  7.90–7.88 (m, 2H), 7.70 (tt, *J*=1.2, 7.6 Hz, 1H), 7.59 (t, J=7.2 Hz, 2H), 7.45-7.41 (m, 2H), 7.11 (tt, J=1.6, 8.0 Hz, 2H), 4.29-4.16 (m, 2H), 3.26 (dt, *J*=4.8, 13.6 Hz, 1H), 3.03 (dt, *J*=3.6, 12.8 Hz, 1H), 2.72 (dt, J=4.4, 14.0 Hz, 1H), 2.61 (dt, J=4.4, 14.0 Hz, 1H), 1.22 (t, J=7.2 Hz, 3H); <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>)  $\delta$  166.0, 163.0 (d,  ${}^{1}J_{CF}$ =249.0 Hz), 138.3, 134.1, 129.5, 128.4 (d,  ${}^{4}J_{CF}$ =3.0 Hz), 128.0, 127.8 (d, <sup>3</sup>J<sub>CF</sub>=8.3 Hz), 116.9, 116.6 (d, <sup>2</sup>J<sub>CF</sub>=22.1 Hz), 63.9, 52.0, 51.7, 30.8, 13.6; HRMS (CI) *m*/*z* calcd for (C<sub>19</sub>H<sub>18</sub>FNO<sub>4</sub>S+H<sup>+</sup>): 376.1019, found: 376.1015; IR (CHCl<sub>3</sub>) v 3070, 2985, 2940, 1745, 1604, 1510, 1447, 1307, 1237, 1087 cm<sup>-1</sup>.

# 4.2.6. Compound (+)-11E (Table 3, entry 5)

This product was obtained as a colorless oil in 95% yield after flash chromatograph (ethyl acetate/hexane=1:6) and 94% ee as determined by HPLC analysis [Daicel chiralcel OD, hexane/IPA=85:15, 1.0 mL/min,  $\lambda$ =220 nm,  $t_R$  (major)=16.75 min,  $t_R$  (minor)=13.57 min] from a reaction catalyzed by Q-**5c** (20 mol %) at -25 °C for 69 h.  $[\alpha]_D^{55}$  +31.9 (*c* 1.10, CHCl<sub>3</sub>); <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$  7.85 (d, *J*=8.8 Hz, 2H), 7.65 (t, *J*=7.2 Hz, 1H), 7.55 (t, *J*=6.4 Hz, 2H), 7.34 (s, 4H), 4.22-4.14 (m, 2H), 3.23 (dt, *J*=4.4, 12.8 Hz, 1H), 2.56 (dt, *J*=4.4, 12.8 Hz, 1H), 1.17 (t, *J*=7.2 Hz, 3H); <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>)  $\delta$  165.8, 138.2, 135.6, 134.1, 131.1, 129.6, 129.4, 127.9, 127.2, 116.6, 63.9, 51.9, 51.8, 30.6, 13.6; HRMS (CI) *m/z* calcd for (C<sub>19</sub>H<sub>18</sub>ClNO<sub>4</sub>S+H<sup>+</sup>): 392.0723, found: 392.0728; IR (CHCl<sub>3</sub>)  $\nu$  3066, 2984, 1745, 1493, 1447, 1322, 1236, 1148 cm<sup>-1</sup>.

# 4.2.7. Compound (-)-11E

This product was obtained as a colorless oil in 94% yield and 89% ee from a reaction catalyzed by QD-**5c** (20 mol %) at -25 °C for 72 h.

# 4.2.8. Compound (+)-11F (Table 3, entry 6)

This product was obtained as a colorless oil in 95% yield after flash chromatography (ethyl acetate/hexane=1:6) and 94% ee as determined by HPLC analysis [Daicel chiralpak AD, hexane/ IPA=85:15, 1.0 mL/min,  $\lambda$ =220 nm,  $t_R$  (major)=22.51 min,  $t_R$ (minor)=26.82 min] from a reaction catalyzed by Q-**5c** (20 mol%) at  $-25 \degree C$  for 66 h.  $[\alpha]_{D}^{25} + 28.8$  (*c* 1.10, CHCl<sub>3</sub>); <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$  7.89 (d, *J*=8.0 Hz, 2H), 7.70 (t, *J*=8.0 Hz, 1H), 7.60 (t, *J*=8.0 Hz, 2H), 7.54 (d, *J*=8.8 Hz, 2H), 7.32 (d, *J*=8.8 Hz, 2H), 4.29–4.16 (m, 2H), 3.25 (dt, *J*=4.4, 13.2 Hz, 1H), 3.01 (dt, *J*=4.0, 13.2 Hz, 1H), 2.71 (dt, *J*=4.4, 13.2 Hz, 1H), 2.60 (dt, *J*=4.0, 13.2 Hz, 1H), 1.23 (t, *J*=7.2 Hz, 3H); <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>)  $\delta$  165.8, 138.2, 134.2, 132.7, 131.6, 129.5, 128.0, 127.5, 123.9, 116.6, 64.0, 52.01, 51.96, 30.7, 13.7; HRMS (CI) *m*/*z* calcd for (C<sub>19</sub>H<sub>18</sub>BrNO<sub>4</sub>S+H<sup>+</sup>): 436.0218, found: 436.0208; IR (CHCl<sub>3</sub>)  $\nu$  2983, 1745, 1489, 1447, 1322, 1236, 1148 cm<sup>-1</sup>.

# 4.2.9. Compound (-)-11F

This product was obtained as a colorless oil in 95% yield and 88% ee from a reaction catalyzed by QD-5c (20 mol %) at -25 °C for 72 h.

#### 4.2.10. Compound (+)-11G (Table 3, entry 7)

This product was obtained as a colorless oil in 96% yield after flash chromatography (ethyl acetate/hexane=1:7) and 93% ee as determined by HPLC analysis [Daicel chiralpak AD, hexane/IPA=85:15, 1.0 mL/min,  $\lambda$ =220 nm,  $t_R$  (major)=15.00 min,  $t_R$  (minor)=12.58 min] from a reaction catalyzed by Q-**5c** (20 mol %) at -25 °C for 60 h. [ $\alpha$ ] $_D^{25}$  +29.5 (*c* 1.00, CHCl<sub>3</sub>); <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$  7.91-7.88 (m, 2H), 7.73-7.68 (m, 1H), 7.60 (t, *J*=7.2 Hz, 2H), 7.42-7.33 (m, 4H), 4.31-4.17 (m, 2H), 3.27 (dt, *J*=4.4, 13.6 Hz, 1H), 3.04 (dt, *J*=3.6, 12.8 Hz, 1H), 2.73 (dt, *J*=4.8, 12.8 Hz, 1H), 2.58 (dt, *J*=4.4, 12.8 Hz, 1H), 1.23 (t, *J*=6.8 Hz, 3H); <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>)  $\delta$  165.7, 138.2, 135.6, 134.5, 134.2, 130.8, 129.8, 129.5, 128.0, 126.1, 124.0, 116.5, 64.1, 52.0, 30.9, 13.7; HRMS (CI) *m/z* calcd for (C<sub>19</sub>H<sub>18</sub>ClNO<sub>4</sub>S+H<sup>+</sup>): 392.0723, found: 392.0721; IR (CHCl<sub>3</sub>)  $\nu$  2984, 1746, 1478, 1447, 1309, 1322, 1237, 1148 cm<sup>-1</sup>.

# 4.2.11. Compound (+)-11H (Table 3, entry 8)

This product was obtained as a colorless oil in 96% yield after flash chromatography (elution gradient: ethyl acetate/hexane=1:6) and 98% ee as determined by HPLC analysis [Daicel chiralcel OD, hexane/IPA=90:10, 1.0 mL/min,  $\lambda$ =220 nm,  $t_{\rm R}$  (major)=23.90 min,  $t_{\rm R}$  (minor)=27.21 min] from a reaction catalyzed by Q-**5c** (20 mol %) at -25 °C for 60 h. [ $\alpha$ ]<sub>D</sub><sup>25</sup> +46.7 (*c* 1.00, CHCl<sub>3</sub>); <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$  7.97 (d, *J*=1.6 Hz, 1H), 7.88-7.84 (m, 5H), 7.68-7.64 (m, 1H), 7.57 (m, 4H), 7.44 (dd, *J*=2.4, 8.4 Hz, 1H), 4.29-4.15 (m, 2H), 3.32 (dt, *J*=4.8, 13.6 Hz, 1H), 3.04 (dt, *J*=4.4, 12.4 Hz, 1H), 2.87-2.71 (m, 2H), 1.20 (t, *J*=7.6 Hz, 3H); <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>)  $\delta$  166.2, 138.3, 134.0, 133.1, 132.9, 129.7, 129.6, 129.4, 128.2, 127.9, 127.6, 127.4, 127.2, 125.9, 122.0, 117.1, 63.8, 52.5, 52.1, 30.6, 13.6; HRMS (CI) *m/z* calcd for (C<sub>23</sub>H<sub>21</sub>NO<sub>4</sub>S+H<sup>+</sup>): 408.1270, found: 408.1266; IR (CHCl<sub>3</sub>)  $\nu$  3061, 2983, 1744, 1321, 1234, 1151, 1087 cm<sup>-1</sup>.

# 4.2.12. Compound (-)-11H

This product was obtained as a colorless oil in 95% yield and 90% ee from a reaction catalyzed by QD-**5c** (20 mol %) at -25 °C for 60 h.

## 4.2.13. Compound (+)-**11I** (Table 3, entry 9)

This product was obtained as a brown oil in 95% yield after flash chromatography (ethyl acetate/hexane=1:6) and 93% ee as determined by HPLC analysis [Daicel chiralcel OD, hexane/IPA=90:10, 1.0 mL/min,  $\lambda$ =220 nm,  $t_R$  (major)=20.54 min,  $t_R$  (minor)=18.50 min] from a reaction catalyzed by Q-**5c** (20 mol %) at -25 °C for 48 h.  $[\alpha]_D^{25}$  +20.2 (c 1.00, CHCl<sub>3</sub>); <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$  7.91-7.88 (m, 2H), 7.72-7.68 (m, 1H), 7.59 (t, J=8.0 Hz, 2H), 7.36 (dd, J=1.2, 4.8 Hz, 2H), 7.20 (dd, J=1.2, 4.0 Hz, 1H), 6.99 (dd, J=3.6, 4.8 Hz, 1H), 4.29-4.23 (m, 2H), 3.27 (dt, J=4.8, 12.8 Hz, 1H), 3.14 (dt, J=4.4, 12.4 Hz, 1H), 2.78-2.61 (m, 2H), 1.27 (t, J=7.6 Hz, 3H); <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>)  $\delta$  165.5, 138.2, 134.9, 134.1, 129.4, 127.9, 127.5, 127.4, 127.1, 116.3, 64.0, 52.0, 49.3, 32.1, 13.6; HRMS (CI) m/z calcd for (C<sub>17</sub>H<sub>17</sub>NO<sub>4</sub>S<sub>2</sub>+H<sup>+</sup>): 364.0677, found: 364.0669.

# 4.2.14. Compound (-)-111

This product was obtained as a brown oil in 91% yield and 88% ee from a reaction catalyzed by QD-**5c** (20 mol %) at -25 °C for 48 h.

# 4.2.15. Compound (+)-12J (Table 4, entry 3)

This product was obtained as a white solid in 76% yield after flash chromatography (ethyl acetate/hexane=1:10) and in 93% ee as determined by HPLC analysis [Daicel Chiralcel OD, hexane/IPA=90:10, 1.0 mL/min,  $\lambda$ =220 nm,  $t_R$  (major)=9.78 min,  $t_R$  (minor)=13.73 min] from the reaction catalyzed by Q-**5c** (20 mol %) at 0 °C for 96 h.  $[\alpha]_D^{25}$  +1.6 (*c* 1.5, CHCl<sub>3</sub>); mp 96–98 °C; <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$  8.36 (s, 2H), 8.18 (s, 1H), 5.66–5.78 (m, 2H), 5.20–5.30 (m, 1H), 4.18–4.32 (m, 2H), 3.36 (dt, *J*=12.8, 4.4 Hz, 1H), 3.32 (dt, *J*=12.8, 4.4 Hz, 1H), 2.68 (dd, *J*=13.6, 6.8 Hz, 1H), 2.65 (dd, *J*=13.6, 6.8 Hz, 1H), 2.40 (dt, *J*=12.4, 4.4 Hz, 1H), 1.29 (t, *J*=6.8, 3H); <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>)  $\delta$  166.8, 141.4, 133.5 (q, *J*=34 Hz), 129.2, 128.5, 127.9, 122.3 (q, *J*=273.0 Hz), 122.2, 117.2, 63.6, 52.3, 47.4, 41.3, 28.0, 13.9; IR (CHCl<sub>3</sub>)  $\nu$  3088, 2987, 1739, 1360, 1279, 1185, 1147 cm<sup>-1</sup>; HRMS (CI) *m/z* calcd for (C<sub>18</sub>H<sub>17</sub>F<sub>6</sub>NO<sub>4</sub>S+H<sup>+</sup>): 458.0861, found 458.0862.

The absolute configuration of adduct (+)-**12J** was determined to be 'S' by X-ray structure analysis, see Section 4.5 for details.

# 4.2.16. Compound (-)-12K (Table 4, entry 4)

This product was obtained as a white solid in 85% yield after flash chromatography (ethyl ether/hexane=1:5) and in 92% ee as determined by HPLC analysis [Daicel Chiralcel OD, hexane/IPA=90:10, 1.0 mL/min,  $\lambda$ =220 nm,  $t_R$  (major)=10.88 min,  $t_R$  (minor)=17.64 min] from the reaction catalyzed by Q-**5c** (20 mol %) at 0 °C for 96 h.  $[\alpha]_D^{25}$  -1.0 (*c* 2.0, CHCl<sub>3</sub>); mp 64–66 °C; <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$  8.40 (s, 2H), 8.21 (s, 1H), 4.28 (dq, *J*=1.6, 7.2 Hz, 2H), 3.38 (dt, *J*=13.6, 4.4 Hz, 1H), 3.31 (dt, *J*=12.8, 4.4 Hz, 1H), 2.45 (dt, *J*=12.8, 4.4 Hz, 1H), 2.27 (dt, *J*=12.8, 4.4 Hz, 1H), 1.67 (s, 3H), 1.32 (t, *J*=6.8 Hz, 3H); <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>)  $\delta$  167.6, 141.4, 133.5 (q, *J*=30 Hz), 128.5, 128.0, 127.9, 122.1 (q, *J*=272.4 Hz), 118.2, 63.1, 52.4, 42.4, 29.8, 23.7, 13.9; IR (CHCl<sub>3</sub>)  $\nu$  3088, 2990, 1745, 1281, 1186, 1147 cm<sup>-1</sup>; HRMS (CI) *m/z* calcd for (C<sub>16</sub>H<sub>15</sub>F<sub>6</sub>NO<sub>4</sub>S+H<sup>+</sup>): 432.0708, found 432.0712.

# 4.2.17. Compound (+)-**12K**

This product was obtained as a white solid in 83% yield and 88% ee from a reaction catalyzed by QD-**5c** (20 mol %) at 0 °C for 96 h.

# 4.3. Typical procedure for enantioselective additions of $\alpha$ -cyanoketones (13A–C) or $\beta$ -ketoesters (14A and B) to vinyl sulfone 4a with 5c

To a mixture of **4a** (0.20 mmol) and **5c** (20 mol%) in toluene (0.40 mL) was added **13** (0.20 mmol) or **14** (0.40 mmol) in one portion. The resulting reaction mixture was kept at either room temperature or 0 °C from 24 h to 96 h. The reaction mixture was purified by silica gel chromatography directly using the eluent specified below.

# 4.3.1. Compound (-)-15A (Table 5, entry 1)

This product was obtained as a white solid in 96% yield after flash chromatography (silica gel, ethyl acetate/hexanes=1:10) and in 95% ee as determined by HPLC [determined by HPLC, Daiso Chiralpak AD, hexanes/IPA=80:20, 1.0 mL/min,  $\lambda$ =220 nm, t<sub>R</sub> (major)=32.66 min, t<sub>R</sub> (minor)=27.03 min] from a reaction of **13A** (43.6 mg, 0.40 mmol) and vinyl sulfone **4a** (33.6 mg, 0.20 mmol) in toluene (400 µL) catalyzed by Q-**5c** (0.04 mmol) at room temperature for 24 h. [ $\alpha$ ]<sub>D</sub><sup>25</sup> –12.8 (*c* 1.0, CHCl<sub>3</sub>); mp 113–115 °C; <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$  1.98–2.18 (m, 4H), 2.25 (dt, *J*=4.4, 12.4 Hz, 1H), 2.36–2.46 (m, 1H), 2.46–2.58 (m, 2H), 3.265 (dt, *J*=4.4, 12.4 Hz, 1H), 3.47 (dt, *J*=4.4, 12.4 Hz, 1H), 7.61 (t, *J*=7.6 Hz, 2H), 7.71 (d, *J*=7.6 Hz, 2H)

1H), 7.90–7.96 (m, 2H);  $^{13}$ C NMR (100 MHz, CDCl<sub>3</sub>)  $\delta$  19.2, 26.6, 35.0, 36.3, 46.7, 51.9, 117.5, 128.0, 129.5, 134.2, 138.4, 207.9; HRMS [M+H]<sup>+</sup> calcd for: C<sub>14</sub>H<sub>16</sub>O<sub>3</sub>SN: 278.0851, found: 278.0854.

# 4.3.2. Compound (+)-15A

This product was obtained as a white solid in 90% isolated yield and in 95% ee from a reaction of **13A** (43.6 mg, 0.40 mmol) and vinyl sulfone **4a** (33.6 mg, 0.20 mmol) in toluene (400  $\mu$ L) catalyzed by QD-**5c** (0.04 mmol) at room temperature for 24 h.

# 4.3.3. Compound (-)-15B (Table 5, entry 2)

This product was obtained as a white solid in 96% yield after flash chromatography and in 95% ee as determined by HPLC [determined by HPLC, Daiso Chiralcel OJ, hexanes/IPA=70:30, 1.0 mL/min,  $\lambda$ =220 nm,  $t_{\rm R}$  (major)=42.83 min,  $t_{\rm R}$  (minor)=36.97 min] from a reaction of **13B** (49.2 mg, 0.40 mmol) and vinyl sulfone **4a** (33.6 mg, 0.20 mmol) in toluene (400 µL) catalyzed by Q-**5c** (0.04 mmol) at room temperature for 24 h.  $[\alpha]_{\rm D}^{25}$  –97.1 (c 1.1, CHCl<sub>3</sub>); mp 68–70 °C; <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$  1.64–1.76 (m, 2H), 1.86–1.96 (m, 1H), 1.98–2.10 (m, 2H), 2.10–2.18 (m, 1H), 2.28–2.36 (m, 2H), 2.46 (dt, *J*=3.6, 14.0 Hz, 1H), 2.79 (dt, *J*=6.4, 20.0 Hz, 1H), 3.22 (dt, *J*=4.4, 13.2 Hz, 1H), 3.39 (dt, *J*=4.4, 13.2 Hz, 1H), 7.61 (t, *J*=7.6 Hz, 2H), 7.71 (m, 1H), 7.90–7.96 (m, 2H); <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>)  $\delta$  22.1, 26.8, 27.7, 39.0, 39.1, 49.9, 52.0, 118.3, 128.0, 129.4, 134.0, 138.5, 201.9; HRMS [M+H]<sup>+</sup> calcd for: C<sub>15</sub>H<sub>18</sub>O<sub>3</sub>SN: 292.1007, found: 292.1008.

## 4.3.4. Compound (+)-15B

This product was obtained as a white solid in 98% isolated yield and in 95% ee from a reaction of 1**3B** (49.2 mg, 0.40 mmol) and vinyl sulfone **4a** (33.6 mg, 0.20 mmol) in toluene (400  $\mu$ L) catalyzed by QD-**5C** (0.04 mmol) at room temperature for 24 h.

## 4.3.5. Compound (-)-15C (Table 5, entry 3)

This product was obtained as a white solid in 91% yield after flash chromatography and in 96% ee as determined by HPLC [determined by HPLC, Daiso Chiralcel OJ, hexanes/IPA=70:30, 1.0 mL/min,  $\lambda$ =220 nm,  $t_R$  (major)=16.27 min,  $t_R$  (minor)=22.27 min] from a reaction of **13C** (54.8 mg, 0.40 mmol) and vinyl sulfone **4a** (33.6 mg, 0.20 mmol) in toluene (400 µL) catalyzed by Q-**5c** (0.04 mmol) at room temperature for 24 h.  $[\alpha]_D^{25}$  –16.2 (*c* 1.0, CHCl<sub>3</sub>); mp 153–156 °C; <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$  1.24–1.36 (m, 1H), 1.58–1.70 (m, 2H), 1.72–1.82 (m, 1H), 1.88–1.98 (m, 3H), 2.10–2.22 (m, 2H), 2.32 (dq, *J*=4.4, 8.0 Hz, 1H), 2.61 (dq, *J*=4.8, 10.8 Hz, 1H), 2.72 (dt, *J*=5.2, 13.6 Hz, 1H), 3.16 (dt, *J*=4.4, 13.6 Hz, 1H), 3.27 (dt, *J*=4.4, 13.6 Hz, 1H), 7.61 (t, *J*=7.6 Hz, 2H), 7.67–7.72 (m, 1H), 7.90–7.94 (m, 2H); <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>)  $\delta$  24.1, 25.8, 28.2, 28.9, 36.2, 40.6, 52.0, 53.2, 118.7, 127.9, 129.5, 134.1, 138.4, 204.0; HRMS [M+H]<sup>+</sup> calcd for: C<sub>16</sub>H<sub>20</sub>O<sub>3</sub>SN: 306.1163, found: 306.1161.

The absolute configuration of (-)-**15C** was determined to be 'S' by X-ray structure analysis, see Section 4.5 for details.

# 4.3.6. Compound (+)-15C

This product was obtained as a white solid in 91% isolated yield and in 97% ee from a reaction of **13C** (54.8 mg, 0.40 mmol) and vinyl sulfone (33.6 mg, 0.20 mmol) in toluene (400  $\mu$ L) catalyzed by QD-**2c** (20 mol%) at room temperature for 24 h.

#### 4.3.7. Compound (-)-16A (Table 5, entry 5)

This product was obtained as a white solid in 93% yield after flash chromatography (silica gel, ethyl ether/hexanes=1:2) and in 96% ee as determined by HPLC [determined by HPLC, Daiso Chiralpak AD, hexanes/IPA=90:10, 1.0 mL/min,  $\lambda$ =220 nm,  $t_R$  (major)=15.64 min,  $t_R$  (minor)=19.24 min] from a reaction of **14a** (73.6 mg, 0.40 mmol) and vinyl sulfone **4a** (33.6 mg, 0.20 mmol) in toluene (400 µL) catalyzed by Q-**5c** (0.04 mmol) at room temperature for 72 h.  $[\alpha]_D^{25}$  –7.3

(c 1.00, CHCl<sub>3</sub>); <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$  0.84 (t, *J*=6.8 Hz, 3H), 1.04–1.54 (m, 8H), 1.56–1.78 (m, 1H), 2.84–2.96 (m, 1H), 3.20 (d, *J*=17.6 Hz, 1H), 3.40 (dd, *J*=6.8, 6.8 Hz, 1H), 3.70 (s, 3H), 3.87 (d, *J*=17.6 Hz, 1H), 4.12 (dd, *J*=6.8, 6.8 Hz, 1H), 7.43 (t, *J*=7.2 Hz, 1H), 7.53 (d, *J*=8.0 Hz, 1H), 7.67 (dt, *J*=1.2, 8.0 Hz, 1H), 7.8 (d, *J*=8.0 Hz, 1H); <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>)  $\delta$  19.5, 26.5, 27.7, 34.4, 33.7, 52.0, 58.6, 82.6, 128.0, 129.3, 133.7, 138.7, 169.6, 214.2; IR (neat)  $\nu$  2977, 1746, 1719, 1147; HRMS [M+Na]<sup>+</sup> calcd for: C<sub>18</sub>H<sub>24</sub>O<sub>5</sub>SNa: 375.1242, found: 375.1233.

# 4.3.8. Compound (+)-16A

This product was obtained as a white solid in 96% isolated yield and in 96% ee from a reaction of **14A** (73.6 mg, 0.40 mmol) and vinyl sulfone **4a** (33.6 mg, 0.20 mmol) in toluene (400  $\mu$ L) catalyzed by QD-**5c** (0.04 mmol) at room temperature for 72 h.

## 4.3.9. Compound (-)-16B (Table 5, entry 6)

This product was obtained as a white solid in 100% yield after flash chromatography (silica gel, ethyl ether/hexanes=2:3) and in 97% ee as determined by HPLC [determined by HPLC, Daiso Chiralpak AD, hexanes/IPA=90:10, 1.0 mL/min,  $\lambda$ =220 nm, t<sub>R</sub> (major)=28.57 min,  $t_R$  (minor)=24.85 min] from a reaction of **14B** (93.0 mg, 0.40 mmol) and vinyl sulfone **4a** (33.6 mg, 0.20 mmol) in toluene (400  $\mu$ L) catalyzed by Q-5c (0.04 mmol) at room temperature for 36 h.  $[\alpha]_D^{25}$  –8.5 (*c* 2.00, CHCl<sub>3</sub>); mp 114–116 °; <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 0.84 (t, J=6.8 Hz, 3H), 1.04–1.54 (m, 8H), 1.56– 1.78 (m, 1H), 2.84–2.96 (m, 1H), 3.20 (d, *J*=17.6 Hz, 1H), 3.40 (dd, *I*=6.8, 6.8 Hz, 1H), 3.70 (s, 3H), 3.87 (d, *I*=17.6 Hz, 1H), 4.12 (dd, *I*=6.8, 6.8 Hz, 1H), 7.43 (t, *I*=7.2 Hz, 1H), 7.53 (d, *I*=8.0 Hz, 1H), 7.67 (dt, *J*=1.2, 8.0 Hz, 1H), 7.8 (d, *J*=8.0 Hz, 1H); <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>) § 26.6, 27.7, 37.86, 52.2, 59.1, 82.6, 124.9, 126.4, 128.1, 129.3, 133.8, 134.7, 135.6, 138.7, 140.0, 152.3, 169.3, 201.7; IR (neat) v 2978, 1733, 1707, 1149; HRMS [M+Na]<sup>+</sup> calcd for: C<sub>22</sub>H<sub>24</sub>O<sub>5</sub>SNa: 423.1242, found: 423.1245.

#### 4.3.10. Compound (+)-16B

This product was obtained as a white solid in 96% isolated yield and in 96% ee from a reaction of **14B** (93.0 mg, 0.40 mmol) and vinyl sulfone **4a** (33.6 mg, 0.20 mmol) in toluene (400  $\mu$ L) catalyzed by QD-**5c** (0.04 mmol) at room temperature for 36 h.

# 4.4. Typical procedure for enantioselective additions of $\alpha$ -cyanocyanoacetates 10, $\alpha$ -cyanoketones 13 and $\beta$ -ketoesters 14 to $\beta$ -substituted vinyl triflones 17 with 5c

To a mixture of **17** (0.10 mmol) and **5c** (0.01 mmol) in dichloromethane (0.2 mL) was added **10** or **13** or **14** in one portion. The resulting reaction mixture was kept at -20 or -50 °C from 24 h to 96 h. The reaction mixture was purified by silica gel chromatography directly using the eluent specified below.

# 4.4.1. Compound (+)-**18Ab** (Table 7, entry 1)

This product was obtained from a Q-**5c** (4.8 mg, 0.01 mmol, 10 mol%) catalyzed reaction of **17b** (23.0 mg, 0.10 mmol) and **10A** (37.8 mg, 0.20 mmol) in methylene chloride (200  $\mu$ L) at  $-50 \degree$ C for 96 h. The dr (5.6:1) was determined by an analysis of the crude product via integrations of one set of <sup>19</sup>F NMR (376 MHz, CF<sub>3</sub>COOH=-78.5 ppm) signals (s,  $\delta$  (major)=-81.86 ppm,  $\delta$  (minor)=-82.06 ppm)]. The crude product was purified by flash chromatography (silica gel, hexanes/ethyl ether=20:1) to give the major diastereomer of the 1,4-adduct as a colorless oil (36 mg, 85% yield, single diastereomer) and in 93% ee [determined by HPLC, Daiso Chiralpak AD, hexanes/IPA=98:2, 1.0 mL/min,  $\lambda$ =220 nm,  $t_R$  (major)=5.70 min,  $t_R$  (minor)=6.75 min]. [ $\alpha$ ]<sub>D</sub><sup>5</sup> +12.0 (*c* 1.00, CHCl<sub>3</sub>); <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$  0.75 (t, *J*=7.2 Hz, 3H), 0.94–1.14 (m, 5H), 1.23 (t, *J*=6.4 Hz, 3H), 1.22–1.34 (m, 1H), 1.42–1.52 (m, 2H), 3.20–3.30

(m, 1H), 3.40–3.48 (m, 2H), 4.18–4.42 (m, 2H), 7.38–7.44 (m, 3H); 7.50–7.58 (m, 2H); <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>)  $\delta$  13.9, 14.0, 22.2, 26.1, 31.1, 31.6, 39.5, 51.8, 59.3, 64.3, 116.9, 119.5 (q, *J*=325 Hz), 126.7, 129.7, 130.0, 132.2, 166.5; <sup>19</sup>F NMR (376 MHz, CF<sub>3</sub>COOH=–78.5 ppm) –81.9; IR (neat)  $\nu$  2959, 1746, 1510, 1374, 1228, 1122; HRMS (ESI) [M+H]<sup>+</sup> calcd for: C<sub>19</sub>H<sub>24</sub>F<sub>4</sub>O<sub>4</sub>S: 438.1362, found: 438.1358.

# 4.4.2. Compound (-)-18Ab

This product was obtained as an colorless oil in 76% yield (single diastereomer) and 87% ee from the corresponding QD-**5c**-catalyzed reaction of **17b** (23.0 mg, 0.10 mmol) and **10A** (37.8 mg, 0.20 mmol) in methylene chloride (200  $\mu$ L) at -50 °C for 96 h. The dr (4.6:1) of the reaction was determined by <sup>1</sup>H NMR analysis of the crude product.

# 4.4.3. Compound (+)-18Db (Table 7, entry 2)

This product was obtained from a Q-5c (4.8 mg, 0.01 mmol, 10 mol %) catalyzed reaction of 17b (23.0 mg, 0.10 mmol) and 10D (41.0 mg, 0.20 mmol) in methylene chloride (200  $\mu$ L) at -50 °C for 96 h. The dr (26:1) was determined by an analysis of the crude product via integration of one set of <sup>19</sup>F NMR (376 MHz, CF<sub>3</sub>COOH=-78.5 ppm) signals (s,  $\delta$  (major)=-81.78 ppm,  $\delta$  (minor)=-82.11 ppm)]. The crude product was purified by flash chromatography (silica gel, hexanes/ethyl ether=30:1) to give adduct **18Db** as a colorless oil (41 mg, 94% yield, single diastereomer) and in 92% ee [determined by HPLC, Daiso Chiralpak AD, hexanes/ IPA=98:2, 1.0 mL/min,  $\lambda$ =220 nm,  $t_{\rm R}$  (major)=5.94 min,  $t_{\rm R}$  (minor)= 8.01 min].  $[\alpha]_D^{25}$  +11.0 (c 1.00, CHCl<sub>3</sub>); <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 0.79 (t, J=6.8 Hz, 3H), 1.00-1.20 (m, 5H), 1.26 (t, J=7.0 Hz, 3H), 1.28-1.38 (m, 1H), 1.42-1.52 (m, 2H), 3.20-3.30 (m, 1H), 3.40-3.48 (m, 2H), 4.20–4.36 (m, 2H), 7.16 (t, *J*=8.4 Hz, 2H), 7.50–7.60 (m, 2H); <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>) § 13.9, 14.0, 22.2, 26.1, 31.0, 31.5, 39.5, 51.6, 58.7, 64.5, 116.7, 116.8 (d, J=22 Hz), 119.52 (q, J=325 Hz), 128.00 (d, *J*=3.0 Hz), 128.79 (d, *J*=8.0 Hz), 163.50 (d, *J*=118 Hz), 166.45; IR (neat) ν 2959, 1746, 1510, 1374, 1228, 1122; <sup>19</sup>F NMR (376 MHz, CF<sub>3</sub>COOH=-78.5 ppm)-81.7, -113.8; HRMS(ESI)[M+H]<sup>+</sup> calcd for: C<sub>19</sub>H<sub>24</sub>F<sub>4</sub>O<sub>4</sub>S: 438.1362, found: 438.1358.

## 4.4.4. Compound (-)-18Db

This product was obtained as an colorless oil in 91% yield (single diastereomer) and 84% ee from the corresponding QD-**5c**-catalyzed reaction of **17b** (23.0 mg, 0.10 mmol) and **10D** (41.0 mg, 0.20 mmol) in methylene chloride (200  $\mu$ L) at -50 °C for 96 h. The dr (12:1) of the reaction was determined by <sup>1</sup>H NMR analysis of the crude product.

# 4.4.5. Compound (+)-18Fb (Table 7, entry 3)

This product was obtained from a Q-5c (4.8 mg, 0.010 mmol, 10 mol%) catalyzed reaction of 17b (23.0 mg, 0.10 mmol) and 10F (53.2 mg, 0.20 mmol, 2.0 equiv) in methylene chloride (200 µL) at -50 °C for 96 h. The dr (49:1) was determined by an analysis of the crude product via integration of one set of <sup>19</sup>F NMR  $(376 \text{ MHz}, \text{CF}_3\text{COOH}=-78.5 \text{ ppm})$  signals  $(s, \delta \text{ (major})=-81.69 \text{ ppm})$ ,  $\delta$  (minor)=-81.95 ppm)]. The crude product was purified by flash chromatography (silica gel, hexanes/ethyl ether=30:1) to give adduct **18Fb** as a colorless oil (46 mg, 92% yield, single diastereomer) and in 81% ee [determined by HPLC, Daiso Chiralpak AD, hexanes/ IPA=98:2, 1.0 mL/min,  $\lambda$ =220 nm,  $t_{\rm R}$  (major)=5.92 min,  $t_{\rm R}$  (minor)= 9.91 min].  $[\alpha]_{D}^{25}$  +4.6 (*c* 2.00, CHCl<sub>3</sub>); <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$  0.79 (t, *J*=6.8 Hz, 3H), 1.04–1.20 (m, 5H), 1.26 (t, *J*=7.0 Hz, 3H), 1.30– 1.38 (m, 1H), 1.41-1.52 (m, 2H), 3.20-3.30 (m, 1H), 3.40-3.50 (m, 2H), 4.20–4.36 (m, 2H), 7.46 (d, *J*=9.2 Hz, 2H), 7.60 (d, *J*=8.4 Hz, 2H); <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>) δ 13.6, 13.7, 22.0, 25.8, 30.8, 31.3, 39.1, 51.3, 58.8, 64.4, 116.2, 119.3 (q, *J*=327 Hz), 124.2, 128.2, 131.0, 132.7, 166.0; <sup>19</sup>F NMR (376 MHz, CF<sub>3</sub>COOH=-78.5 ppm) -81.7; IR (neat) *v* 2932, 1746, 1373, 1225, 1122.

# 4.4.6. Compound (-)-**18Fb**

This product was obtained as an colorless oil in 92% yield (single diastereomer) and 86% ee from the corresponding QD-**5c**-catalyzed reaction of **17b** (23.0 mg, 0.10 mmol) and **10F** (53.2 mg, 0.20 mmol) in methylene chloride (200  $\mu$ L) at -50 °C for 96 h. The dr (26:1) of the reaction was determined by <sup>1</sup>H NMR analysis of the crude product.

## 4.4.7. Compound (+)-19Eb (Table 7, entry 4)

This product was obtained from a Q-5c (4.8 mg, 0.010 mmol, 10 mol%) catalyzed reaction of 13E (15.7 mg, 0.10 mmol) and 17b (25.3 mg, 0.11 mmol) in methylene chloride (400  $\mu$ L) at  $-50 \degree$ C for 96 h. The dr (>49:1) was determined by an analysis of the crude product via integration of one set of <sup>19</sup>F NMR (376 MHz, CF<sub>3</sub>COOH=-78.5 ppm) signals (s,  $\delta$  (major)=-81.50 ppm,  $\delta$  (minor)= -81.24 ppm)]. The crude product was purified by flash chromatography (silica gel, hexanes/ethyl acetate=3:1) to give adduct 19Eb as a colorless oil (39 mg, 95% yield, single diastereomer) and in 99% ee [determined by HPLC, Daiso Chiralcel OJ, hexanes/ IPA=90:10, 1.0 mL/min,  $\lambda$ =220 nm,  $t_{\rm R}$  (major)=19.15 min,  $t_{\rm R}$ (minor)=24.02 min].  $[\alpha]_D^{25}$  +75.0 (*c* 1.00, CHCl<sub>3</sub>); <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 0.89 (t, J=6.8 Hz, 3H), 1.20-1.42 (m, 5H), 1.50-1.64 (m, 1H), 1.68–1.88 (m, 2H), 2.52–2.62 (m, 1H), 3.37 (dd, J=6.8, 6.8 Hz, 1H), 3.47 (d, J=17.2 Hz, 1H), 3.73 (d, J=7.2 Hz, 1H), 4.33 (dd, J=3.6, 6.8 Hz, 1H), 7.52 (dd, J=8.0, 8.0 Hz, 2H), 7.75 (dd, J=8.0, 8.0 Hz, 1H), 7.87 (d, J=8.0 Hz, 1H); <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>=77.0 ppm) δ 14.1, 22.5, 26.4, 31.7, 32.6, 37.7, 38.0, 49.2, 50.7, 63.9, 118.2, 119.5 (q, J=325 Hz), 126.27, 126.66, 129.42, 133.44. 137.27, 150.09, 196.55; <sup>19</sup>F NMR (376 MHz, CF<sub>3</sub>COOH=-78.5 ppm) -81.5. IR (neat) v 2934, 1725, 1370, 1206, 1121; HRMS (ESI) [M+H]+ calcd for: C<sub>18</sub>H<sub>21</sub>NF<sub>3</sub>O<sub>5</sub>S: 388.1194, found: 388.1188.

# 4.4.8. Compound (-)-19Eb

This product was obtained as an colorless oil in 94% yield (single diastereomer) and 94% ee from the corresponding QD-**5c**-catalyzed reaction of **17b** (25.3 mg, 0.11 mmol) and **13E** (15.7 mg, 0.10 mmol) in methylene chloride (200  $\mu$ L) at -50 °C for 96 h. The dr (>49:1) of the reaction was determined by <sup>1</sup>H NMR analysis of the crude product.

# 4.4.9. Compound (+)-20Db (Table 7, entry 5)

This product was obtained from a Q-5c (4.8 mg, 0.010 mmol, 10 mol%) catalyzed reaction of 14D (19.0 mg, 0.10 mmol) and 17b (23 mg, 0.10 mmol) in methylene chloride  $(200 \mu L)$  at  $-20 \degree$ C for 96 h. The dr (32:1) was determined by an analysis of the crude product via integration of one set of <sup>19</sup>F NMR (376 MHz, CF<sub>3</sub>COOH=-78.5 ppm) signals (s,  $\delta$  (major)=-81.87 ppm,  $\delta$  (minor)=-81.79 ppm)]. The crude product was purified by flash chromatography (silica gel, hexanes/ethyl ether=5:1) to give adduct 20Db as a colorless oil (38 mg, 90% yield, single diastereomer) and in 98% ee [determined by HPLC, Daiso Chiralcel OD+Regis (R,R)-whelk 1 O, hexanes/IPA=99:1, 1.0 mL/min,  $\lambda = 220$  nm,  $t_R$  (major)=20.15 min,  $t_R$  (minor)=29.10 min].  $[\alpha]_D^{25}$  +116.4 (c 1.00, CHCl<sub>3</sub>); <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$  0.84 (t, *I*=6.8 Hz, 3H), 1.06–1.54 (m, 7H), 1.56–1.70 (m, 1H), 2.82–2.96 (m, 1H), 3.20(d, J=17.6 Hz, 1H), 3.40(dd, J=6.8, 6.8 Hz, 1H), 3.70(s, 3H), 3.87(d, J=17.6 Hz, 1H), 4.12 (dd, J=3.2, 14.4 Hz, 1H), 7.43 (t, J=7.2 Hz, 1H), 7.53 (d, J=8.0 Hz, 1H), 7.67 (dt, J=1.2, 8.0 Hz, 1H), 7.8 (d, J=8.0 Hz, 1H); <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>=77.0 ppm)  $\delta$  14.1, 22.5, 27.4, 31.7, 32.3, 35.8, 37,3, 50.4, 53.4, 63.9, 119.6 (q, J=325 Hz), 125.4, 126.6, 128.4, 135.0, 136.2, 152.9, 170.0, 200.4; <sup>19</sup>F NMR (376 MHz, CF<sub>3</sub>COOH=-78.5 ppm) -81.8; IR (neat) v 2957, 1742, 1717, 1367, 1200, 1122; HRMS (ESI) [M+H]<sup>+</sup> calcd for: C<sub>19</sub>H<sub>24</sub>F<sub>3</sub>O<sub>5</sub>S: 421.1297, found: 421.1289.

# 4.4.10. Compound (-)-20Db

This product was obtained as an colorless oil in 93% yield (contains minor diastereomer, dr=90:10) and 97% ee from the corresponding QD-**5c**-catalyzed reaction of **17b** (23.0 mg, 0.10 mmol) and **10F** (53.2 mg, 0.20 mmol) in methylene chloride

(200  $\mu$ L) at -50 °C for 96 h. The dr (9:1) of the reaction was determined by <sup>1</sup>H NMR analysis of the crude product.

## 4.4.11. Compound **20Dc** (Table 7, entry 6)

This product was obtained from a Q-5c (4.8 mg, 0.010 mmol, 10 mol %) catalyzed reaction of 14D (38.0 mg, 0.20 mmol) and 17c (21.6 mg, 0.10 mmol) in methylene chloride (200  $\mu$ L) at  $-20 \degree$ C for 72 h. The crude product was purified by flash chromatography (silica gel, hexanes/methylene chloride=1:1) to give the adduct **20Dc** as a colorless oil and a diastereomeric mixture (41 mg, 100% yield, dr=24:1). The ee of the major diastereomer is determined to be 99%. Both dr and ee were determined by HPLC [Daiso Chiralcel AD+Chiralcel AD, hexanes/IPA=98:2, 1.0 mL/min,  $\lambda$ =220 nm,  $t_{\rm R}$ (major)=21.10 min,  $t_{\rm R}$  (minor)=27.65 min].  $[\alpha]_{\rm D}^{25}$  +153.0 (c 1.00, CHCl<sub>3</sub>); <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>, major diastereomer was reported) δ 0.90 (d, *J*=6.4 Hz, 3H), 0.91 (d, *J*=6.4 Hz, 3H), 1.02–1.12 (m, 1H), 1.60– 1.76 (m, 2H), 2.92–3.06 (m, 1H), 3.18 (d, *J*=17.2 Hz, 1H), 3.36 (dd, *J*=6.8, 14.4 Hz, 1H), 3.71 (s, 3H), 3.88 (d, J=17.2 Hz, 1H), 4.12 (dd, J=6.8, 14.4 Hz, 1H), 7.44 (t, J=7.2 Hz, 1H), 7.53 (d, J=7.20 Hz, 1H), 7.68 (t, J=6.8 Hz, 1H), 7.80 (d, J=8.0 Hz, 1H); <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>)  $\delta$  21.1, 23.7, 25.1, 34.9, 35.7, 41.3, 50.2, 53.2, 63.6, 119.4 (q, J=327 Hz), 125.2, 126.3, 128.2, 134.7, 134.0, 152.7, 169.7, 200.1; <sup>19</sup>F NMR (376 MHz, CF<sub>3</sub>COOH=-78.5 ppm) -82.0; IR (neat) v 2960, 1742, 1715, 1367, 1200, 1121; HRMS (ESI) [M+H]<sup>+</sup> calcd for: C<sub>18</sub>H<sub>22</sub>F<sub>3</sub>O<sub>5</sub>S: 407.1140, found: 407.1150.

The QD-**5c** (4.8 mg, 0.010 mmol, 10 mol %) catalyzed reaction of **14E** (38.0 mg, 0.20 mmol) and **17c** (21.6 mg, 0.10 mmol) in methylene chloride (200  $\mu$ L) was carried out at -20 °C for 72 h to furnish the crude product (dr=8:1). The crude product was purified by flash chromatography to give the adduct **20Dc** as a colorless oil and a diastereomeric mixture (40 mg, 98% yield, dr=8:1). The ee of the major diastereomer is determined to be 95%.

## 4.4.12. Compound (-)-20Ed (Table 7, entry 7)

This product was obtained from a Q-5c (4.8 mg, 0.010 mmol, 10 mol%) catalyzed reaction of 14E (28.4 mg, 0.20 mmol) and 17d (26.4 mg, 0.10 mmol) in methylene chloride (100  $\mu L)$  at  $-20\ensuremath{\,^\circ C}$  for 96 h. The crude product (single diastereomer by <sup>1</sup>H NMR analysis) [determined by integration of one set of <sup>19</sup>F NMR (376 MHz, CF=-78.5 ppm) signals (s,  $\delta$  (major)=-81.57 ppm)]. The crude product was purified by flash chromatography (silica gel, hexanes/ methylene chloride=2:1) to give adduct 20Ed as a colorless oil (37 mg, 91% yield, single diastereomer) and in 95% ee [determined by HPLC, Daiso Chiralcel OJ, hexanes/IPA=90:10, 1.0 mL/min,  $\lambda = 220 \text{ nm}, t_{\text{R}}(\text{major}) = 17.45 \text{ min}, t_{\text{R}}(\text{minor}) = 14.30 \text{ min}]. [\alpha]_{\text{D}}^{25} - 57.4$ (c 1.00, CHCl<sub>3</sub>); <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 1.70–2.00 (m, 5H), 2.30– 2.42 (m, 2H), 2.52-2.70 (m, 3H), 2.80-2.92 (m, 1H), 3.32 (dd, J=6.4, 14.4 Hz, 1H), 3.71 (s, 3H), 4.25 (dd, J=4.0, 14.4 Hz, 1H), 7.12-7.32 (m, 5H); <sup>13</sup>C NMR (100 MHz, CDCl<sub>3</sub>=77.0 ppm)  $\delta$  19.3, 32.6, 34.0, 34.8, 35.4, 38.4, 50.6, 53.1, 63.2, 119.7 (q, J=325 Hz), 126.5, 128.7, 128.8, 140.9, 170.0, 213.1; IR (neat) v 1751, 1725, 1364, 1200, 1120; HRMS  $(ESI) [M+H]^+$  calcd for: C<sub>18</sub>H<sub>22</sub>F<sub>3</sub>O<sub>5</sub>S: 407.1140, found: 407.1145.

# 4.4.13. Compound (+)-**20Ed**

The QD-**5c** (4.8 mg, 0.010 mmol, 10 mol %) catalyzed reaction of **14E** (38.0 mg, 0.20 mmol) and **17c** (21.6 mg, 0.10 mmol) in methylene chloride (200  $\mu$ L) was carried out at -20 °C for 72 h to furnish the crude product as a single diastereomer (by <sup>1</sup>H NMR analysis). The crude product was purified by flash chromatography to give the adduct **20Ed** as a colorless oil and a single diastereomer (33 mg, 80% yield) in 89% ee.

# **4.5.** X-ray analysis for the determination of the absolute configurations of (+)-12J and (-)-15C

X-ray analysis was performed by Richard J. Staples at the Department of Chemistry and Chemical Biology, Harvard University. Crystallographic data (excluding structure factors) for the structures in this paper have been deposited with the Cambridge Crystallographic Data Centre as supplementary publication nos. CCDC 709230 [(+)-**12J**] and CCDC 709231 [(-)-**15C**]. The absolute configurations of compounds (+)-**12J** and (-)-**15C** were established using the anomalous scattering technique; Flack parameters were -0.06(10) and -0.06(6), respectively. Copies of the data can be obtained, free of charge, on application to CCDC, 12 Union Road, Cambridge CB2 1EZ, UK (fax: +44 (0)1223 336033 or e-mail: deposit@ccdc.cam.ac.uk).

# Acknowledgements

We gratefully acknowledge the financial support of NIH (GM-61951) and Daiso Inc.

### **References and notes**

- 1. Simpkins, N. S. Sulphones in Organic Synthesis; Pergamon: Oxford, 1993.
- For selective examples of asymmetric conjugate additions to α,β-unsaturated sulfones catalyzed by organic catalysts, see: (a) Pinheiro, S.; Guingant, A.; Desmaële, D.; d'Angelo, J. *Tetrahedron: Asymmetry* **1992**, 3, 1003; (b) d'Angelo, J.; Revial, G. *Tetrahedron: Asymmetry* **1991**, 2, 199; (c) Enders, D.; Müller, S. F.; Raabe, G.; Runsink, J. *Eur. J. Org. Chem.* **2000**, 879; (d) Mauleón, P.; Carretero, J. C. *Org. Lett.* **2004**, 6, 3195; (e) Mauleón, P.; Carretero, J. C. *Chem. Commun.* **2005**, 4961; (f) Mosse, S.; Alexakis, A. *Org. Lett.* **2005**, 7, 4361; (g) Sulzer-Mosse, S.; Tissot, M.;

Alexakis, A. Org. Lett. **2007**, *9*, 3749; (h) Zhu, Q.; Lu, Y. Org. Lett. **2008**, *10*, 4803; For examples involving chiral metallic catalysts, see: (i) Llamas, T.; Arrayas, R. G.; Carretero, J. C. Angew. Chem., Int. Ed. **2007**, *46*, 3329; (j) Desrosiers, J.-N.; Charette, A. B. Angew. Chem., Int. Ed. **2007**, *46*, 5955; (k) Mauleón, P.; Alonso, I.; Rivero, M. R.; Carretero, J. C. J. Org. Chem. **2007**, *72*, 9924; (l) Desrosiers, J.-N.; Bechara, W. S.; Charette, A. B. Org. Lett. **2008**, *10*, 2315; (m) Bos, P. H.; Minnaard, A. J.; Feringa, B. L. Org. Lett. **2008**, *10*, 4219.

- For a preliminary communication of our studies of 5c-catalyzed conjugate additions to vinylsulfones, see: (a) Li, H.; Song, J.; Liu, X.; Deng, L. J. Am. Chem. Soc. 2005, 127, 8948.
- For select reviews of catalytic enantioselective construction of all carbon quaternary chiral centers, see: (a) Douglas, C. J.; Overman, L. E. Proc. Natl. Acad. Sci. U.S.A. 2004, 101, 5363; (b) Christoffers, J.; Baro, A. Angew. Chem., Int. Ed. 2003, 42, 1688; (c) Christoffers, J. Chem.—Eur. J. 2003, 9, 4862; (d) Christoffers, J.; Mann, A. Angew. Chem., Int. Ed. 2001, 40, 4591; (e) Corey, E. J.; Guzman-Perez, A. Angew. Chem., Int. Ed. 1998, 37, 388.
- (a) Li, H.; Wang, Y.; Tang, L.; Deng, L. J. Am. Chem. Soc. 2004, 126, 9906; (b) Li, H.; Wang, Y.; Tang, L.; Wu, F.; Liu, X.; Guo, C.; Foxman, B. M.; Deng, L. Angew. Chem., Int. Ed. 2005, 44, 105; For other select examples of 5c-catalyzed other asymmetric reactions, see: (c) Wu, F.; Hong, R.; Khan, J.; Liu, X.; Deng, L. Angew. Chem., Int. Ed. 2006, 45, 4301; (d) Wang, Y.; Liu, X.; Deng, L. J. Am. Chem. Soc. 2006, 128, 3928; (e) Li, H.; Wang, B.; Deng, L. J. Am. Chem. Soc. 2006, 128, 732; (f) Li, H.; Wang, Y.-Q.; Deng, L. Org. Lett. 2006, 8, 4063; (g) Wang, Y.; Li, H.; Wang, Y.-Q.; Liu, Y.; Foxman, B. M.; Deng, L. J. Am. Chem. Soc. 2007, 129, 6364.
- For a minireview of asymmetric catalysis with C6'–OH cinchona alkaloids (cupreines and cupreidines), see: Marcelli, T.; van Maarseveen, J.; Hiemstra, H. Angew. Chem., Int. Ed. 2006, 45, 7496; For a review of asymmetric catalysis by hydrogen-bond donors, see: Doyle, A.; Jacobsen, E. N. Chem. Rev. 2007, 107, 5713.
- 7. For the synthesis of **4b**, see: Alonso, D. A.; Nájera, C.; Varea, M. *Synthesis* **2003**, 277.
- For the synthesis of 17b-d, see: Mahadevan, A.; Fuchs, P. L. Tetrahedron Lett. 1994, 33, 6025.