


Absolute Stereochemistry of  
Amphidinolide QYohei Takahashi,<sup>†</sup> Takaaki Kubota,<sup>†</sup> Eri Fukushi,<sup>§</sup> Jun Kawabata,<sup>§</sup> and  
Jun'ichi Kobayashi<sup>\*,†</sup>Graduate School of Pharmaceutical Sciences, Hokkaido University,  
Sapporo 060-0812, Japan, and Graduate School of Agriculture,  
Hokkaido University, Sapporo 060-0859, Japan

jkobay@pharm.hokudai.ac.jp

Received June 10, 2008

## ABSTRACT




amphidinolide Q (1)

The absolute configurations at five chiral centers in amphidinolide Q (1), a cytotoxic 12-membered macrolide isolated from a marine dinoflagellate *Amphidinium* sp., were elucidated to be 4*R*, 7*R*, 9*S*, 11*R*, and 13*R* on the basis of NMR analyses and a modified Mosher's method.

Amphidinolides are a series of cytotoxic macrolides possessing unique structural features isolated from laboratory cultured marine dinoflagellates *Amphidinium* sp.<sup>1</sup> Amphidinolide Q (1) isolated from a dinoflagellate *Amphidinium* sp. (Y-5 strain) is a cytotoxic 12-membered macrolide having C1 branches at vicinal carbons (C-13 and C-14) and an  $\alpha,\beta$ -unsaturated ester moiety.<sup>2</sup> The gross structure of 1 has been elucidated by 2D NMR, whereas the stereochemistry remains unsolved due to lack of sample.<sup>2</sup> Quite recently, we could reisolate amphidinolide Q (1) from the same strain so that this sample was used to elucidate the stereochemistry of 1. Here, we describe the elucidation of the absolute configurations at five chiral centers in 1 on the basis of the NMR analysis and a modified Mosher's method.

The relative stereochemistry for amphidinolide Q (1) was elucidated on the basis of the *J*-based configuration analysis

(JBCA) method<sup>3</sup> and NOESY correlations. The geminal, vicinal, and long-range *J*(C,H) coupling constants were obtained by the resolution-enhanced <sup>1</sup>H spectrum, the HET-LOC,<sup>4</sup> and *J*-IMPEACH-MBC<sup>5</sup> spectra in C<sub>6</sub>D<sub>6</sub>. A small value for <sup>3</sup>*J*(H-4/H-5a) and a large value for <sup>2</sup>*J*(H-5b/C-4) indicated that H-4/H-5a and 4-OH/H-5b were both *gauche* (Figure 1). A middle value for <sup>3</sup>*J*(H-4/H-5b) suggested that



**Figure 1.** Rotation models for the C-4/C-5 bond of amphidinolide Q (1).

the C-4/C-5 bond existed as two conformers, while NOESY correlations for H-4/H-5a, H-4/H-5b, and H-5a/H<sub>3</sub>-17 in C<sub>6</sub>D<sub>6</sub> indicated that the C-4/C-5 bond existed mainly as a


<sup>†</sup> Graduate School of Pharmaceutical Sciences.

<sup>§</sup> Graduate School of Agriculture.

(1) (a) Kobayashi, J.; Kubota, T. *J. Nat. Prod.* **2007**, *70*, 451–460. (b) Kobayashi, J. *J. Antibiot.* **2008**, *61*, 271–284. (c) Takahashi, Y.; Kubota, T.; Kobayashi, J. *J. Antibiot.* **2007**, *60*, 376–379. (d) Oguchi, K.; Tsuda, M.; Iwamoto, R.; Okamoto, Y.; Kobayashi, J.; Fukushi, E.; Kawabata, J.; Ozawa, T.; Masuda, A.; Kitaya, Y.; Omasa, K. *J. Org. Chem.* **2008**, *73*, 1567–1570.

(2) Kobayashi, J.; Takahashi, M.; Ishibashi, M. *Tetrahedron Lett.* **1996**, *37*, 1449–1450.


conformer with *gauche* relationships for C-3/H-5a and H-4/H-5b. The relative stereochemistry of the C-7 to C-11 moiety in **1** was deduced from analyses of the HETLOC, *J*-IMPEACH-MBC, and NOESY spectra in C<sub>6</sub>D<sub>6</sub>. The middle values for  $^3J(\text{H-7/H-8b})$  and  $^3J(\text{H-9/H-10a})$  indicated that C-7/C-8 and C-9/C-10 bonds adapted two conformers, while NOESY cross-peaks of H-7/H-9, H-8a/H-10a, H-9/H-11, and H-10b/H<sub>3</sub>-19 as well as small *J*-values for H-7/H-8a, H-8b/C-6, H-9/H-10b, and H-10a/C-19 allowed us to assign the major conformer for C-7/C-9 and C-9/C-10 bonds as shown in Figure 2a and 2c, respectively. NOESY correlations for




**Figure 2.** Rotation models for (a) C-7/C-8, (b) C-8/C-9, (c) C-9/C-10, and (d) C-10/C-11 bonds of amphidinolide Q (**1**).

H-7/H-9, H-8a/H-10a, H-9/H-11, and H-10b/H-11 and values for  $^3J(\text{H-9/H-8a})$ ,  $^3J(\text{H-9/H-8b})$ ,  $^3J(\text{H-8a/C-19})$ ,  $^3J(\text{H-11/H-10a})$ ,  $^3J(\text{H-11/H-10b})$ , and  $^2J(\text{H-10b/C-11})$  revealed that C-8/C-9 and C-10/C-11 bonds existed as a single conformer as shown in Figure 2b and 2d, respectively.

Relative configurations at C-4 and C-7 were deduced from NOESY correlations for H-2/H-8a, H<sub>3</sub>-17/H-5a, H-4/H<sub>2</sub>-5, and H-5a/H-7 for **1** as shown in Figure 3.




The molecular mechanics calculations of **1** were carried out by using Sybyl 6.5 (MMFF94 force-field).<sup>6</sup> One of the most stable conformers explained well the NOESY correlations observed for **1** (Figure 5). Thus, the relative stereochemistry of amphidinolide Q (**1**) was elucidated as shown.



**Figure 5.** Selected NOESY correlations and a stable conformer for macrocyclic ring of amphidinolide Q (**1**) obtained from the molecular mechanics calculations (hydrogen atoms of methyl groups were omitted).


To elucidate the absolute configurations of amphidinolide Q (**1**), a modified Mosher's method<sup>7</sup> was applied as follows. Treatment of **1** with (*R*)-(-)- and (*S*)-(+)-2-methoxy-2-trifluoro-2-phenylacetyl chloride (MTPACl) gave the (*S*)- and (*R*)-MTPA esters (**2a** and **2b**, respectively) of **1**.  $\Delta\delta$  values ( $\Delta\delta = \delta_S - \delta_R$ ) obtained from <sup>1</sup>H NMR data of **2a** and **2b** are shown in Figure 6. The  $\Delta\delta$  values for H<sub>2</sub>-5 and H<sub>3</sub>-18 are positive, while negative  $\Delta\delta$  values are observed for H-2, H<sub>2</sub>-10, H-13, H<sub>3</sub>-16, H<sub>3</sub>-17, H<sub>3</sub>-19, H<sub>3</sub>-20, and H<sub>2</sub>-21.<sup>8</sup> These results indicated that the absolute configuration at C-4 was *R*.

The relative stereochemistry of **1** was elucidated as described above, while the absolute configuration at C-11 was elucidated directly by application of a modified Mosher method to the linear methyl ester derivatives of amphidinolide Q (**1**). Treatment of **1** with K<sub>2</sub>CO<sub>3</sub> in MeOH yielded a



**Figure 6.**  $\Delta\delta$  values [ $\Delta\delta$  (in ppm) =  $\delta_S - \delta_R$ ] obtained for the (*S*)- and (*R*)-MTPA esters (**2a** and **2b**, respectively) of amphidinolide Q (**1**).

mixture of two linear methyl esters generated by dehydration of C-4 and C-5 and epimerization of C-7. The mixture was treated with (*R*)- and (*S*)-MTPACl, and the resulting bis- (*S*)- and (*R*)-MTPA esters were obtained by HPLC purification.  $\Delta\delta$  values for **3a** and **3b** suggested the absolute configuration at C-11 to be *R* (Figure 7). Thus, the absolute configurations at the five chiral centers in **1** were concluded to be 4*R*, 7*R*, 9*S*, 11*R*, and 13*R*, respectively.



**Figure 7.**  $\Delta\delta$  values [ $\Delta\delta$  (in ppm) =  $\delta_S - \delta_R$ ] obtained for the (*S*)- and (*R*)-MTPA esters (**3a** and **3b**, respectively) of the linear methyl ester derivatives of amphidinolide Q (**1**).

**Acknowledgment.** The authors thank Ms. S. Oka, Center for Instrumental Analysis, Hokkaido University, for measurements of ESIMS. This work was partly supported by grants from a Takeda Science Foundation, a Mochida Memorial Foundation for Medical and Pharmaceutical Research, and a Grant-in-Aid for Scientific Research from the Ministry of Education, Culture, Sports, Science and Technology of Japan.

**Supporting Information Available:** Experimental procedures and one- and two-dimensional NMR spectra for **1**, **2a**, **2b**, **3a**, and **3b**. This material is available free of charge via the Internet at <http://pubs.acs.org>.

OL8013123

(6) Halgren, T. A. *J. Am. Chem. Soc.* **1992**, *114*, 7827–7643.

(7) Ohtani, I.; Kusumi, T.; Kashman, Y.; Kakisawa, H. *J. Am. Chem. Soc.* **1991**, *113*, 4092–4095.

(8) A positive  $\Delta\delta$  value for H-11 seems to be due to the anisotropic effect of the ester carbonyl group at C-1.