A Convenient Preparation of 3-Substituted Furans: Synthesis of Perillene and Dendrolasin

D. K. Barma,[†] Abhijit Kundu,[†] Rachid Baati,[‡] Charles Mioskowski,^{*,‡,§} and J. R. Falck^{*,†}

Department of Biochemistry, University of Texas Southwestern Medical Center, Dallas, Texas 75390-9038, and Université Louis Pasteur, Faculté de Pharmacie, Laboratoire de Synthèse Bio-Organique, associe au CNRS 67401 Illkirch, France

j.falck@utsouthwestern.edu

Received February 12, 2002

ORGANIC LETTERS 2002 Vol. 4, No. 8 1387–1389



A variety of 3-substituted furans, including the natural products perillene and dendrolasin, are obtained in good yield via reductive annulation of 1,1,1-trichloroethyl propargyl ethers using catalytic Cr(II) regenerated by Mn/TMSCI.

Furans are common subunits in pharmaceuticals,¹ natural products,² fragrances,³ and flavors.⁴ They have also found utility as synthetic intermediates or synthons for numerous functional groups, inter alia, carboxylic acids, β -keto-esters, and aromatics.⁵ While electrophilic aromatic substitution or metalation⁶ of furans permit ready functionalization at C(2)/(5), these methods are not generally applicable to the preparation of C(3)-substituted furans. The latter are typically obtained by elaboration of an existing C(3)-substituent⁷ or accessed from acyclic precursors via multistep sequences.⁸ In continuation of our exploration of the chemistry of

- § E-mail: mioskow@aspirine.u-strasbg.fr.
- (1) (a) Bock, I.; Bornowski, H.; Ranft, A.; Theis, H. *Tetrahedron.* **1990**, *46*, 1199–1210. (b) Mortensen, D. S.; Rodriguez, A. L.; Carlson, K. E.; Sun, J.; Katzenellenbogen, B. S.; Katzenellenbogen, J. A. *J. Med. Chem.* **2001**, *44*, 3838–3848.
 - (2) Murray, R. D. H. Aromat. Heteroaromat. Chem. 1978, 6, 258-281.
- (3) Levisalles, J. Perfumery Essent. Oil Record 1958, 49, 627–635.
- (4) Naim, M.; Zuker, I.; Zehavi, U.; Rouseff, R. L. J. Agric. Food Chem. 1993, 41, 1359–1361.
- (5) Meyers, A. I. *Heterocycles in Organic Synthesis*; Wiley-Interscience: New York, 1974.
- (6) (a) Kojima, Y.; Wakita, S.; Kato, N. *Tetrahedron Lett.* **1979**, 4577–4580.
 (b) Tanis, S. P.; Herrinton, P. M. J. Org. Chem. **1983**, 48, 4572–4580.
- (7) Mandai, T.; Takeshita, M.; Mori, K.; Kawada, M.; Otera, J. *Chem. Lett.* **1983**, 1909–1910.

10.1021/ol025708s CCC: \$22.00 © 2002 American Chemical Society Published on Web 03/19/2002

organochromium reagents,⁹ we describe herein a convenient synthesis of C(3)-substituted furan by reductive annulation of 1,1,1-trichloroethyl propargyl ethers¹⁰ mediated by catalytic Cr(II) coupled to a Mn/TMSCl regeneration system (eq 1).¹¹

To study the efficacy of the annulation, a representative panel of 1,1,1-trichloroalkynes was prepared by alkylation of 2,2,2-trichloroethanol with various propargyl bromides at

[†] University of Texas Southwestern.

[‡] Université Louis Pasteur.

⁽⁸⁾ Review of furan syntheses: Hou, X. L.; Cheung, H. Y.; Hon, T. Y.; Kwan, P. L.; Lo, T. H.; Tong, S. Y.; Wong, H. N. C. *Tetrahedron* **1998**, *54*, 1955–2020.

⁽⁹⁾ Recent CrCl₂-based methodology: (a) Falck, J. R.; Barma, D. K.; Venkataraman, S. K.; Baati, R.; Mioskowski, C. *Tetrahedron Lett.* **2002**, *43*, 963–966. (b) Baati, R.; Barma, D. K.; Krishna, U. M.; Mioskowski, C.; Falck, J. R. *Tetrahedron Lett.* **2002**, *43*, 959–961. (c) Baati, R.; Barma, D. K.; Falck, J. R.; Mioskowski, C. J. Am. Chem. Soc. **2001**, *123*, 9196– 9197. (d) Barma, D. K.; Baati, R.; Valleix, A.; Mioskowski, C.; Falck, J. R. Org. Lett. **2001**, *3*, 4237–4238. (e) Falck, J. R.; Barma, D. K.; Baati, R.; Mioskowski, C. Angew. Chem., Int. Ed. **2001**, *40*, 1281–1283. (f) Baati, R.; Valleix, A.; Mioskowski, C.; Barma, D. K.; Falck, J. R. Org. Lett. **2000**, *2*, 485–487. (g) Falck, J. R.; Barma, D. K.; Mioskowski, C.; Schlama, T. Tetrahedron Lett. **1999**, *40*, 2091–2094.

room temperature in the presence of NaOH/*n*-Bu₄NI in CH₂Cl₂.¹² Stirring the structurally simplest of these, **1**, with stoichiometric (4–6 equiv) CrCl₂ proved disappointing and never generated more than a 30% yield of furan **2**, regardless of solvent or temperature, with the remaining material balance being principally Z-chloroenol ether **30**. In stark contrast, **2** was obtained¹³ in good yield (Table 1, entry 1)

 Table 1. Synthesis of 3-Substituted Furans



by warming a THF solution of **1** with a catalytic amount of $CrCl_2$, or more conveniently $CrCl_3$, coupled to a Mn powder/ TMSCl regeneration system as prescribed by Fürstner.¹⁴ Other solvents, e.g., HMPA, DMF, CH_2Cl_2 , Et_2O , *p*-dioxane, and EtOAc, were less satisfactory. Notably, the corresponding tribromoethyl ether led to a complex product mixture containing little, if any, **2**.

Likewise, the naturally occurring¹⁵ furanyl terpenoids perillene (4) and dendrolasin (6) were synthesized in a single step from isoprenoids 3 and 5 (entries 2 and 3, respectively)

using catalytic CrCl₂. No olefin isomerization or reduction was observed during the transformation of styrene 7 to 8(entry 4). Phenylacetylene 9, on the other hand, was consumed under the standard catalytic conditions but afforded a poor yield ($\sim 20\%$) of **10**. However, the reaction efficiency was significantly improved using 6 equiv of CrCl₂ at 60 °C for 12 h and furnished 10 in 70% yield (entry 5). The presence of heteroatoms or leaving groups adjacent to the distal side of the acetylene resulted in low yields, for example, 11 to 12 (entry 6), although β -functionality was well tolerated (13 to 14, entry 7). The reaction conditions proved compatible with most other common functional groups as well, inter alia, methyl ester 15, benzyl ether 17, acetate 19, silvloxy 21, and free alcohol 23 which gave rise to 3-substituted furans 16 (entry 8), 18 (entry 9), 20 (entry 10), 22 (entry 11), and 24 (entry 12), respectively. A mechanistic interpretation of the preceding observations is summarized in Scheme 1. One electron transfer¹⁶ from Cr(II)



to the trichloroalkyne substrate with concomitant loss of chloride spawns the key radical intermediate **25** which undergoes cyclization¹⁷ to give vinylchromium **26** under catalytic Cr(II) conditions (pathway a). Further transmetalation and aromatization lead to furan **27**. The latter was inferred by quenching with D₂O, which resulted in **28**, incorporating one atom of deuterium. Alternatively, in the presence of high concentrations of Cr(II), i.e., stoichiometric conditions, reduction of **25** to **29** is competitive (pathway b) and Z-chloroenol ether **30** is formed as previously described.^{9a} Control experiments demonstrated that both Mn and TMSCI

(13) General procedure: 1,1,1-Trichloroethyl propargyl ether (1 mmol) in THF (2 mL) is added to a stirring, room-temperature suspension of anhydrous $CrCl_3$ (15 mol %), Mn powder (4 mmol), and freshly distilled TMSCl (4 mmol) in THF (8 mL) under argon. After complete addition, the reaction mixture was heated at 60 °C. After 12–15 h, the reaction mixture was cooled to ambient, quenched with an equal volume of water, and extracted thrice with ether. The combined ethereal extracts were evaporated in vacuo and the residue was purified by SiO₂ chromatography to give 3-substituted furans in the indicated yields (Table 1).

(14) Fürstner, A.; Shi, N. J. Am. Chem. Soc. 1996, 118, 12349-12357.
 (15) Bernardi, R.; Cardani, C.; Selva, D.; Baggini, A.; Pavan, M. Tetrahedron Lett. 1967, 8, 3893-3896.

(16) Okazoe, T.; Takai, K.; Utimoto, K. J. Am. Chem. Soc. 1987, 109, 951.

(17) Review of heterocyclic syntheses via radical cyclization: Bowman, W. R.; Cloonan, M. O.; Krintel, S. L. *J. Chem. Soc., Perkin Trans. 1* **2001**, 2885–29.

⁽¹⁰⁾ Multistep synthesis of 3-substituted furans using 1,1,1-trichloroethyl allyl ethers: Ram, R. N.; Charles, I. *Chem. Commun.* **1999**, 2267–2268.

⁽¹¹⁾ CrCl₂-based synthesis of 2,5-disubstituted furans: Takai, K.; Morita, R.; Sakamoto, S. *Synlett* **2001**, 1614–1616.

⁽¹²⁾ Morimoto, T.; Sekiya, M. Synthesis 1981, 308-310.

are need for the catalytic procedure to succeed. However, the role of the TMSCl is obscure in this instance, but is not a source of HCl as proposed¹¹ in other systems since exogenously added acid was detrimental to the yield.

In summary, we report a convenient, high-yield synthesis of 3-substituted furans from readily available acyclic precursors utilizing, in most cases, catalytic CrCl₂. Extensions of this methodology to the preparation of di- and trisubstituted furans are under active investigation, and the results will be the subject of future publications.

Acknowledgment. The authors thank Dr. E. R. Fogel for helpful discussions. Financial support was provided by the Robert A. Welch Foundation, the NIH (GM 31278, DK38226), and the Instituts de Recherche Pierre Fabre, CNRS (to R.B.).

Supporting Information Available: Spectral data for all new compounds. This material is available free of charge via the Internet at http://pubs.asc.org.

OL025708S