This article was downloaded by: [University of California, San Francisco] On: 29 December 2014, At: 10:43 Publisher: Taylor & Francis Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Phosphorus, Sulfur, and Silicon and the Related Elements

Publication details, including instructions for authors and subscription information: http://www.tandfonline.com/loi/gpss20

Synthesis and Biological Activities of O, O-Dialkyldithiophosphoryl-S-acetoxy Triazolo Compounds

Wen-Bin Chen^a & Gui-Yu Jin^a ^a Nankai University, Tianjin, China Published online: 27 Oct 2010.

To cite this article: Wen-Bin Chen & Gui-Yu Jin (2003) Synthesis and Biological Activities of O, O-Dialkyl-dithiophosphoryl-S-acetoxy Triazolo Compounds, Phosphorus, Sulfur, and Silicon and the Related Elements, 178:2, 311-317, DOI: <u>10.1080/10426500307939</u>

To link to this article: http://dx.doi.org/10.1080/10426500307939

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at http://www.tandfonline.com/page/terms-and-conditions



SYNTHESIS AND BIOLOGICAL ACTIVITIES OF O, O-DIALKYL-DITHIOPHOSPHORYL-S-ACETOXY TRIAZOLO COMPOUNDS

Wen-Bin Chen and Gui-Yu Jin Nankai University, Tianjin, China

(Received May 10, 2002; accepted June 9, 2002)

In this article, we introduce organic phosphorus moiety into triazole ring to prepare O-dialkyl-dithiophosphoryl-S-acetoxy triazolo compounds and characterize their structures by elementary analysis and ¹HNMR and IR spectral data. From the results of biological activity screening, we found that the O-dialkyl-dithiophosphoryl-S-acetoxy triazolo compounds show lower fungicidal activities and their structures need to be optimized for the better.

Keywords: Biological activities; organic phosphorus; synthesis; triazole

INTRODUCTION

Organic phosphorus compounds are biologically versatile as they are pesticides, fungicides, and herbicides,¹⁻⁴ and many compounds have been commercialized. At the same time, triazole compounds also possess pesticidal, herbicidal, and fungicidal activities and plant growth regulation activities.⁵⁻¹¹ The introduction of organic phosphorus in triazole compounds not only reduces the toxicity of organic phosphorus compounds on mammalian, but also expands the application scope of triazole compounds. However, triazole compounds containing organic phosphorus are seldom reported.^{12,13} To study the biological activities of all kinds of triazole compounds to prepare O-dialkyl-dithiophosphoryl-S-acetoxy triazolo compounds. The syntheses of these compounds are outlined in Schemes 1, 2, and 3.

The authors wish to thank the National Natural Science Foundation of China (No. 29832050) for support of this project.

Address correspondence to Gui-Yu Jin, Institute of Elemento-Organic Chemistry, Nankai University, Tianjin 300071, People's Republic of China. E-mail: wbchen@ eyou.com



SCHEME 3

RESULTS AND DISCUSSION

Preparation of O-Dialkyl-dithiophosphoryl-S-acetoxy Triazolo Compounds (5)

During the course of our experiment, we found that the intermediate **3** was unstable and decomposed for some time to a unknown solid. Thus, compounds **3** must be freshly prepared before using.

The base is necessary to this reaction. Results show that no product was obtained when the base was absent. At the same time, we tested such bases as triethylamine, pyridine, and potassium carbonate. Here, the reactant became very complex and no obvious product was detected using potassium carbonate as a base. When we used pyridine as a base, the same result was observed. Only using triethylamine, were we able to obtain the desired result. On the other hand, the reaction temperature is an important factor in the reaction. A higher temperature lead to dark color and complicated unknown compounds; at room temperature or a lower temperature, desired results were observed. In the end, we found that the intermediate **3b** is more reactive than **3a**, as it reacts at lower temperature.

	$\operatorname{IR}(v/\operatorname{cm}^{-1},\operatorname{film})$							
Compd.	С=0	C=N (triazole)	C—H (triazole)	P=S	Р-О-С			
5a 5b 5c 5d 5e 5f	$\begin{array}{c} 1727.7,1751.3\\ 1726.3,1751.8\\ 1723.5,1754.4\\ 1723.8,1758.4\\ 1704.6,1758.1\\ 1704.7,1756.1 \end{array}$	$\begin{array}{c} 1505.9,1476.9\\ 1504.9,1475.3\\ 1503.0,1467.3\\ 1503.4,1478.9\\ 1507.0,1440.4\\ 1506.8,1459.3 \end{array}$	3102.5 3109.0 3108.0 3112.0 3108.5 3103.5	$\begin{array}{c} 654.1 \\ 668.3 \\ 658.5 \\ 653.1 \\ 654.2 \\ 656.8 \end{array}$	$\begin{array}{c} 1009.4,1173.7,771.2\\ 1000.4,1180.1,793.0\\ 993.4,1180.5,757.6\\ 994.7,1182.5,749.4\\ 1006.1,1153.9,783.7\\ 977.4,1154.2,747.6\end{array}$			
5g	1707.8, 1752.8	1506.9, 1461.8	3100.0	649.2	995.6, 1154.6, 771.3			

TABLE I The IR Data of Compounds 5

Characterizations of O-Dialkyl-dithiophosphoryl-S-acetoxy Triazolo Compounds

The IR spectrum of the O-dialkyl-dithiophosphoryl-S-acetoxy triazolo compounds exhibited two sharp bands around $1700 \sim 1750 \text{ cm}^{-1}$ due to the two C=O functions, whereas the bands around 650 cm⁻¹ and 1000 cm⁻¹ were attributed to P=S and P-O-C groups respectively. The other data was outlined in Table I.

With respect to the ¹HNMR spectra, there is much difference because of the different substituents R and R'. When R is *i*-Pr, the two *i*-propoxy bonded to phosphorus atom are not equal to appear two doublet peaks. But the difference is so small that triplet peaks were observed. While R is Me, Et, and *n*-Pr, the two alkyoxy groups are equal and the same chemical shifts were observed. At the same time, different substituent R' also affect the chemical shift of the methenyl group bonded to triazole ring. When R' is *i*-butyl, it appeared as singlet peak at δ 7.3, while R' is 4-FC₆H₄, the singlet peak was observed at δ 8.5–8.7. The detailed data was shown in Table II.

Biological Activities Results

We selected some compounds **5** to test their fungicidal and insecticidal activities and found that they possess some fungicidal but no insectididal activities. Some results of these compounds are outlined in Table III.

EXPERIMENTAL

¹HNMR spectra were recorded in CDCl₃ as solvent on an AC-P200 instrument using TMS as an internal standard. IR spectra were measured

TABLE II The IR Data of Compounds 5

Compd.	¹ HNMR δ (ppm), J _{P-H} (Hz) (CDCl ₃ /TMS)
5a	1.15 (s, 9H, $3 \times CH_3$), 3.73 (d, 2H, CH ₂ S, J _{P-H} =15.33), 3.78 (d, 6H, $2 \times CH_3$ O, J _P = -15.22), 7.36 (c, 1H, CH), 8.05 (c, 1H, Tr, H), 8.51 (c, 1H, Tr, H)
5b	1.13 (s, 9H, $3 \times CH_3$), 1.29 (t, 6H, $2 \times CH_3CH_2O$), 3.73 (d, 2H, CH ₂ S, J _{P-H} =17.77), 4.15 (m, 4H, $2 \times CH_3CH_2O$), J _{P-H} =15.22), 7.31 (s, 1H, CH), 8.00 (s, 1H, Tr-H), 8.36 (s, 1H, Tr-H)
5c	0.96 (t, 6H, $2 \times \underline{CH_3CH_2CH_2O}$), 1.18 (ts, 9H, $3 \times CH_3$), 1.72 (m, 4H, $2 \times \underline{CH_3CH_2CH_2O}$), 3.79 (d, 2H, $\underline{CH_2S}$, $\underline{J_{P-H}} = 17.73$), 3.98 (m, 4H, $2 \times \underline{CH_3CH_2CH_2O}$), 7.38 (s, 1H, CH), 8.08 (s, 1H, Tr-H), 8.55 (s, 1H, Tr-H)
5d	1.19 (s, 9H, $3 \times CH_3$), 1.34 (t, 12H, $2 \times (\underline{CH_3})_2CH$), 3.82 (d, 2H, CH_2S , $J_{P-H} = 17.39$), 4.81 (m, 2H, $2 \times (CH_3)_2\underline{CH}$), 7.39 (s, 1H, CH), 8.09 (s, 1H, Tr-H). 8.52 (s, 1H, Tr-H)
5e	1.33 (t, 6H, 2 × \underline{CH}_3CH_2O), 3.80 (d, 2H, CH_2S , $J_{P-H} = 18.34$), 4.13 (m, 4H, 2 × $CH_3\underline{CH}_2O$), 7.12~7.22, 7.91~7.98 (m, 4H, C_6H_4), 7.82 (s, 1H, Tr-H), 8.05 (s, 1H, Tr-H), 8.56 (s, 1H, CH)
5f	$\begin{array}{l} \textbf{0.95 (t, 6H, 2 \times \underline{CH_3}CH_2CH_2O), 1.71 (m, 4H, 2 \times CH_3CH_2 CH_2O), 3.81} \\ \textbf{(d, 2H, CH_2S, J_{P-H} = 17.87), 4.06 (m, 4H, 2 \times CH_3CH_2\underline{CH_2}O), 7.13 \sim 7.22, } \\ \textbf{7.91} \sim 8.02 (m, 4H, C_6H_4), 7.84 (s, 1H, Tr-H), 8.07 (s, 1H, Tr-H), 8.71 \\ \textbf{(s, 1H, CH)} \end{array}$
5g	$ \begin{array}{l} 1.31 \; (t, 12H, 2 \times (\underline{CH_3})_2 CHO), 3.83 \; (d, 2H, CH_2 S, J_{P\!-\!H} = \! 17.54), 4.81 \; (m, 2H, \\ 2 \times (CH_3)_{\underline{2}} \underline{CHO}), 7.12 \sim \! 7.23, 7.93 \sim \! 8.01 \; (m, 4H, C_6 H_4), 7.85 \; (s, 1H, Tr\!-\!H), \\ 8.07 \; (s, 1H, Tr\!-\!H), 8.68 \; (s, 1H, CH) \end{array} $

on a Nicolet 5DX IR spectrometer. Elemental analyses were conducted on an MF-3 automatic analyzer. Melting points were determined on an MP-500 melting point apparatus. Refractive indices were obtained by use of Abbe refractometer. All temperatures and pressures are uncorrected.

Compd.	Fungicidal activities (%)									
	Ir	In vitro (50 ppm)								
	S. sclerotiorum	P. recibduta	B. cinerea	A. solani	P. asparagi	P. piricola	G. zeae	C. arachidicola		
5a	5	0	0	4.8	0	7.7	2.8	0		
5b	0	0	0	9.5	10	11.5	2.8	0		
5c	10	10	0	0	0	3.8	5.6	0		
5d	0	10	0	0	0	11.5	2.8	6.7		
5e	15	0	49.6	9.5	0	7.7	5.6	6.7		

TABLE III The Fungicidal Activities of Compounds 5

Compd.	R	m.p. (°C)	Yield (%)	Compd.	R	n_{D}^{25}	Yield (%)
1a	Me	$190{\sim}2$	$52.3 \\ 48.7 \\ 44.1 \\ 40.2$	2a	Me	1.5070	98.0
1b	Et	$213{\sim}4$		2b	Et	1.5078	96.0
1c	<i>n</i> -Pr	$168{\sim}70$		2c	<i>n</i> -Pr	1.5056	86.9
1d	<i>i</i> -Pr	$204{\sim}6$		2d	<i>i</i> -Pr	1.5046	97.1

TABLE IV Some Data of Compounds of 1 and 2

Preparations of Intermediates

Preparation of O,O-Dialkylphosphinothioylthioacetic Acetic Acid 2^{14,15}

According to the literature procedure, potassium O,O-diethylphosphorodithioate¹⁴ was treated with chloracetic acid in acetone to give the liquid that was purified by silicon gel column chromatography using petroleum ether as eluent to give the compound **2b**, yield 96.0%, n_D^{25} 1.5078, ¹HNMR (δ /ppm, CDCl₃): 134 (t, 6H, 2 × <u>CH₃CH₂O</u>), 3.67 (d, 2H, CH₂S J_{P-H} =17.54 Hz), 4.11–4.23 (m, 4H, 2 × CH₃<u>CH₂O</u>), 9.98(brs, 1H, COOH).

Similarly, other compounds 2 were prepared from the corresponding potassium, and some data of compounds 1 and 2 were shown in Table IV.

Preparations of Bromo Ketone 4¹⁶

Triazolo ketone **3** was treated with bromine in acetic acid and sodium acetate anhydride to give the crude products **4a** and **4b** which not purified for the next step, yield 84.4% and 95.4%.

Preparations of O-Dialkyl-dithiophosphoryl-S-acetoxy Triazolo Compounds 5

To the mixture of O, O-dimethylphosphinothioylthioacetic acetic acid **2a** 1.08 g (5.0 mmol), and 30 mL of dry acetone, was added 0.56 g (5.5 mmol) Et₃N by stirring at room temperature. After addition, the mixture was cooled to $0 \sim 5^{\circ}$ C with ice-water bath, 1.23 g (5.0 mmol) 3, 3-dimethyl-1-bromo-1-(1H-1, 2, 4-triazolo-1-yl)-2-butanone **3a** in 5 mL of dry acetone was added to the reactant, then triethylamine hydrobromide appeared. After the addition was completed, the ice-water bath was removed and the mixture was stirred for 4 h at room temperature. The solvent was evaporated off under reduced pressure, and the residue was diluted with 30 mL of CH₂Cl₂ and washed with water and dried over MgSO₄. After removal of solvent, the residue was separated by silicon gel chromatography using ethyl acetate/petroleum ether as eluent to give the viscous liquid, 1.54 g, yield 80.8%. The other compounds **5b–5d** were prepared by the same procedure.

		R'	m.p. (°C)	Yield (%)	Elementary analyses (% calcd.)		
Compd.	R				С	Н	Ν
5a	Me	$(\mathrm{CH}_3)_3\mathrm{C}$	Viscous liquid	80.8	37.80 (37.79)	5.09 (5.25)	10.89 (11.02)
5b	Et	$(\mathrm{CH}_3)_3\mathrm{C}$	Viscous liquid	80.2	41.10 (41.08)	5.62 (5.87)	10.17 (10.27)
5 c	<i>n</i> -Pr	$(\mathrm{CH}_3)_3\mathrm{C}$	Viscous liquid	65.6	43.67 (43.94)	6.19 (6.41)	9.34 (9.61)
5d	<i>i</i> -Pr	$(\mathrm{CH}_3)_3\mathrm{C}$	$46 \sim \!\! 48$	62.5	43.98 (43.94)	6.16 (6.41)	9.33 (9.61)
5e	Et	$4\text{-}\mathrm{FC}_{6}\mathrm{H}_{4}$	Viscous liquid	77.8	43.04 (42.95)	3.99 (4.25)	8.87 (9.39)
5f	<i>n</i> -Pr	$4\text{-}\mathrm{FC}_{6}\mathrm{H}_{4}$	Viscous liquid	61.3	45.08 (45.47)	4.71 (4.84)	8.68 (8.84)
5g	<i>i-</i> Pr	$4\text{-}\mathrm{FC}_{6}\mathrm{H}_{4}$	Viscous liquid	59.5	45.64 (45.47)	4.47 (4.84)	8.40 (8.84)

TABLE V The Data of Compounds 5

Similarly, to the mixture of 1.22 g (5.0 mmol) O, O-diethylphosphinothioylthioacetic acetic acid **2b**, 0.56 g (5.5 mmol) Et₃N, and 30 mL of acetone, was added 1.42 g (5.0 mmol) α -(1H-1,2,4-triazolo-1yl)- α -bromo-*para*-fluoroacetonphenone **3b** in 5 mL of acetone at $0 \sim 5^{\circ}$ C. After the addition, the mixture was stirred for 2 h under this condition. Then the reactant was poured into 50 mL of ice-water, extracted with 3×20 mL of chloroform, washed with water twice, and dried over MgSO₄. After removal of solvent, the residue was isolated by silicon gel chromatography using ethyl acetate/petroleum ether as eluent to give viscous liquid **5e**, 1.74 g, yield 77.8%.

According to the same procedure, the title compound 5f, 5g were obtained and all compounds prepared were characterized by element analysis, HNMR and IR spectral data, their relevant data being listed in Tables I, II, and V.

REFERENCES

- [1] N. G. Smith, B. S. Watson, and F. S. Fischer, J. Agric. Food. Chem., 15, 132 (1967).
- [2] J. H. Vinopal and T. R. Fukuto, Pest Biochem. Physiol., 1, 44 (1971).
- [3] H. Leader and J. E. Casida, J. Agric. Chem., 30, 546 (1982).
- [4] S. Kimura, K. Toeda, T. Miyamoto, and L. Yamaoto, J. Pest. Sci., 9, 137 (1984).
- [5] R. K. Varma and S. C. Vyas, *Pesticides*, 3(24), 38 (1979).
- [6] H. Buchenauer, Pestic. Sci., 9, 507 (1978).
- [7] B. C. Baldwin and T. E. Wiggins, Pestic. Sci., 5(2), 150, 106 (1984).
- [8] Nihon Tokushu Noyaku Seizo K. K., Japn. Kokai. Tokyo Koho JP 5661302 (1981).
- [9] L. Klaus and R. Walf, Pestic. Sci., 19, 153 (1987).

- [10] H. Takano, Y. Oguri, T. Kato, et al., Nippon Noyaku Gakkaishi, 8(4), 575 (1983).
- [11] Y. Funaki, J. Pestic. Sci., 9, 229 (1984).
- [12] Y. N. Shi, J. X. Fang, and L. Z. Xu, Chem. J. Chinese Univ., 12(9), 1190 (1991).
- [13] Y. N. Shi, J. X. Fang, and L. Z. Xu, Chem. J. Chinese Univ., 13(8), 1084 (1992).
- [14] P. L. Carter, A. J. Lambie, and D. W. J. Lane, J. Appl. Chem., 18(9), 257 (1968).
- [15] E. I. Hoegberg and J. T. Cassaday, J. Am. Chem. Soc., 73, 557 (1951).
- [16] G. Holmwood, S. Oeckl, and J. Stetter, Ger. Offen., 2937595 (1981).