


Catalytic Regioselective Control in the Diastereoselective 1,3-Dipolar Cycloaddition Reactions of 1-(1-Alkynyl)cyclopropyl Ketones with Nitrones

Yanqing Zhang,^[a] Feng Liu,^[a] and Junliang Zhang*^[a, b]

One of the challenges in modern synthesis is the creation of distinct types of complex molecules from identical starting materials by subtly altering the choice of catalyst.^[1] Recently, much attention has been paid to 1-(1-alkynyl)cyclopropyl ketones due to their unique structures and reactivities. For example, Schmalz^[2a] and Zhang^[2b-c] have successfully developed gold-catalyzed tandem reactions of 1-(1-alkynyl)cyclopropyl ketones to efficiently construct highly substituted furans and carbobicycles. In addition, we very recently developed a Rh^I-catalyzed carbonylation reaction of 1-(1-alkynyl)cyclopropyl ketones that leads to fused 5,5-bicyclic furans.^[3]

After our study of the gold-catalyzed cycloaddition reactions of 2-(1-alkynyl)-2-alken-1-ones with nitrones and as a continuation of our efforts to design and develop novel regiodivergent reactions,^[4] we became interested in the Lewis acid catalyzed 1,3-dipolar cycloaddition reactions of 1-(1-alkynyl)cyclopropyl ketones **1** with nitrones **2**.^[5] We envisaged that these reactions might provide two different types of adduct by two regioselective cycloaddition reactions (Scheme 1): 1) tetrahydro-1,2-oxazines **3** through 1,3-dipolar (formal [3+3]) cycloaddition of nitrones to the cyclopropane moiety of **1**^[6,7] and 2) 5,7-fused heterobicyclic furo[3,4-*d*]-[1,2]oxazepines **4** through tandem double cyclization (formal [4+3] cycloaddition) of nitrone **2** with **1**.^[8,9] How to efficiently control these two types of cycloaddition reaction



Scheme 1. Proposed reaction pattern for 1-(1-alkynyl)cyclopropyl ketones with nitrones.

is an interesting, but troublesome, issue. Herein, we report our recent result that this regioselectivity can be controlled by subtle changes in the choice of catalyst. Furthermore, a kinetic resolution study and the transformation of an optically active substrate provide evidence supporting the proposed mechanism of the gold(I)-catalyzed tandem cyclization/[4+3] cycloaddition.


This hypothesis was initially tested by reacting 1-(1-alkynyl)cyclopropyl ketone **1a** with nitrone **2a** under catalysis by a series of Lewis acids. After numerous attempts, we were pleased to find that the reaction gave tetrahydro-1,2-oxazine **3aa** in 90% isolated yield with a diastereomeric ratio of 92:8 under conditions A: Sc(OTf)₃ (10 mol %), Tf = triflate), 1,10-phenanthroline (10 mol %), 4 Å molecular sieves (MS), 1,2-dichloroethane (DCE), 28–32 °C (Table 1).^[10] In contrast, the corresponding 5,7-fused bicyclic furo[3,4-*d*][1,2]oxazepine **4aa** was isolated in 96% yield, as a single diastereomer (d.r. > 20:1), under the catalysis of Ph₃PAuOTf (2 mol %) in CH₂Cl₂ at room temperature (conditions B). Interestingly, the reaction gave both formal [3+3] cycloadduct **3aa** (25% ¹H NMR yield) and [4+3] cycloadduct **4aa** (25% ¹H NMR yield) under the catalysis of AgOTf after 2 days with 50% conversion. Other Lewis

[a] Y. Zhang, F. Liu, Prof. Dr. J. Zhang
Shanghai Key Laboratory of Green Chemistry and
Chemical Processes
Department of Chemistry, East China Normal University
3663 N. Zhongshan Road, Shanghai 200062 (P.R. China)
Fax: (+86) 21-6223-5039
E-mail: jlzhang@chem.ecnu.edu.cn

[b] Prof. Dr. J. Zhang
State Key Laboratory of Organometallic Chemistry
Shanghai Institute of Organic Chemistry
Chinese Academy of Sciences
354 Fenglin Road, Shanghai 200032 (P.R. China)

Supporting information for this article is available on the WWW
under <http://dx.doi.org/10.1002/chem.200903342>.

Table 1. Testing the reaction of **1a** and **2a** with various Lewis acid catalysts.^[a]




Conditions	3aa		4aa	
	Yield [%]	d.r.	Yield [%]	d.r.
A	90	11.5:1	0	–
B	0	–	96	<20:1
C	25 ^[b]	10:1	25 ^[b]	–

[a] Conditions A: 1,10-phenanthroline (10 mol %), Sc(OTf)₃ (10 mol %), MS 4 Å, DCE, 28–32°C, 9.5 h; conditions B: Ph₃PAuOTf (2 mol %), CH₂Cl₂, RT, 10 min; conditions C: AgOTf (10 mol %), MS 4 Å, DCE, 30°C, 2 d. [b] Yield derived from the ¹H NMR spectra.

acids, such as Yb(OTf)₃, NiClO₄, and Sc(OTf)₃ without ligands were also tested and gave **3aa** in lower yields.

With the optimal reaction conditions in hand, we studied the scope of these Lewis acid catalyzed, regioselectively tunable, cycloaddition reactions (Tables 2 and 3). The gold(I)-catalyzed,^[11] formal [4+3] cycloaddition reactions of ketones **1** with nitrones **2**, which form compounds **4**, are very fast and most are completed within 10–20 min, the exceptions being nitrone **2f** (Table 2, entry 10) and ketone **1j** (Table 3, entry 18). On the other hand, the Sc(OTf)₃-catalyzed formal [3+3] cycloaddition reactions to form compounds **3** require several hours to consume all of ketone **1**. Although the diastereoselectivity of the Sc(OTf)₃-catalyzed transformation is good (d.r.=2.2–15.7:1, with most >5:1), the gold(I)-catalyzed reaction gives the cycloadducts with excellent diastereoselectivity (>20:1). It is worth noting that under both conditions A and B the cycloadditions are regiospecific, that is, no [4+3] cycloadduct was detected under conditions A and vice versa. Various functional groups, such as cyclohexenyl, cyclopropane, and esters are tolerated, which indicates that the reactions, under both sets of conditions, are highly chemoselective. Substituents on the nitron (R⁴ and R⁵) have a greater impact on the diastereoselectivity of the reaction than

Table 2. Tuning the regioselectivity of the 1,3-dipolar cycloadditions of 1-(1-alkynyl)-cyclopropyl ketone **1a** with nitrones **2**.^[a]




R ⁴	R ⁵	2	Conditions	t [h]	Yield [%] ^[b]	d.r. ^[c]
1	1-furanyl	Ph 2b	A	5.5	3ab 85	15.7:1
2		2b	B	0.3	4ab 93	>20:1
3	4-NO ₂ C ₆ H ₄	Ph 2c	A	6.5	3ac 68	4.3:1
4		2c	B	0.3	4ac 85	>20:1
5	4-MeOC ₆ H ₄	Ph 2d	A	9.5	3ad 97	13.3:1
6		2d	B	0.3	4ad 89	>20:1
7	styryl	Ph 2e	A	27	3ae 69	4.0:1
8		2e	B	0.3	4ae 92	>20:1
9	Ph	Bn 2f	A	17.5	3af 82	2.2:1
10		2f	B	6	4af 73	>20:1

[a] Conditions A: Sc(OTf)₃ (10 mol %), 1,10-phenanthroline monohydrate (10 mol %), 4 Å MS, DCE, 28–32°C; conditions B: Ph₃PAuOTf (2 mol %), DCM, RT; 1.5 equivalents of the nitron was used under conditions A and 1.1 equivalents of the nitron was used under conditions B. [b] Isolated yield. [c] Diastereoselectivity was determined by ¹H NMR analysis of the crude product.

those on the ketone (R¹, R², and R³) in the Sc(OTf)₃-catalyzed case. However, for the substrate in which R³=H (**1j**),

Table 3. Tuning the regioselectivity of the 1,3-dipolar cycloadditions of 1-(1-alkynyl)-cyclopropyl ketones **1** with nitrone **2a**.^[a]



R ¹	R ²	R ³	1	Conditions	t [h]	Yield [%] ^[b]	d.r. ^[c]
1	Me	cyclopropyl	Ph	1b A	7	3ba 73	6.1:1
2			Ph	1b B	0.3	4ba 99	>20:1
3	Me	<i>n</i> -C ₄ H ₉	Ph	1c A	48	3ca 52	6.7:1
4			Ph	1c B	0.3	4ca 81	>20:1
5	Me	1-cyclohexenyl	Ph	1d A	21	3da 73	13.3:1
6			Ph	1d B	0.3	4da 94	>20:1
7	Me	AcOC ₂ H ₄	Ph	1e A	8	3ea 65	7.3:1
8			Ph	1e B	0.25	4ea 71	>20:1
9	Me	1-naphthyl	Ph	1f A	21	3fa 84	5.3:1
10			Ph	1f B	0.3	4fa 73	>20:1
11	Me	4-MeOC ₆ H ₄	4-MeOC ₆ H ₄	1g A	5	3ga 72	9.0:1
12			4-MeOC ₆ H ₄	1g B	0.6	4ga 86	>20:1
13	Me	Ph	4-MeOC ₆ H ₄	1h A	5.5	3ha 93	1.5:1
14			Ph	1h B	0.25	4ha 85	>20:1
15	Ph	Ph	Ph	1i A	3.5	3ia 40	9.0:1
16			Ph	1i B	0.3	4ia 81	>20:1
17	Et	Ph	H	1j A	24	3ja 0	–
18			H	1j B	1	4ja 62	>20:1

[a] Conditions A: Sc(OTf)₃ (10 mol %), 1,10-phenanthroline monohydrate (10 mol %), 4 Å MS, DCE, 28–32°C; conditions B: Ph₃PAuOTf (2 mol %), CH₂Cl₂, RT. [b] Isolated yield. [c] Diastereoselectivity was determined by ¹H NMR analysis of the crude product.

the reaction did not occur at all under conditions A, but did give the [4+3] cycloadduct **4ja** in 62% yield under conditions B (Table 2, entries 17 and 18). The structures and relative stereochemistries of **4ga** and the major diastereomer of **3ga** were further confirmed by single-crystal X-ray diffraction analysis (Figure 1).^[12]





Figure 1. X-ray crystal structures of compounds **3ga** (upper) and **4ga** (lower).

Furthermore, the gold-catalyst loading can be reduced to only 0.2 mol% and the reaction still proceeds smoothly with no reduction in either the yield, or the diastereoselectivity of a 2.5 mmol scale reaction, which makes this method more practical. For example, the formal [4+3] cycloaddition reaction of **1a** (2.5 mmol) with **2a** (2.75 mmol) can be performed in 10 min to give a 94% yield of **4aa**.


Plausible mechanisms that account for these regioselectively tunable cycloadditions are depicted in Scheme 2. Under conditions A (cycle 1), the Sc(OTf)₃/1,10-phenanthroline complex coordinates selectively to the carbonyl group giving intermediate **A**, which makes the cyclopropane reactive enough to regioselectively react with the nitron and form enolate intermediate **B**.^[6] Subsequent ring closure gives the formal [3+3] cycloadduct and regenerates the catalyst. The major diastereomer may be formed from the more favorable chair-like conformation of intermediate **B**. In contrast, the cationic gold(I) species preferentially binds to the



Scheme 2. Plausible mechanisms for this catalytically tunable, regioselective 1,3-dipolar cycloaddition.

alkyne,^[14] rather than the carbonyl, and, thus, mediates nucleophilic attack to give oxonium-containing vinyl-gold intermediate **C** (cycle 2). This is followed by the regioselective homo-Michael addition of nitron **2** at the more substituted position of the cyclopropyl ring,^[6,2a,15] which produces furanyl-gold intermediate **D**; in turn, this, upon ring closure, gives the formal [4+3] cycloadduct and regenerates the gold catalyst. The high diastereoselectivity of this reaction may result from the more favorable chair-like conformation of intermediate **D**. An alternative, plausible catalytic cycle (cycle 3) for the formation of the formal [4+3] cycloadduct, could not be ruled out by the results described so far. In this catalytic cycle, a key carbocationic furanyl-gold intermediate **E** would be formed. Subsequent nucleophilic addition of the nitron to the carbocation would produce intermediate **D**, which, in turn, would give the final product by the same ring closure step as in cycle 2.

In order to gain an insight into the mechanism of the formation of compounds **4**, a control experiment without the addition of the nitron was performed. This showed that the cycloisomerization of 1-(1-alkynyl)cyclopropyl ketone **1a** into the trisubstituted furan is very slow under conditions B, indicating that the nitron is involved in the cyclopropyl ring-opening step, which is consistent with Schmalz's work.^[2a] Further studies (Scheme 3) on the kinetic resolution^[6f] of a racemic mixture of **1a** and the transformation of optically active *ent*-**1a** provided further strong supporting evidence for the proposed cycle (cycle 2, Scheme 2). For example, the reaction of racemic **1a** with nitron **2a** under the catalysis of the (*R*)-C₃-TunePhos-derived^[16] gold complex was quenched, before completion, after 29 h and gave a 55% yield of *ent*-**4aa** in 38% *ee* and 19% of *ent*-**1a**, in 84% *ee*, was recovered. However, if the reaction was allowed to go to completion a racemic mixture of cycloadduct **4aa** was formed. Interestingly, optically active *ent*-**1a** reacted smoothly with nitron **2a** under the catalysis of



Scheme 3. Kinetic resolution of *rac*-**1a** and transformation of the recovered *ent*-**1a**; L* = (*R*)-C₃-TunePhos.

Ph₃PAuCl/AgOTf to give the optically active *ent*-**4aa'** in high yield with the same level of enantiomeric purity as the starting materials. With these results, the reaction pathway through carbocationic furanyl–gold intermediate **E** (cycle 3) can be ruled out.

In summary, we have developed a Lewis acid catalyzed, regioselective, 1,3-dipolar cycloaddition of nitrones to 1-(1-alkynyl)cyclopropyl ketones, in which the cycloaddition pattern can be tuned by the selection of the catalyst. In the presence of the Sc(OTf)₃/1,10-phenanthroline catalyst, the reaction undergoes formal [3+3] cycloaddition to afford highly substituted, multifunctionalized tetrahydro-1,2-oxazines in moderate to excellent yields with up to 15.7:1 diastereomeric ratio, whereas 5,7-fused bicyclic furo[3,4-*d*]-[1,2]oxazepines can be obtained in good to excellent yields with excellent diastereoselectivity from a gold-catalyzed formal [4+3] cycloaddition. Further studies, including kinetic resolution and synthetic applications, are ongoing in our laboratories and will be reported in due course.

Experimental Section

For preparative procedures and spectroscopic data for all new compounds, see the Supporting Information.

Synthesis of 3aa (conditions A): In a glove box, Sc(OTf)₃ (0.025 mmol, 12.3 mg), 1,10-phenanthroline monohydrate (0.025 mmol, 5.0 mg), 4 Å MS (50 mg), and DCE (2 mL) were added to a dry Schlenk tube. After the solution was stirred for 15 min, nitron **2a** (74.0 mg, 0.375 mmol) and ketone **1a** (0.25 mmol, 65.0 mg) were added. Following stirring for 7 h at 28–32 °C, the reaction was complete, as determined by TLC analysis. Filtration and concentration under reduced pressure provided the residue, which was purified by flash column chromatography on silica gel (hexanes/diethyl ether = 30:1) to afford the pure product **3aa** (103.0 mg, d.r. = 11.5:1) in 90% yield.

Synthesis of 4aa (conditions B): A solution of Ph₃PAuOTf (1 mL, 0.01 M in CH₂Cl₂, 2 mol%) was added to a solution of ketone **1a** (130 mg, 0.5 mmol) and nitron **2a** (102.9 mg, 0.55 mmol, 1.1 equivalents) in CH₂Cl₂ (1 mL) at room temperature. The resulting mixture was stirred

for 10 min at room temperature until the reaction was complete, as determined by TLC analysis. After filtration and concentration under reduced pressure, the residue was purified by flash column chromatography on silica gel (hexanes/diethyl ether = 50:1) to afford the pure product **4aa** (219.1 mg, d.r. = >20:1) in 96% yield as a white solid.

Acknowledgements

We are grateful to the NSFC (20702015) for financial support. This work was also sponsored by the 973 program (2009CB825300), the Shanghai Shuguang Program (07SG27), the Shanghai Pujiang Program (07pj14039), and the Shanghai Leading Academic Discipline Project (B409).

Keywords: cycloaddition • cyclopropane • Lewis acids • regioselectivity • stereoselectivity

- [1] For selected recent examples, see: a) S. Ma, J. Zhang, *Angew. Chem.* **2003**, *115*, 193; *Angew. Chem. Int. Ed.* **2003**, *42*, 183; b) S. Ma, J. Zhang, *J. Am. Chem. Soc.* **2003**, *125*, 12386; c) S. Ma, L. Lu, J. Zhang, *J. Am. Chem. Soc.* **2004**, *126*, 9645; d) S. Kamijo, C. Kanazawa, Y. Yamamoto, *J. Am. Chem. Soc.* **2005**, *127*, 9260; e) B. Alcaide, P. Almendros, T. Martínez del Campo, *Angew. Chem.* **2007**, *119*, 6804; *Angew. Chem. Int. Ed.* **2007**, *46*, 6684; f) S. Chuprakov, V. Gevorgyan, *Org. Lett.* **2007**, *9*, 4463; g) A. S. Dudnik, A. W. Sromek, M. Rubina, J. T. Kim, A. V. Kel'in, V. Gevorgyan, *J. Am. Chem. Soc.* **2008**, *130*, 1440; h) X. Jiang, X. Ma, Z. Zheng, S. Ma, *Chem. Eur. J.* **2008**, *14*, 8572.
- [2] a) J. Zhang, H.-G. Schmalz, *Angew. Chem.* **2006**, *118*, 6856; *Angew. Chem. Int. Ed.* **2006**, *45*, 6704; b) G. Zhang, X. Huang, G. Li, L. Zhang, *J. Am. Chem. Soc.* **2008**, *130*, 1814; c) G. Li, X. Huang, L. Zhang, *J. Am. Chem. Soc.* **2008**, *130*, 6944.
- [3] Y. Zhang, Z. Chen, Y. Xiao, J. Zhang, *Chem. Eur. J.* **2009**, *15*, 5208.
- [4] a) H. Gao, J. Zhang, *Adv. Synth. Catal.* **2009**, *351*, 85; b) Y. Xiao, J. Zhang, *Chem. Commun.* **2009**, 3594; c) L. Liu, J. Zhang, *Angew. Chem.* **2009**, *121*, 6209; *Angew. Chem. Int. Ed.* **2009**, *48*, 6093.
- [5] F. Liu, Y. Yu, J. Zhang, *Angew. Chem.* **2009**, *121*, 5613; *Angew. Chem. Int. Ed.* **2009**, *48*, 5505.
- [6] For the Lewis acid catalyzed formal [3+3] cycloadditions of donor-acceptor cyclopropanes with nitrones, see: a) I. S. Young, M. A. Kerr, *Angew. Chem.* **2003**, *115*, 3131; *Angew. Chem. Int. Ed.* **2003**, *42*, 3023; b) M. D. Ganton, M. A. Kerr, *J. Org. Chem.* **2004**, *69*, 8554; for the total synthesis of (+)-phyllantidine and nakadomarin A from a cyclopropane, see: c) C. A. Carson, M. A. Kerr, *Angew. Chem.* **2006**, *118*, 6710; *Angew. Chem. Int. Ed.* **2006**, *45*, 6560; d) I. S. Young, M. A. Kerr, *J. Am. Chem. Soc.* **2007**, *129*, 1465; e) for asymmetric catalysis, see: M. P. Sibi, Z. Ma, C. P. Jasperse, *J. Am. Chem. Soc.* **2005**, *127*, 5764; f) Y.-B. Kang, X.-L. Sun, Y. Tang, *Angew. Chem.* **2007**, *119*, 3992; *Angew. Chem. Int. Ed.* **2007**, *46*, 3918; g) for a minireview, see: F. Cardona, A. Goti, *Angew. Chem.* **2005**, *117*, 8042; *Angew. Chem. Int. Ed.* **2005**, *44*, 7832.
- [7] a) N,O-heterocycles, such as 1,2-oxazines and oxazepanes, are of great interest due to their presence in a number of biologically active compounds, see: D. Letnicer, *Organic Chemistry of Drug Synthesis*, Vol. 7, Wiley, New York, **2008**; for reviews on the synthesis of 1,2-oxazines, see: b) T. L. Gilchrist, J. E. Wood in *Comprehensive Heterocyclic Chemistry II*, Vol. 6, Elsevier, Oxford, **1996**, pp. 279–299; c) T. L. Gilchrist, J. E. Wood in *Comprehensive Heterocyclic Chemistry II*, Vol. 6, Elsevier, Oxford, **1996**, pp. 1177–1307; d) P. G. Tsoungas, *Heterocycles* **2002**, *57*, 1149; for the synthesis of 1,2-oxazines, see: e) S. C. Yoon, K. Kim, Y. J. Park, *J. Org. Chem.* **2001**, *66*, 7334; f) R. Pulz, S. Cicchi, A. Brandi, H.-U. Reißig, *Eur. J. Org. Chem.* **2003**, 1153; g) Y. Yamamoto, N. Momiyama, H. Yamamoto,

- J. Am. Chem. Soc.* **2004**, *126*, 5962; h) S. Kumarn, D. M. Shaw, D. A. Longbottom, S. V. Ley, *Org. Lett.* **2005**, *7*, 4189; i) M. Helms, W. Schade, R. Pulz, T. Watanabe, A. Al-Harrasi, L. Fišera, I. Hlobilová, G. Zahn, H.-U. Reißig, *Eur. J. Org. Chem.* **2005**, 1003; j) S. Kumarn, D. M. Shaw, S. V. Ley, *Chem. Commun.* **2006**, 3211; k) C. Winter, N. Krause, *Angew. Chem.* **2009**, *121*, 6457; *Angew. Chem. Int. Ed.* **2009**, *48*, 6339.
- [8] For recent reviews, accounts, and highlights dealing with the synthesis of furans, see: a) R. C. D. Brown, *Angew. Chem.* **2005**, *117*, 872; *Angew. Chem. Int. Ed.* **2005**, *44*, 850; b) S. F. Kirsch, *Org. Biomol. Chem.* **2006**, *4*, 2076; c) D. M. D'Souza, T. J. J. Müller, *Chem. Soc. Rev.* **2007**, *36*, 1095; d) N. T. Patil, Y. Yamamoto, *ARKIVOC* **2007**, x, 121; e) G. Balme, D. Bouyssi, N. Monteiro, *Heterocycles* **2007**, *73*, 87; f) X. L. Hou, Z. Yang, K.-S. Y. H. N. C. Wong in *Heterocyclic Chemistry, Vol. 19* (Eds.: G. W. Gribble, J. A. Joule), Pergamon Press, Oxford, **2008**, pp. 176; g) S. F. Kirsch, *Synthesis* **2008**, 3183.
- [9] After the submission of an earlier version of this manuscript, Wang and his co-workers reported a similar work on a [4+3] cycloaddition by the employment of AuCl₃ (10 mol%) or Cu(OTf)₂ (20–30 mol%), see: Y. Bai, J. Fang, J. Ren, Z. Wang, *Chem. Eur. J.* **2009**, *15*, 8975.
- [10] The reaction is run in a glove box, which gives reproducible results, since the reaction is sensitive to water, and the temperature inside is varied from 28–32 °C. For more reaction conditions, see the Supporting Information.
- [11] For reviews on gold-catalyzed reactions published since 2008, see: a) Z. Li, C. Brouwer, C. He, *Chem. Rev.* **2008**, *108*, 3239; b) A. Arcadi, *Chem. Rev.* **2008**, *108*, 3266; c) E. Jimenez-Nunez, A. M. Echavarren, *Chem. Rev.* **2008**, *108*, 3326; d) D. J. Gorin, B. D. Sherry, F. D. Toste, *Chem. Rev.* **2008**, *108*, 3351; e) N. T. Patil, Y. Yamamoto, *Chem. Rev.* **2008**, *108*, 3395; f) R. A. Widenhoefer, *Chem. Eur. J.* **2008**, *14*, 5382; g) H. C. Shen, *Tetrahedron* **2008**, *64*, 3885; h) A. S. K. Hashmi, *Angew. Chem.* **2008**, *120*, 6856; *Angew. Chem. Int. Ed.* **2008**, *47*, 6754; i) N. Marion, S. P. Nolan, *Chem. Soc. Rev.* **2008**, *37*, 1776; j) A. S. K. Hashmi, M. Rudolph, *Chem. Soc. Rev.* **2008**, *37*, 1766; k) N. Bongers, N. Krause, *Angew. Chem.* **2008**, *120*, 2208; *Angew. Chem. Int. Ed.* **2008**, *47*, 2178.
- [12] CCDC-741978 (**3ga**) and 741926 (**4ga**) contain the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/data_request/cif.
- [13] B. T. Shireman, M. J. Miller, *J. Org. Chem.* **2001**, *66*, 4809.
- [14] For recent examples of gold-activated alkynes, see: a) N. Asao, K. Takahashi, S. Lee, T. Kasahara, Y. Yamamoto, *J. Am. Chem. Soc.* **2002**, *124*, 12650; b) J. P. Markham, S. T. Staben, F. D. Toste, *J. Am. Chem. Soc.* **2005**, *127*, 9708; c) E. Jiménez-Núñez, C. K. Claverie, C. Nieto-Oberhuber, A. M. Echavarren, *Angew. Chem.* **2006**, *118*, 5578; *Angew. Chem. Int. Ed.* **2006**, *45*, 5452; d) S. F. Kirsch, J. T. Binder, C. Liébert, H. Menz, *Angew. Chem.* **2006**, *118*, 6010; *Angew. Chem. Int. Ed.* **2006**, *45*, 5878; e) P. Dubé, F. D. Toste, *J. Am. Chem. Soc.* **2006**, *128*, 12062; f) J.-J. Lian, P.-C. Chen, Y.-P. Lin, H.-C. Ting, R.-S. Liu, *J. Am. Chem. Soc.* **2006**, *128*, 11372; g) Y. Yamamoto, *J. Org. Chem.* **2007**, *72*, 7817; h) T. Jin, Y. Yamamoto, *Org. Lett.* **2007**, *9*, 5259; i) J.-M. Tang, S. Bhunia, S. M. A. Sohel, M.-Y. Lin, H.-Y. Liao, S. Datta, A. Das, R.-S. Liu, *J. Am. Chem. Soc.* **2007**, *129*, 15677; j) S. F. Kirsch, J. T. Binder, B. Crone, A. Duschek, T. T. Haug, C. Liébert, H. Menz, *Angew. Chem.* **2007**, *119*, 2360; *Angew. Chem. Int. Ed.* **2007**, *46*, 2310; k) G. Li, L. Zhang, *Angew. Chem.* **2007**, *119*, 5248; *Angew. Chem. Int. Ed.* **2007**, *46*, 5156; l) N. D. Shapiro, F. D. Toste, *J. Am. Chem. Soc.* **2008**, *130*, 9244; m) J. T. Binder, B. Crone, T. T. Haug, H. Menz, S. F. Kirsch, *Org. Lett.* **2008**, *10*, 1025; n) T. Jin, Y. Yamaoto, *Org. Lett.* **2008**, *10*, 3137; o) B. Baskar, H. J. Bae, S. E. An, J. Y. Cheong, Y. H. Rhee, A. Duschek, S. F. Kirsch, *Org. Lett.* **2008**, *10*, 2605; p) A. S. K. Hashmi, M. Rudolph, J. Huck, W. Frey, J. W. Bats, M. Hamzić, *Angew. Chem.* **2009**, *121*, 5962; *Angew. Chem. Int. Ed.* **2009**, *48*, 5848.
- [15] For DFT studies on the mechanism and regioselectivity of the gold(I)-catalyzed reactions of 1-(1-alkynyl)cyclopropyl ketones with nucleophiles (ref. [2a]), see: a) R. Fang, C.-Y. Su, C. Zhao, D. L. Phillips, *Organometallics* **2009**, *28*, 741; b) J. Zhang, W. Shen, L. Li, M. Li, *Organometallics* **2009**, *28*, 3129.
- [16] (*R*)-C₃-TunePhos is a commercially available ligand; for its synthesis, see: Z. Zhang, H. Qian, J. Longmire, X. Zhang, *J. Org. Chem.* **2000**, *65*, 6223.

Received: December 5, 2009
Published online: April 15, 2010