

Novel 2- and 4-Substituted 1*H*-Imidazo[4,5-*c*]quinolin-4-amine Derivatives as Allosteric Modulators of the A₃ Adenosine Receptor

Yoonkyung Kim,^{†,‡,§} Sonia de Castro,^{†,‡} Zhan-Guo Gao,[†] Adriaan P. IJzerman,^{||} and Kenneth A. Jacobson^{*,†}

Molecular Recognition Section, Laboratory of Bioorganic Chemistry, National Institute of Diabetes and Digestive and Kidney Diseases, National Institutes of Health, Bethesda, Maryland 20892, and Division of Medicinal Chemistry, Leiden/Amsterdam Center for Drug Research, Leiden University, P.O. Box 9502, 2300 RA, Leiden, The Netherlands

Received December 31, 2008


4-Arylamino and 2- cycloalkyl (including amino substitution) modifications were made in a series of 1*H*-imidazo[4,5-*c*]quinolin-4-amine derivatives as allosteric modulators of the human A₃ adenosine receptor (AR). In addition to allosteric modulation of the maximum functional efficacy (in [³⁵S]GTPγS G protein binding assay) of the A₃AR agonist Cl-IB-MECA (**15**), some analogues also weakly inhibited equilibrium radioligand binding at ARs. 4-(3,5-Dichlorophenylamino) (**6**) or 2-(1-adamantyl) (**20**) substitution produced allosteric enhancement (twice the maximal agonist efficacy), with minimal inhibition of orthosteric AR binding. 2-(4-Tetrahydropyranyl) substitution abolished allosteric enhancement but preserved inhibition of orthosteric binding. Introduction of nitrogen in the six-membered ring at the 2 position, to improve aqueous solubility and provide a derivatization site, greatly reduced the allosteric enhancement. 2-(4-(Benzoylamino)cyclohexyl) analogues **23** and **24** were weak negative A₃AR modulators. Thus, consistent with previous findings, the allosteric and orthosteric inhibitory A₃AR effects in imidazoquinolines are structurally separable, suggesting the possible design of additional derivatives with enhanced positive or negative allosteric A₃AR activity and improved selectivity in comparison to inhibition of orthosteric binding.

Introduction

The adenosine receptors (ARs^a), of which A₁, A_{2A}, A_{2B}, and A₃ subtypes have been defined, represent a physiologically important family of G-protein-coupled receptors (GPCRs).^{1,2} ARs are important pharmacological targets in the treatment of a variety of diseases because of their key roles in controlling numerous cellular processes. For example, A₃AR agonists are of interest for the treatment of cardiac ischemia, bowel inflammation, protection of skeletal muscle, cancer, and rheumatoid arthritis.^{3–7} However, therapeutic intervention using a selective AR agonist is subject to side effects related, in part, to the widespread occurrence of the corresponding receptor throughout the body.

Native agonists of a given GPCR bind at a principal (orthosteric) site on the receptor protein to effect its activation. However, an allosteric modulator would bind to a distinct site on the receptor to either enhance (positive modulator) or impede (negative modulator) the action of a native agonist.^{8–11} The therapeutic application of allosteric modulation has advantages over directly acting orthosteric GPCR agonists. In a particular

Chart 1. Structures of a Series of 2-Cycloalkylimidazoquinoline Derivatives Previously Found To Be Positive Allosteric Modulators of the Human A₃AR



disease state, the effect of an endogenous agonist, which may be insufficient to fully compensate for an imbalance, may be magnified in a temporally and/or spatially specific manner by a positive allosteric modulator for therapeutic benefit.¹² The allosteric modulator theoretically would have no effect of its own on the unoccupied receptor. An additional advantage of allosteric modulators is that they typically are found to display higher subtype-selectivity than orthosteric ligands of the same receptor. Thus, allosteric action that is dependent on the simultaneous presence of an endogenous ligand ideally can produce a more selective drug action and prevent side effects and possible overdose associated with the administration of a conventional orthosteric agonist.

Allosteric modulators of several subtypes of ARs have been reported and their structure–activity relationships (SARs) explored.^{8,13,14} *N*-(3,4-Dichlorophenyl)-2-cyclohexyl-1*H*-imidazo[4,5-*c*]quinolin-4-amine (LUF6000, **3**, Chart 1)^{15–17} is an allosteric modulator of the human A₃AR that increases the maximum efficacy of the agonist 2-chloro-*N*⁶-(3-iodobenzyl)adenosine-5'-*N*-methylcarboxamide (Cl-IB-MECA, **15**). A six-membered ring provided the optimal A₃AR enhancement in the homologous series of 2-cycloalkyl derivatives **1–4**. Compound **3** enhanced the A₃AR agonist efficacy in a functional assay and decreased the agonist dissociation rate without influencing

* To whom correspondence should be addressed. Phone: 301-496-9024. Fax: 301-480-8422. E-mail: kajacobs@helix.nih.gov.


[†] National Institute of Diabetes and Digestive and Kidney Diseases.

[‡] Contributed equally to this work.

[§] Current address: Biomedical Research Center, Korea Institute of Science and Technology, 39-1 Hawolgok-dong, Seongbuk-gu, Seoul, 136-791, Korea.

^{||} Leiden/Amsterdam Center for Drug Research.

^a Abbreviations: AR, adenosine receptor; Cl-IB-MECA, 2-chloro-*N*⁶-(3-iodobenzyl)adenosine-5'-*N*-methylcarboxamide; CCPA, 2-chloro-*N*⁶-cyclopentyladenosine; CHAPS, 3-[(3-cholamidopropyl)-dimethylammonio]-1-propanesulfonate; DMEM, Dulbecco's modified Eagle's medium; DMF, *N,N*-dimethylformamide; DMSO, dimethyl sulfoxide; ESI, electrospray ionization; GPCR, G-protein-coupled receptor; GTPγS, guanosine 5'-(γ-thiotriphosphate); I-AB-MECA, *N*⁶-(4-amino-3-iodobenzyl)adenosine-5'-*N*-methylcarboxamide; *m*-CPBA, 3-chloroperoxybenzoic acid; NECA, 5'-*N*-ethylcarboxamidoadenosine; SAR, structure–activity relationship; TLC, thin layer chromatography.

Scheme 1. Synthesis of Novel 1*H*-Imidazo[4,5-*c*]quinolin-4-amine Derivatives with Structural Variation at the 4 Position^a

^a Reagents: (i) *m*-CPBA, CHCl₃/CH₂Cl₂/MeOH; (ii) POCl₃, toluene/DMF; (iii) R-PhNH₂, DMF.

agonist potency, because of decreased interaction with the orthosteric binding site on the A₃AR. Since the structural requirements for allosteric enhancement at the A₃AR are distinct from the requirements to inhibit equilibrium binding, structural manipulation of this family of imidazoquinolines might achieve even greater selectivity.

In the present study we have extended our search for highly effective allosteric enhancer ligands for the A₃AR, by modifying the substitutions around the 4-arylamino and 2-cycloalkyl moieties of the imidazoquinoline scaffold. We have also identified weak negative allosteric modulators in the same structural series.

Results and Discussion

The novel 4-substituted imidazoquinoline derivatives **5–14** (Scheme 1) were prepared from a key 4-chloro-2-cyclohexyl intermediate **29**. Oxidation of **27**¹⁵ with 3-chloroperoxybenzoic acid (*m*-CPBA) afforded the 5-oxide derivative **28**, which was subsequently converted with phosphorus oxychloride into a 4-chloro species **29**.^{15,18} Reaction of **29** with the appropriately substituted aniline provided the desired 4-amino derivatives **5–14** in varying yields (4–100%), where the 3-aminopyridyl derivative **14** has the lowest yield. Attempts to prepare an *o*-pyridyl derivative (i.e., using 2-aminopyridine) under the same reaction condition with **29** failed.


The novel 2-substituted imidazoquinoline derivatives **16–26** (Scheme 2) were prepared similarly. However, here different 2-cycloalkyl groups were appended, each as its carboxylic acid form, to a common intermediate **30** in an earlier synthetic step to give the corresponding imidazoquinoline derivatives **31a–f**. In fact, compound **30** also served as a precursor for **27**.¹⁵ Next, compounds **31a–f** were treated with *m*-CPBA to afford 5-oxide derivatives **32a–f**, followed by phosphorus oxychloride to give the respective 4-chloro compounds **33a–f**.^{15,18} Reaction of the 4-chloro-2-cycloalkyl derivatives **33a–f** with 3,4-dichloroaniline afforded the desired compounds (**16–21**, **23**). Aniline was used instead in the synthesis of compound **24**.

Several alkylamino derivatives, **22** and **26**, were included. In the case of a 2-(4-*cis*-aminocyclohexyl) analogue **22**, it was necessary to remove the amine-protecting group of **21** in order to obtain the final compound. The deprotection was accomplished using methylamine in ethanol. In the case of the piperidine derivative **26**, the reduction of the carbonyl group of **23** with lithium aluminum hydride afforded the *N*-benzyl derivative **25**, which was subsequently converted into the desired compound **26** by hydrogenation.


To obtain a general procedure for the aniline substitution reaction, we first attempted the synthesis following the method recently described by Göblyös et al.,¹⁵ using a microwave reactor (in ethanol under nitrogen, prestirring for 60 s, at 120 °C for 40 min, normal sample absorption, fixed hold time). Although the reaction was performed on a very small scale, we were able to determine preferred conditions to be *N,N*-dimethylformamide (DMF) as a solvent and using 2–3 equiv of corresponding aniline with heating at 140 °C overnight under a N₂ atmosphere in a tightly sealed Biotage reaction vial. In an earlier attempted reaction at 105 °C under the same conditions, no desired product was obtained.¹⁸

The structures of the imidazoquinoline derivatives were analyzed by NMR in dimethyl sulfoxide (DMSO-*d*₆). In all final compounds except the 2-*exo*-norbornyl-4-(3,4-dichlorophenyl)-amino analogue **18** that was attached in an axial position, the central imidazoquinoline system was clearly attached to the six-membered ring in an equatorial position. This was demonstrated based on the *J* coupling of the proton at the junction (*J*_{ax-ax} ≈ 12–11 and *J*_{ax-eq} ≈ 5–3). We also noted the fact that potentially there are two possible annular tautomers (1*H* or 3*H* at the imidazole ring), which may affect the GPCR binding equilibrium depending on the stability of each form in the aqueous media. 2D NOESY experiments in DMSO-*d*₆ (see Supporting Information) suggested the 3*H*-tautomer as an exclusive form for compounds **10** and **11**. Interestingly, for compound **14**, two tautomers were observed as correlated by NOE cross-peaks (1*H*-/3*H*- ≈ 88:12 by NMR integration in DMSO-*d*₆). Unlike other compounds, detection of two competing tautomers on the NMR time scale could be due to the formation of a hydrogen bond between the pyridyl nitrogen (acceptor) and the imidazole NH (donor) as a less preferred 3*H*-tautomer.

All of the imidazoquinoline derivatives were evaluated for interaction with the human A₃ receptors and two other ARs as listed in Tables 1 and 2. We first tested the effect of these compounds on the equilibrium binding at A₁, A_{2A}, and A₃ARs using standard agonist radioligands^{19–21} [³H]2-Chloro-*N*⁶-cyclopentyladenosine ([³H]CCPA, **34**), [³H]2-[4-(2-carboxylethyl)phenylethylamino]-5'-*N*-ethylcarboxamidoadenosine ([³H]CGS21680, **35**), and [¹²⁵I]*N*⁶-(4-amino-3-iodobenzyl)adenosine-5'-*N*-methylcarboxamide ([¹²⁵I]I-AB-MECA, **36**), respectively. For compounds **5–14** and **16–26**, only the percent inhibition of orthosteric radioligand binding was reported rather than *K*_i values because the affinity at all three subtypes is weak and close to the solubility limits of the compounds. It is unknown

Scheme 2. Synthesis of Novel 1*H*-Imidazo[4,5-*c*]quinolin-4-amine Derivatives with Structural Variation at the 2 Position^a

^a Reagents: (i) polyphosphoric acid, R¹CO₂H; (ii) *m*-CPBA, CHCl₃/CH₂Cl₂/MeOH; (iii) POCl₃, toluene/DMF; (iv) R²-PhNH₂, DMF; (v) LiAlH₄, THF; (vi) MeNH₂, EtOH; (vii) H₂/Pd, MeOH.

Table 1. Potency of 4-Anilino Substituted 1*H*-Imidazo[4,5-*c*]quinolin-4-amine Derivatives in Binding Assays at the Human A₁ and A₃ARs Expressed in CHO Cells and at the Human A_{2A} in HEK-293 Cells and the Allosteric Effects at the Human A₃AR^a



compd	R	A ₁ AR, % displ at 10 μM	A _{2A} AR, % displ at 10 μM	A ₃ AR, % displ at 10 μM	A ₃ AR ag dissociation, ^b % at 10 μM	[³⁵ S]GTPγS binding in A ₃ AR cells, ^c % at 10 μM
3	3,4-Cl ₂ -PhNH	37 ± 9	-11.2 ± 2.8	40.5 ± 13.4	192 ± 7	208 ± 14
5	3-Cl-PhNH	53.3 ± 3.8	42.8 ± 3.7	79.2 ± 1.3	194 ± 3	200 ± 13
6	3,5-Cl ₂ -PhNH	16.7 ± 2.9	-11.4 ± 10.9	32.7 ± 4.8	184 ± 5	209 ± 22
7	2,4-Cl ₂ -PhNH	48.0 ± 1.1	67.2 ± 2.4	43.9 ± 0.1	174 ± 5	187 ± 20
8	3,4-F ₂ -PhNH	14.6 ± 4.8	26.5 ± 7.3	68.8 ± 1.2	187 ± 1	197 ± 17
9	3,5-F ₂ -PhNH	36.6 ± 5.3	17.2 ± 2.1	70.1 ± 7.9	188 ± 3	171 ± 7
10	3,5-(CF ₃) ₂ -PhNH	7.2 ± 2.3	15.6 ± 0.3	21.8 ± 1.4	137 ± 5	143 ± 3
11	3,4-(CN) ₂ -PhNH	33.1 ± 4.1	37.2 ± 1.5	42.2 ± 2.9	129 ± 7	138 ± 7
12	3,4-(OMe) ₂ -PhNH	75.6 ± 2.1	93.8 ± 1.6	98.4 ± 0.1	173 ± 1	182 ± 14
13	3,4-O ₂ CH ₂ -PhNH	65.4 ± 3.3	49.0 ± 0.7	83.9 ± 1.1	185 ± 6	166 ± 11
14	3-pyridyl-NH	73.1 ± 2.5	78.1 ± 1.8	64.8 ± 6.9	114 ± 5	119 ± 1

^a All experiments were performed using adherent mammalian cells stably transfected with cDNA encoding the human ARs. Binding at human A₁, A_{2A}, and A₃ARs in this study was carried out as described in the Experimental Procedures using [³H]**34**, [³H]**35**, or [¹²⁵I]**36** as a radioligand.^{19–21} Values from the present study are expressed as the mean ± SEM, *n* = 3–5. Percentage of inhibition at A₁, A_{2A}, or A₃ receptors is expressed as the mean value from two to four separate experiments with similar results performed in duplicate. ^b Dissociation: % decrease of [¹²⁵I]**36** dissociation at 60 min (control = 100%). ^c Increase of efficacy in the stimulation of the binding of [³⁵S]GTPγS compared to maximal effect induced by 10 μM **15** alone (set to 100%). It is noted that the *E*_{max} of **15** in this functional assay was recently demonstrated to be about 50% of that of **37**.¹⁷ In the adenylate cyclase assay, **15** and **37** were both full agonists.¹⁵

whether the observed inhibition is of an allosteric or nonallosteric character.

Ability to allosterically modulate the A₃AR was determined using two methods: effects of the imidazoquinoline on the dissociation rate of **36** ([¹²⁵I]I-AB-MECA) and on the binding

to the G protein of the stable GTP analogue [³⁵S]guanosine-5'-(γ-thiotriphosphate) ([³⁵S]GTPγS).²² Depending on the functional assay used, **15** (Cl-IB-MECA) may appear to be either a full¹⁵ or partial agonist^{17,23–25} at the A₃AR. The earlier series of imidazoquinoline derivatives displayed dual

Table 2. Potency of 2-Cycloalkyl Substituted 1*H*-Imidazo[4,5-*c*]quinolin-4-amine Derivatives in Binding Assays at the Human A₁ and A₃ARs Expressed in CHO Cells and at the Human A_{2A} in HEK-293 Cells and the Allosteric Effects at the Human A₃AR^a

R' = 3,4-dichloro, unless otherwise noted

No.	R =	A ₁ AR, %displ. at 10 μM	A _{2A} AR, %displ. at 10 μM	A ₃ AR, %displ. at 10 μM	A ₃ AR ag. dissociation, ^c % at 10 μM	[³⁵ S]GTPγS binding in A ₃ AR cells, ^d % at 10 μM
1 ^b		-5.2	0.4	52	116±3	126±3 ^b
2 ^b		15	0	67 ^c	144±9	141±5 ^b
3		37±9	-11.2±2.8	40.5±13.4	192±7	208±14
4 ^b		-4.2	-2.2	68	130±2	115±7 ^b
16		37.2±8.6	-11.9±2.8	69.6±1.4	115±12	114±7
17		77.9±12.2	5.8±1.0	61.2±1.4	187±16	201±26
18		44.8±10.5	39.4±1.9	50.5±5.5	168±8	168±13
19		14.5±2.7	-17.3±6.9	3.3±4.1	139±10	156±2
20		13.8±6.9	-4.4±12.5	17.2±6.7	196±13	210±12
21		21.4±4.3	56.8±1.7	5.6±4.9	102±5	96±5
22		31.7±16.2	15.8±1.2	4.2±6.1	107±11	111±6
23		20.2±10.1	8.8±1.9	29.8±0.6	91±9	93±7
24		12.4±1.1	43.4±5.5	35.3±1.5	92±5	93±1
25		17.8±3.6	-3.1±1.6	46.3±0.8	93±6	97±5
26		10.9±8.6	7.0±0.7	52.8±1.1	101±7	99±3

^a All experiments were performed using adherent mammalian cells stably transfected with cDNA encoding the human ARs. Binding at human A₁, A_{2A}, and A₃ARs in this study was carried out as described in the Experimental Procedures using [³H]34, [³H]35, or [¹²⁵I]36 as a radioligand.^{19–21} Values from the present study are expressed as the mean ± SEM, *n* = 3–5. Percentage inhibition at A₁, A_{2A}, or A₃ receptors is expressed as the mean value from two to four separate experiments with similar results performed in duplicate. ^b Values from Göblyös et al.,¹⁵ the functional enhancement of which was measured with the cyclic AMP assay.²⁶ ^c Dissociation: % decrease of [¹²⁵I]36 dissociation at 60 min (control = 100%). ^d Increase of efficacy in the stimulation of the binding of [³⁵S]GTPγS compared to maximal effect induced by 10 μM 15 alone (set to 100%). It should be noted that the *E*_{max} of 15 in this functional assay was recently demonstrated to be about 50% of that of 37.¹⁷ In the adenylate cyclase assay, 15 and 37 were both full agonists.¹⁵ ^e *K*_i value in a binding assay:¹⁵ 4690 ± 970 nM.




Figure 1. Allosteric modulation of the human A₃AR by compound **6**. (a) Functional assay of the human A₃AR: % stimulation of binding of [³⁵S]GTPγS by increasing concentrations of **15** under control conditions or in the presence of 10 μM compound **3** or **6**. (b) Radioligand binding studies on the human A₃AR: study of the dissociation kinetics of the agonist radioligand [¹²⁵I]**36** under control conditions and in the presence of 10 μM compound **3** or **6**.

and apparently opposite actions as positive allosteric modulators of agonist action and as inhibitors of radioligand binding.¹⁵ In the previous study, compound **3** induced a substantial functional enhancement of the effects of the A₃AR agonist **15** as determined using an agonist radioligand dissociation kinetic assay and a cyclic AMP assay,^{15,26} while there was only a weak inhibition of equilibrium radioligand binding at ARs at the concentration used.

The potentiation of the maximum efficacy of the agonist **15** by arylhalo derivatives **5–9** (Table 1) was high and similar to that observed with the lead compound **3**. The percent of maximal functional effect ranged from 171% for **9** to 209% for **6**. Among these halo analogues, **6** had the least inhibition of orthosteric binding to ARs. The degree of enhancement of [³⁵S]GTPγS binding by **6** over a range of concentrations of **15** up to 10 μM was indistinguishable from the effect of the lead compound **3** (Figure 1a). Similarly, the marked decrease in dissociation rate of the A₃AR agonist radioligand [¹²⁵I]**36** produced by **6** was indistinguishable from the effect of compound **3** (Figure 1b). A 3,5-di(trifluoromethyl) analogue **10** and another aniline derivative **11** that was disubstituted with electron withdrawing groups displayed an intermediate degree of allosteric enhancement. Other 3,4-disubstituted anilines **12** and **13** that contained electron-donating groups displayed a high degree of allosteric enhancement but also substantial inhibition of orthosteric binding to the A₃AR and other ARs. The 3-pyridinylamine **14** had a low degree of allosteric enhancement.

Highly variable biological activities were observed for the 2 position derivatives **16–26** (Table 2). Replacement of the distal methylene group of the 2-cyclohexyl ring of **3** with an ether oxygen in the tetrahydropyranyl derivative **16** abolished allosteric modulation of the A₃AR but retained a similar degree of orthosteric inhibition. Inclusion of a methylene bridge across the 2-cyclohexyl ring of **3** in **17** and **18** resulted in considerable




Figure 2. Allosteric modulation of the human A₃AR by compound **20** or **24**. (a) Functional assay of the human A₃AR: % stimulation of binding of [³⁵S]GTPγS by increasing concentrations of **15** under control conditions or in the presence of 10 μM compound **3** or **20**. (b) Radioligand binding studies on the human A₃AR: study of the dissociation kinetics of the agonist radioligand [¹²⁵I]**36** under control conditions and in the presence of 10 μM compound **20** or **24**.

allosteric potentiation. Multiply bridged cycloalkyl substitution in the 2-(1-adamantyl)-4-(3,4-dichlorophenyl)amino analogue **20** resulted in high allosteric potentiation with minimal effects on the binding of orthosteric ligands at A₁, A_{2A}, and A₃ARs. The 2-*exo*-noradamantyl-4-(3,4-dichlorophenyl)amino analogue **19** displayed only moderate allosteric potentiation. The (decreasing) order of allosteric enhancement by these bridged cycloalkyl derivatives was **20**, **17** > **18** > **19**. The adamantyl derivative **20** increased [³⁵S]GTPγS binding over a range of concentrations of **15** up to 10 μM in a manner indistinguishable from the effect of **3** (Figure 2a).

Introduction of a nitrogen atom in (or as substituted at) the six-membered ring in **21–26** in an attempt to improve aqueous solubility and to provide a site for further derivatization greatly reduced the allosteric enhancement of this series. Variable degrees of inhibition of orthosteric binding at the AR were observed in this group with the most pronounced inhibition found for **21** and **24** at the A_{2A}AR and for **25** and **26** at the A₃AR. The slightly decreased agonist efficacy (91–93% of control in both functional assays, Table 2) of 2-(*N*-benzoyl-4-piperidinyl) analogues **23** and **24** was suggestive of possible negative allosteric modulation of the A₃AR. The effect of compound **24** on the dissociation kinetics of the agonist radioligand was only slightly different from that of the control (Figure 2b).

Conclusions

In this study, structural modification of the 1*H*-imidazo[4,5-*c*]quinolin-4-amine **3** has demonstrated that a limited set of substituents at the 2 and 4 positions are tolerated to preserve the allosteric enhancement of agonist action at the A₃AR. Notably, the haloanilino derivatives **5**, **6**, and **8** and the bridged 2-cycloalkyl analogues **17** and **20** approximately doubled the

maximum efficacy of the agonist **15** in the [^{35}S]GTP γ S assay. The highest potentiation of the maximum efficacy of the agonist **15**, without increased inhibition of orthosteric binding, was observed for the 2-(1-adamantyl)-4-(3,4-dichlorophenyl)amino analogue **20**, with 210% activity of control in the [^{35}S]GTP γ S binding assay. Compounds **6** and **20** were preferred as selective allosteric enhancers of the A₃AR because of the minimal effect on binding at the orthosteric sites of the three ARs examined. Substitution of a 4-tetrahydropyran moiety at the 2 position completely abolished allosteric enhancement but preserved inhibition of orthosteric binding. Thus, as extension of previous findings, the allosteric and orthosteric inhibitory effects at the A₃AR in this series of imidazoquinolines are structurally separable. These biological results suggest that it will be possible to design additional derivatives that would display enhanced positive or negative allosteric activity at this receptor and improved selectivity in comparison to inhibition of orthosteric binding.

Experimental Procedures

General. Glassware was oven-dried and cooled in a desiccator before use. All reactions were carried out under a dry nitrogen atmosphere. Solvents were purchased as anhydrous grade and used without further purification. Suppliers of some commercial compounds are listed as follows: 3,4-diaminoquinoline (**30**) was purchased from Tyger Scientific, Inc.; 3,5-bis(trifluoromethyl)aniline and 4-aminophthalonitrile were purchased from Acros Organics; polyphosphoric acid, *m*-CPBA, phosphorus oxychloride (POCl₃), chloroform (CHCl₃), methylene chloride (CH₂Cl₂), DMF, toluene, diisopropyl ether, methanol (MeOH), tetrahydrofuran (THF), and most of other reagents and solvents were purchased from Sigma-Aldrich; dimethyl sulfoxide (DMSO-*d*₆), chloroform-*d* (CDCl₃), and CD₃OD were purchased from Cambridge Isotope Laboratories. All reagents were of commercial grade and were used without further purification unless otherwise noted.

NMR spectra were recorded on either a Varian Inova/Gemini 300 or a Bruker DRX-600 spectrometer at 25.0 °C under an optimized parameter setting for each sample, unless otherwise mentioned. For compounds **5–14**, **28**, and **29**, ¹H NMR chemical shifts were measured relative to the residual solvent peak at 2.50 ppm in DMSO-*d*₆ and at 3.31 ppm in CD₃OD or in a mixture of CD₃OD/CDCl₃. For compounds **16–26**, **31a–f**, **32a–f**, and **33a–f**, ¹H NMR chemical shifts were measured relative to tetramethylsilane at 0.00 ppm in CDCl₃ and the residual water peak at 3.30 ppm in CD₃OD. ¹³C NMR chemical shifts were measured relative to the residual solvent peak at 49.15 ppm in CD₃OD or in a mixture of CD₃OD/CDCl₃. Suggested NMR peak assignments of some target compounds are shown in Supporting Information and are based on 2D COSY and NOESY experiments.

Analytical or preparative thin layer chromatography (TLC) was performed on either 0.2 mm silica coated sheets with F₂₅₄ indicator (Sigma-Aldrich) or 0.2 mm reversed-phase C18 silica coated sheets (pore size of 60 Å, Whatman Inc.). Visualization of the products on the TLC plate was aided by the use of UV light, ninhydrin, or potassium permanganate. Column chromatography was performed on 230–400 mesh silica gel (pore size of 60 Å, Sigma-Aldrich). The tested imidazoquinoline derivatives were confirmed by HPLC to possess a $\geq 96\%$ purity.

The electrospray ionization (ESI) MS experiments were performed on a Micromass/Waters LCT Premier electrospray time-of-flight (TOF) mass spectrometer coupled with a Waters HPLC system at the Mass Spectrometry Facility, NIDDK, NIH.

General Procedure for 2-Substituted 1H-Imidazo[4,5-*c*]quinoline (31a–f). Polyphosphoric acid (1.3 mL/mmol) was added to 3,4-diaminoquinoline (**30**) (100 mg, 0.63 mmol) and the appropriate carboxylic acid (1.2 equiv). The mixture was stirred at 100 °C for 5 h. Then the mixture was cooled to 0 °C and to it was slowly added NH₄OH till pH 8–9. The mixture was extracted with ethyl acetate (3 \times 15 mL), and the combined organic extracts were

washed with water, brine, and again with water, and then dried over MgSO₄. The solution was filtered, the solvent was evaporated, and the residue was dried in vacuo. The residue obtained was subjected to preparative silica gel column chromatography (100:1 to 8:1 CHCl₃/MeOH).

2-(Tetrahydro-2H-pyran-4-yl)-1H-imidazo[4,5-*c*]quinoline (31a). Yield: 107.6 mg (67%). ¹H NMR (300 MHz, CD₃OD/CDCl₃) δ 13.40 (s, 1H), 9.13 (s, 1H), 8.35 (s, 1H), 8.08 (m, 1H), 7.65 (m, 2H), 3.98 (m, 2H), 3.35 (dd, 2H, *J* = 11.4, 10.2 Hz), 3.28 (t, 1H, *J* = 11.3 Hz), 2.05 (m, 2H), 1.95 (dd, 1H, *J* = 11.6, 4.2 Hz), 1.93 (dd, 1H, *J* = 11.6, 4.2 Hz); ¹³C NMR (75 MHz, CD₃OD/CDCl₃) δ 143.4, 129.5, 126.3, 121.6, 66.7, 31.07; HRMS (ESI) calcd for C₁₅H₁₆N₃O⁺ (*M* + *H*⁺) 254.1288, found 254.1293.

2-(2-Norbornanyl)-1H-imidazo[4,5-*c*]quinoline (31b). Yield: 86.43 mg (52%). ¹H NMR (300 MHz, CD₃OD/CDCl₃) δ 13.33 (bs, 1H), 9.11 (s, 1H), 8.32 (s, 1H), 8.08 (m, 1H), 7.64 (m, 2H), 3.45 (m, 1H), 3.36 (s, 2H), 3.17 (s, 2H), 3.06 (s, 1H), 2.72 (m, 1H), 2.58 (m, 1H), 2.50 (s, 1H), 2.38 (m, 1H), 1.44 (m, 7H); ¹³C NMR (75 MHz, CD₃OD/CDCl₃) δ 143.85, 129.57, 125.98, 121.43, 48.59, 42.59, 36.61, 35.78, 29.10, 28.59, 23.60; HRMS (ESI) calcd for C₁₇H₁₈N₃⁺ (*M* + *H*⁺) 264.1495, found 264.0561.

2-(4-(1,3-Dioxoisindolin-2-yl)cyclohexyl)-1H-imidazo[4,5-*c*]quinoline (31c). Yield: 88.0 mg (35%). ¹H NMR (300 MHz, CD₃OD/CDCl₃) δ 13.35 (bs, 1H), 9.13 (s, 1H), 8.35 (s, 1H), 8.07 (m, 1H), 7.69 (m, 6H), 4.16 (m, 1H), 3.06 (t, 1H, *J* = 12.2 Hz), 2.36 (m, 4H), 1.88 (m, 4H); ¹³C NMR (75 MHz, CD₃OD/CDCl₃) δ 167.9, 167.7, 143.4, 134.3, 129.5, 126.1, 122.9, 121.7, 49.5, 36.9, 30.7, 28.8; HRMS (ESI) calcd for C₂₄H₂₁N₄O₂⁺ (*M* + *H*⁺) 397.1659, found 397.1234.

2-(1-Benzoylpiperidin-4-yl)-1H-imidazo[4,5-*c*]quinoline (31d). Yield: 142 mg (67%). ¹H NMR (300 MHz, CD₃OD/CDCl₃) δ 13.38 (s, 1H), 9.13 (s, 1H), 8.35 (s, 1H), 8.09 (s, 1H), 7.66 (s, 1H), 7.46 (s, 1H), 4.56 (s, 1H), 3.71 (s, 1H), 3.13 (s, 1H), 2.17 (s, 1H), 1.91 (s, 1H); ¹³C NMR (75 MHz, CD₃OD/CDCl₃) δ 169.1, 143.4, 136.3, 129.4, 128.5, 126.6, 121.4, 59.8, 46.8, 41.2, 35.8, 30.4; HRMS (ESI) calcd for C₂₂H₂₁N₄O⁺ (*M* + *H*⁺) 357.1710, found 357.1732.

2-(3-Noradamantanyl)-1H-imidazo[4,5-*c*]quinoline (31e). Yield: 118.7 mg (68%). ¹H NMR (300 MHz, CD₃OD/CDCl₃) δ 8.98 (s, 1H), 8.36 (s, 1H), 8.03 (d, 1H, *J* = 7.8 Hz), 7.52 (m, 2H), 2.77 (t, 1H, *J* = 6.9 Hz), 2.33 (s, 2H), 2.20 (m, 2H), 2.06 (m, 3H), 1.99 (m, 2H), 1.68 (m, 4H); ¹³C NMR (75 MHz, CD₃OD/CDCl₃) δ 162.7, 143.2, 128.3, 121.5, 45.6, 43.6, 38.1, 37.8, 37.5, 37.3, 34.3, 30.4; HRMS (ESI) calcd for C₁₉H₂₀N₃⁺ (*M* + *H*⁺) 290.1652, found 290.1670.

2-(1-Adamantanyl)-1H-imidazo[4,5-*c*]quinoline (31f). Yield: 75.8 mg (39%). ¹H NMR (300 MHz, CD₃OD/CDCl₃) δ 9.11 (s, 1H), 8.30 (s, 1H), 8.15 (d, 1H, *J* = 8.4 Hz), 7.53 (t, 1H, *J* = 7.2 Hz), 7.43 (d, 1H, *J* = 7.2 Hz), 2.21 (s, 6H), 2.05 (s, 4H), 1.72 (m, 6H); ¹³C NMR (75 MHz, CD₃OD/CDCl₃) δ 143.6, 129.1, 127.4, 126.6, 121.9, 41.6, 36.5, 36.0, 228.3, 1.2; HRMS (ESI) calcd for C₂₀H₂₂N₃⁺ (*M* + *H*⁺) 304.1808, found 304.1823.

General Procedure for 2-Substituted 1H-Imidazo[4,5-*c*]quinoline 5-Oxide (28, 32a–f). The appropriate 2-substituted 1H-imidazo[4,5-*c*]quinoline derivatives (**27**, **31a–f**) in a mixture of CHCl₃ (2.5 mL/mmol), CH₂Cl₂ (2.5 mL/mmol), and MeOH (0.25 mL/mmol) was heated for dissolution. To this mixture was added *m*-CPBA (2.5 equiv), which was then refluxed for 30 min. The mixture was cooled to room temperature, Na₂CO₃ (0.04 g/mmol) was added in one portion as a solid, and then the mixture was refluxed for 1 h. The solvent was removed in vacuo, and the crude product was chromatographed on silica gel (20:1 to 7:1 CH₂Cl₂/MeOH for **28**, 100:1 to 8:1 CHCl₃/MeOH for **32a–f**) to give the desired compound.

2-Cyclohexyl-1H-imidazo[4,5-*c*]quinoline 5-Oxide (28).¹⁵ Scale: 0.612 mmol. Yield: 150 mg (91%). *R*_f = 0.43 [silica gel, 10:1 CH₂Cl₂/MeOH]. ¹H NMR (300 MHz, CD₃OD) δ 9.03 (s, 1H), 8.78–8.72 (m, 1H), 8.47 (m, 1H), 7.89–7.83 (m, 2H), 3.06 (tt, 1H, *J* = 11.8, 3.5 Hz), 2.20–2.14 (m, 2H), 1.98–1.91 (m, 2H), 1.86–1.69 (m, 3H), 1.60–1.32 (m, 3H); ¹³C NMR (75 MHz,

CD₃OD) δ 130.9, 130.6, 130.4, 129.0, 123.5, 121.2, 40.2, 32.9, 27.2, 27.0; HRMS (ESI) calcd for C₁₆H₁₈N₃O (M + H⁺) 268.1450, found 268.1451.

2-(Tetrahydro-2H-pyran-4-yl)-1H-imidazo[4,5-c]quinoline 5-Oxide (32a). Scale: 0.42 mmol. Yield: 65 mg (57%). ¹H NMR (300 MHz, CD₃OD/CDCl₃) δ 9.18 (s, 1H), 8.80 (m, 1H), 8.53 (m, 1H), 7.85 (m, 2H), 4.17 (m, 2H), 3.68 (m, 2H), 3.36 (m, 1H), 2.15 (m, 4H); HRMS (ESI) calcd for C₁₅H₁₆N₃O₂⁺ (M + H⁺) 270.1237, found 270.1032.

2-(2-Norbornanyl)-1H-imidazo[4,5-c]quinoline 5-Oxide (32b). Scale: 0.25 mmol. Yield: 44.4 mg (60%). ¹H NMR (300 MHz, CD₃OD/CDCl₃) δ 8.77 (s, 1H), 8.05 (s, 1H), 7.95 (s, 1H), 7.81 (m, 2H), 3.13 (m, 1H), 2.81 (m, 1H), 2.68 (s, 1H), 2.52 (s, 1H), 2.14 (m, 1H), 1.93 (m, 1H), 1.58 (m, 7H); HRMS (ESI) calcd for C₁₇H₁₈N₃O⁺ (M + H⁺) 280.1444, found 280.1443.

2-(4-(1,3-Dioxoisindolin-2-yl)cyclohexyl)-1H-imidazo[4,5-c]quinoline 5-Oxide (32c). Scale: 0.40 mmol. Yield: 65 mg (44%). ¹H NMR (300 MHz, CD₃OD/CDCl₃) δ 8.96 (s, 1H), 8.63 (m, 1H), 8.32 (m, 1H), 7.75 (m, 6H), 4.18 (m, 1H), 3.04 (t, 1H, *J* = 11.6 Hz), 2.24 (m, 4H), 1.84 (m, 4H); HRMS (ESI) calculated for C₂₄H₂₁N₄O₂⁺ (M + H⁺) 413.1608, found 413.1463.

2-(1-Benzoylpiperidin-4-yl)-1H-imidazo[4,5-c]quinoline 5-Oxide (32d). Scale: 0.30 mmol. Yield: 74 mg (67%). ¹H NMR (300 MHz, CD₃OD/CDCl₃) δ 8.79 (s, 1H), 8.57 (m, 1H), 8.18 (bs, 1H), 7.76 (s, 1H), 7.68 (m, 1H), 7.54 (m, 5H), 4.59 (m, 1H), 3.76 (m, 1H), 2.93 (m, 2H), 2.02 (m, 2H), 1.85 (m, 4H); HRMS (ESI) calcd for C₂₂H₂₀N₄O₂⁺ (M + H⁺) 372.1586, found 373.1665.

2-(3-Noradamantanyl)-1H-imidazo[4,5-c]quinoline 5-Oxide (32e). Scale: 0.41 mmol. Yield: 89 mg (71%). ¹H NMR (300 MHz, CD₃OD/CDCl₃) δ 8.77 (s, 1H), 8.52 (m, 1H), 8.29 (s, 1H), 7.71 (m, 1H), 7.53 (m, 2H), 2.67 (t, 1H, *J* = 6.7 Hz), 2.27 (s, 2H), 2.09 (d, 2H, *J* = 11.4 Hz), 1.95 (d, 1H, *J* = 10.8 Hz), 1.87 (m, 3H), 1.58 (m, 4H); HRMS (ESI) calcd for C₁₉H₂₀N₃O⁺ (M + H⁺) 306.1601, found 306.1606.

2-(1-Adamantanyl)-1H-imidazo[4,5-c]quinoline 5-Oxide (32f). Scale: 0.25 mmol. Yield: 63 mg (78%). ¹H NMR (300 MHz, CD₃OD/CDCl₃) δ 8.91 (m, 1H), 8.60 (m, 1H), 8.31 (m, 1H), 7.90 (s, 1H), 7.80 (d, 1H, *J* = 7.5 Hz), 7.54 (m, 1H), 2.08 (s, 10H), 1.76 (s, 6H); HRMS (ESI) calcd for C₂₀H₂₂N₃O⁺ (M + H⁺) 320.1757, found 320.1753.

4-Chloro-2-cyclohexyl-1H-imidazo[4,5-c]quinoline (29).¹⁵ Compound **28** (150 mg, 560 μ mol) was suspended in a mixture of toluene (0.30 mL) and DMF (0.60 mL) and then was treated with 140 μ L (1.50 mmol) of POCl₃ at 0 °C with stirring. The ice bath was removed, and the mixture was heated at 100 °C for 1.5 h. The mixture was cooled to room temperature, and a few small pieces of ice were added with stirring. Subsequently, the pH was adjusted to 6–7 with solid NaHCO₃. The mixture was sonicated and filtered, rinsing with water and diisopropyl ether, and the collected solid was dried in vacuo to give 97.1 mg (340 μ mol, 61%) of **29** as a beige solid. *R*_f = 0.56 [silica gel, 10:1 CH₂Cl₂/MeOH]; ¹H NMR (600 MHz, DMSO-*d*₆) δ 8.38–8.35 (m, 1H), 8.02–7.99 (m, 1H), 7.71–7.62 (m, 2H), 3.07 (tt, 1H, *J* = 12.0, 3.5 Hz), 2.16–2.10 (m, 2H), 1.97–1.91 (m, 2H), 1.88–1.75 (m, 3H), 1.59–1.35 (m, 3H); ¹³C NMR (75 MHz, CD₃OD) δ 129.5, 129.3, 128.2, 122.8, 40.4, 33.0, 27.4, 27.0; HRMS (ESI) calcd for C₁₆H₁₇ClN₃ (M + H⁺) 286.1111, found 286.1106.

General Procedure for 2-Substituted 4-Chloro-1H-imidazo[4,5-c]quinolines (33a–f). A mixture of toluene (0.45 mL/mmol) and DMF (0.90 mL/mmol) was cooled in an ice bath, and phosphorus oxychloride (2.6 equiv) was added. After 10 min, the appropriate 1H-imidazo[4,5-c]quinolin-5-oxide was added, and the solution was stirred at room temperature for 10 min. Subsequently, the solution was heated to 100 °C for 30 min. When the mixture was cooled, the solvent was evaporated, and the resulting syrup was poured on chipped ice while stirring. The mixture was then warmed to room temperature and carefully adjusted to pH 6–7 with solid NaHCO₃. After 2 h, the formed solid was filtered off, washed with water and diisopropyl ether, and subsequently dried. The residue obtained was subjected to preparative silica gel column chromatography (100:1 to 8:1 CHCl₃/MeOH).

4-Chloro-2-(tetrahydro-2H-pyran-4-yl)-1H-imidazo[4,5-c]quinoline (33a). Scale: 0.27 mmol. Yield: 41 mg (53%). ¹H NMR (300 MHz, CD₃OD/CDCl₃) δ 13.66 (br s, 1H), 8.35 (m, 1H), 8.02 (m, 1H), 7.70 (m, 2H), 4.04 (m, 2H), 3.52 (dd, 2H, *J* = 11.1, 10.8 Hz), 3.25 (m, 1H), 1.95 (m, 4H); ¹³C NMR (75 MHz, CD₃OD/CDCl₃) δ 143.2, 128.1, 126.8, 121.2, 67.6, 36.1, 31.1; HRMS (ESI) calcd for C₁₅H₁₅ClN₃O⁺ (M + H⁺) 288.0898, found 288.0713.

4-Chloro-2-(2-norbornanyl)-1H-imidazo[4,5-c]quinoline (33b). Scale: 0.25 mmol. Yield: 44.4 mg (60%). ¹H NMR (300 MHz, CD₃OD/CDCl₃) δ 8.29 (m, 1H), 7.96 (m, 2H), 7.56 (m, 2H), 3.31 (m, 1H), 3.05 (s, 2H), 2.91 (s, 2H), 2.83 (m, 2H), 2.41 (m, 1H), 2.06 (m, 1H), 1.62 (m, 7H); ¹³C NMR (75 MHz, CD₃OD/CDCl₃) δ 143.0, 127.8, 126.5, 121.4, 42.8, 42.0, 36.3, 36.1, 29.6, 28.6; HRMS (ESI) calcd for C₁₇H₁₇ClN₃⁺ (M + H⁺) 298.1106, found 298.1104.

4-Chloro-2-(4-(1,3-dioxoisindolin-2-yl)-1H-imidazo[4,5-c]quinoline (33c). Scale: 0.27 mmol. Yield: 87 mg (74%). ¹H NMR (300 MHz, CD₃OD/CDCl₃) δ 8.56 (m, 1H), 8.22 (d, 1H, *J* = 7.9 Hz), 8.09 (d, 1H, *J* = 8.5 Hz), 7.70 (m, 6H), 4.25 (m, 1H), 3.35 (m, 1H), 2.42 (m, 4H), 1.85 (m, 4H); ¹³C NMR (75 MHz, CD₃OD/CDCl₃) δ 168.5, 158.7, 144.2, 143.6, 134.1, 132.5, 132.3, 132.2, 129.0, 128.8, 123.3, 50.0, 37.8, 31.3, 29.3; HRMS (ESI) calcd for C₂₄H₂₀ClN₄O₂⁺ (M + H⁺) 431.1269, found 431.1260.

4-Chloro-2-(1-benzoylpiperidin-4-yl)-1H-imidazo[4,5-c]quinoline (33d). Scale: 0.20 mmol. Yield: 42.4 mg (54%). ¹H NMR (300 MHz, CD₃OD/CDCl₃) δ 7.99 (s, 1H), 7.86 (d, 1H, *J* = 7.5 Hz), 7.48 (m, 3H), 7.26 (m, 5H), 4.63 (m, 1H), 3.76 (m, 1H), 3.18 (m, 1H), 2.87 (m, 1H), 2.05 (m, 2H), 1.92 (m, 4H); ¹³C NMR (75 MHz, CD₃OD/CDCl₃) δ 135.9, 129.9, 128.6, 128.3, 128.0, 127.0, 126.7, 121.4, 53.6, 36.8, 30.6; HRMS (ESI) calcd for C₂₂H₂₀ClN₄O⁺ (M + H⁺) 391.1320, found 391.1317.

4-Chloro-2-(3-noradamantanyl)-1H-imidazo[4,5-c]quinoline (33e). Scale: 0.23 mmol. Yield: 60 mg (79%). ¹H NMR (300 MHz, CD₃OD/CDCl₃) δ 8.37 (s, 1H), 8.10 (d, 1H, *J* = 7.8 Hz), 7.78 (m, 1H), 7.38 (m, 2H), 2.67 (m, 1H), 2.21 (s, 2H), 2.08 (m, 2H), 1.90 (m, 5H), 1.54 (s, 4H); ¹³C NMR (75 MHz, CD₃OD/CDCl₃) δ 162.9, 142.9, 128.2, 127.7, 121.4, 45.5, 43.5, 38.1, 37.7, 37.4, 37.2, 36.4, 34.3, 31.2; HRMS (ESI) calcd for C₁₉H₁₉ClN₃⁺ (M + H⁺) 324.1262, found 324.1268.

4-Chloro-2-(1-adamantanyl)-1H-imidazo[4,5-c]quinoline (33f). Scale: 0.20 mmol. Yield: 39 mg (60%). ¹H NMR (300 MHz, CD₃OD/CDCl₃) δ 8.34 (s, 1H), 7.95 (m, 2H), 7.45 (m, 2H), 2.15 (m, 10H), 1.78 (m, 6H); ¹³C NMR (75 MHz, CD₃OD/CDCl₃) δ 163.1, 143.0, 132.6, 127.8, 126.5, 121.5, 40.8, 36.4, 36.2, 35.8, 32.2, 28.1; HRMS (ESI) calcd for C₂₀H₂₁ClN₃⁺ (M + H⁺) 338.1419, found 338.1414.

General Procedure for N-Substituted 2-Cyclohexyl-1H-imidazo[4,5-c]quinolin-4-amine (5–14). A solution of compound **29** and an appropriate aniline in DMF was heated at 140 °C overnight under a dry nitrogen atmosphere in a tightly sealed Biotage microwave vial (size, 0.2–0.5 mL or 0.5–2.0 mL) equipped with a magnetic stir bar. The reaction mixture was cooled, the solvent was removed in vacuo, and the crude product was purified by a preparative TLC.

N-(3-Chlorophenyl)-2-cyclohexyl-1H-imidazo[4,5-c]quinolin-4-amine (5). Compound **29** (6.7 mg, 23 μ mol) was reacted with 3-chloroaniline (10 μ L, 94 μ mol) in DMF (50 μ L). The crude product was chromatographed on silica gel (3:1 petroleum ether/EtOAc) to give 5.5 mg (15 μ mol, 62%) of **5**. *R*_f = 0.43 [silica gel, 2:1 petroleum ether/EtOAc]; ¹H NMR (600 MHz, 4:1 CDCl₃/CD₃OD) δ 8.30 (s, 1H), 7.94 (d, 1H, *J* = 7.4 Hz), 7.89 (d, 1H, *J* = 8.5 Hz), 7.75 (d, 1H, *J* = 8.0 Hz), 7.48 (t, 1H, *J* = 7.5 Hz), 7.38 (t, 1H, *J* = 7.3 Hz), 7.24 (t, 1H, *J* = 7.9 Hz), 6.94 (d, 1H, *J* = 8.3 Hz), 2.92 (t, 1H, *J* = 11.9 Hz), 2.13 (d, 2H, *J* = 12.6 Hz), 1.90 (d, 2H, *J* = 13.3 Hz), 1.78 (d, 1H, *J* = 12.6 Hz), 1.67 (q, 2H, *J* = 11.7 Hz), 1.44 (q, 2H, *J* = 12.7 Hz), 1.32 (q, 1H, *J* = 12.9 Hz); ¹³C NMR (150 MHz, 4:1 CDCl₃/CD₃OD) δ 147.1, 143.9, 134.7, 130.1, 127.7, 127.6, 123.3, 121.8, 120.7, 118.8, 117.0, 115.7, 38.8, 32.2, 26.4, 26.1; HRMS (ESI) calcd for C₂₂H₂₂ClN₄ (M + H⁺) 377.1533, found 377.1528.

2-Cyclohexyl-*N*-(3,5-dichlorophenyl)-1*H*-imidazo[4,5-*c*]quinolin-4-amine (6). Compound **29** (7.4 mg, 26 μ mol) was reacted with 3,5-dichloroaniline (13 mg, 80 μ mol) in DMF (50 μ L). The crude product was chromatographed on silica gel (3:1 petroleum ether/EtOAc) to give 6.0 mg (15 μ mol, 56%) of **6**. R_f = 0.50 [silica gel, 2:1 petroleum ether/EtOAc]; ^1H NMR (600 MHz, 4:1 $\text{CDCl}_3/\text{CD}_3\text{OD}$) δ 8.05 (s, 2H), 7.95 (d, 1H, J = 7.6 Hz), 7.91 (d, 1H, J = 8.3 Hz), 7.49 (t, 1H, J = 7.6 Hz), 7.32 (t, 1H, J = 7.4 Hz), 6.94 (s, 1H), 2.91 (t, 1H, J = 12.0 Hz), 2.13 (d, 2H, J = 11.9 Hz), 1.89 (d, 2H, J = 13.9 Hz), 1.78 (d, 1H, J = 12.2 Hz), 1.66 (q, 2H, J = 12.5 Hz), 1.44 (q, 2H, J = 13.0 Hz), 1.32 (q, 1H, J = 13.1 Hz); ^{13}C NMR (150 MHz, 4:1 $\text{CDCl}_3/\text{CD}_3\text{OD}$) δ 158.1, 146.7, 143.7, 143.2, 135.1, 134.8, 127.8, 127.7, 126.7, 123.6, 121.4, 120.8, 116.9, 115.8, 38.8, 32.2, 26.4, 26.1; HRMS (ESI) calcd for $\text{C}_{22}\text{H}_{21}\text{Cl}_2\text{N}_4$ ($\text{M} + \text{H}^+$) 411.1143, found 411.1140.

2-Cyclohexyl-*N*-(2,4-dichlorophenyl)-1*H*-imidazo[4,5-*c*]quinolin-4-amine (7). Compound **29** (8.2 mg, 29 μ mol) was reacted with 2,4-dichloroaniline (14 mg, 87 μ mol) in DMF (50 μ L). The crude product was chromatographed on silica gel (3:1 hexane/EtOAc) to give 5.0 mg (12 μ mol, 42%) of **7**. R_f = 0.63 [silica gel, 2:1 petroleum ether/EtOAc]; ^1H NMR (600 MHz, 4:1 $\text{CDCl}_3/\text{CD}_3\text{OD}$) δ 8.89 (d, 1H, J = 9.4 Hz), 7.98 (d, 1H, J = 7.7 Hz), 7.84 (d, 1H, J = 8.1 Hz), 7.47 (t, 1H, J = 7.6 Hz), 7.39 (d, 1H, J = 2.2 Hz), 7.33 (t, 1H, J = 7.4 Hz), 7.27 (dd, 1H, J = 8.6, 2.3 Hz), 2.96 (tt, 1H, J = 11.9, 3.5 Hz), 2.11 (d, 2H, J = 12.1 Hz), 1.89 (d, 2H, J = 13.7 Hz), 1.77 (d, 1H, J = 12.3 Hz), 1.70 (q, 2H, J = 12.3 Hz), 1.44 (q, 2H, J = 12.8 Hz), 1.32 (q, 1H, J = 12.7 Hz); ^{13}C NMR (150 MHz, 4:1 $\text{CDCl}_3/\text{CD}_3\text{OD}$) δ 146.7, 136.1, 135.0, 129.0, 127.7, 127.6 (127.615), 127.6 (127.570), 123.6, 122.1, 120.9, 116.1, 39.1, 32.2, 26.4, 26.1; HRMS (ESI) calcd for $\text{C}_{22}\text{H}_{21}\text{Cl}_2\text{N}_4$ ($\text{M} + \text{H}^+$) 411.1143, found 411.1132.

2-Cyclohexyl-*N*-(3,4-difluorophenyl)-1*H*-imidazo[4,5-*c*]quinolin-4-amine (8). Compound **29** (7.0 mg, 24 μ mol) was reacted with 3,4-difluoroaniline (20 μ L, 200 μ mol) in DMF (30 μ L). The crude product was chromatographed on silica gel (7:4 petroleum ether/EtOAc) to give 7.1 mg (19 μ mol, 77%) of **8**. R_f = 0.36 [silica gel, 2:1 petroleum ether/EtOAc]; ^1H NMR (600 MHz, 4:1 $\text{CDCl}_3/\text{CD}_3\text{OD}$) δ 8.31 (m, 1H), 7.93 (d, 1H, J = 7.9 Hz), 7.87 (d, 1H, J = 8.4 Hz), 7.47 (t, 1H, J = 7.7 Hz), 7.44 (d, 1H, J = 8.8 Hz), 7.30 (t, 1H, J = 7.4 Hz), 7.09 (q, 1H, J = 9.3 Hz), 2.91 (t, 1H, J = 11.8 Hz), 2.13 (d, 2H, J = 11.9 Hz), 1.89 (d, 2H, J = 13.2 Hz), 1.78 (d, 1H, J = 12.7 Hz), 1.66 (q, 2H, J = 11.4 Hz), 1.44 (q, 2H, J = 12.8 Hz), 1.32 (q, 1H, J = 13.0 Hz); ^{13}C NMR (150 MHz, 4:1 $\text{CDCl}_3/\text{CD}_3\text{OD}$) δ 175.5, 163.8, 158.0, 147.1, 143.9, 134.7, 127.6, 123.2, 120.8, 117.2, 117.1, 115.7, 114.4, 108.3, 108.1, 38.8, 32.2, 26.4, 26.1; HRMS (ESI) calcd for $\text{C}_{22}\text{H}_{21}\text{F}_2\text{N}_4$ ($\text{M} + \text{H}^+$) 379.1734, found 379.1739.

2-Cyclohexyl-*N*-(3,5-difluorophenyl)-1*H*-imidazo[4,5-*c*]quinolin-4-amine (9). Compound **29** (5.1 mg, 18 μ mol) was reacted with 3,5-difluoroaniline (9.5 mg, 72 μ mol) in DMF (50 μ L). The crude product was chromatographed on silica gel (20:1 $\text{CH}_2\text{Cl}_2/\text{MeOH}$, 20:1 $\text{CHCl}_3/\text{MeOH}$, and then 15:1 $\text{CH}_2\text{Cl}_2/\text{MeOH}$) to give 7.4 mg (20 μ mol, 100%) of **9**. R_f = 0.54 [silica gel, 20:1 $\text{CH}_2\text{Cl}_2/\text{MeOH}$]; ^1H NMR (600 MHz, $\text{DMSO}-d_6$) δ 13.19 (s, 1H), 9.58 (s, 1H), 8.16 (m, 3H), 7.83 (d, 1H, J = 8.2 Hz), 7.53 (t, 1H, J = 7.7 Hz), 7.41 (t, 1H, J = 8.0 Hz), 6.73 (t, 1H, J = 8.8 Hz), 3.00 (tt, 1H, J = 11.7, 3.3 Hz), 2.08 (d, 2H, J = 12.2 Hz), 1.87 (d, 2H, J = 13.1 Hz), 1.77–1.71 (m, 3H), 1.44 (qt, 2H, J = 12.8, 3.0 Hz), 1.32 (qt, 1H, J = 12.5, 3.3 Hz); ^{13}C NMR (150 MHz, 4:1 $\text{CDCl}_3/\text{CD}_3\text{OD}$) δ 164.6, 164.4, 162.9, 162.8, 146.8, 143.7, 143.6, 134.8, 127.8, 127.7, 123.5, 120.8, 115.8, 101.6, 101.4, 96.8, 96.6, 96.5, 38.8, 32.2, 26.4, 26.1; HRMS (ESI) calcd for $\text{C}_{22}\text{H}_{21}\text{F}_2\text{N}_4$ ($\text{M} + \text{H}^+$) 379.1734, found 379.1740.

***N*-(3,5-Bis(trifluoromethyl)phenyl)-2-cyclohexyl-1*H*-imidazo[4,5-*c*]quinolin-4-amine (10).** Compound **29** (4.35 mg, 15.2 μ mol) was reacted with 3,5-bis(trifluoromethyl)aniline (10.0 μ L, 62.7 μ mol) in DMF (50 μ L). The crude product was chromatographed on silica gel (2:1 hexane/EtOAc, 30:1 $\text{CHCl}_3/\text{MeOH}$, 80:20:1 hexane/EtOAc/triethylamine, 5:2 hexane/EtOAc, and 35:1 $\text{CHCl}_3/\text{MeOH}$ with a trace amount of citric acid) to give 1.40 mg (2.93 μ mol, 19%) of **10**. R_f = 0.43 [silica gel, 2:1 hexane/EtOAc]; ^1H NMR (600 MHz,

$\text{DMSO}-d_6$) δ 13.25 (s, 1H), 10.00 (s, 1H), 9.17 (s, 2H), 8.19 (d, 1H, J = 7.7 Hz), 7.76 (d, 1H, J = 8.2 Hz), 7.58 (s, 1H), 7.56 (t, 1H, J = 7.9 Hz), 7.44 (t, 1H, J = 7.6 Hz), 3.02 (tt, 1H, J = 11.8, 3.4 Hz), 2.11 (d, 2H, J = 11.9 Hz), 1.88 (d, 2H, J = 13.5 Hz), 1.78–1.72 (m, 3H), 1.45 (qt, 2H, J = 12.8, 3.3 Hz), 1.32 (qt, 1H, J = 12.6, 3.3 Hz); ^{13}C NMR (150 MHz, 4:1 $\text{CDCl}_3/\text{CD}_3\text{OD}$) δ 143.6, 142.9, 135.0, 128.0, 127.8, 123.8, 120.8, 118.2, 115.9, 114.3, 38.8, 32.2, 26.4, 26.1; HRMS (ESI) calcd for $\text{C}_{24}\text{H}_{21}\text{F}_6\text{N}_4$ ($\text{M} + \text{H}^+$) 479.1670, found 479.1680.

4-(2-Cyclohexyl-1*H*-imidazo[4,5-*c*]quinolin-4-ylamino)phthalonitrile (11). Compound **29** (4.50 mg, 15.7 μ mol) was reacted with 4-aminophthalonitrile (11.6 mg, 78.8 μ mol) in DMF (50 μ L). The crude product was chromatographed on reversed-phase C18 silica gel (9:1 MeOH/ H_2O) and normal-phase silica gel (15:1 $\text{CHCl}_3/\text{MeOH}$) to give 0.45 mg (1.1 μ mol, 7.3%) of **11**. R_f = 0.80 [silica gel, 10:1 $\text{CHCl}_3/\text{MeOH}$]; ^1H NMR (600 MHz, $\text{DMSO}-d_6$) δ 13.30 (s, 1H), 10.19 (s, 1H), 9.07 (s, 1H), 8.78 (d, 1H, J = 7.6 Hz), 8.21 (d, 1H, J = 7.2 Hz), 8.04 (d, 1H, J = 8.9 Hz), 7.89 (d, 1H, J = 8.2 Hz), 7.58 (t, 1H, J = 7.6 Hz), 7.48 (t, 1H, J = 7.1 Hz), 3.02 (tt, 1H, J = 12.1, 3.4 Hz), 2.09 (d, 2H, J = 10.7 Hz), 1.87 (d, 2H, J = 13.2 Hz), 1.78–1.72 (m, 3H), 1.45 (qt, 2H, J = 12.8, 3.3 Hz), 1.32 (qt, 1H, J = 12.9, 3.5 Hz); ^{13}C NMR (150 MHz, 4:1 $\text{CDCl}_3/\text{CD}_3\text{OD}$) δ 158.5, 146.2, 145.7, 143.3, 134.6, 130.0, 128.2, 128.0, 124.4, 122.6, 121.8, 120.9, 116.9, 116.7, 116.5, 116.1, 38.8, 32.2, 26.4, 26.1; HRMS (ESI) calcd for $\text{C}_{24}\text{H}_{21}\text{N}_6$ ($\text{M} + \text{H}^+$) 393.1828, found 393.1835.

2-Cyclohexyl-*N*-(3,4-dimethoxyphenyl)-1*H*-imidazo[4,5-*c*]quinolin-4-amine (12). Compound **29** (7.8 mg, 27 μ mol) was reacted with 3,4-dimethoxyaniline (13 mg, 83 μ mol) in DMF (50 μ L). The crude product was chromatographed on silica gel (2:3 hexane/EtOAc) to give 2.5 mg (6.2 μ mol, 23%) of **12**. R_f = 0.33 [silica gel, 1:1 petroleum ether/EtOAc]; ^1H NMR (600 MHz, 4:1 $\text{CDCl}_3/\text{CD}_3\text{OD}$) δ 7.98 (s, 1H), 7.92 (d, 1H, J = 6.6 Hz), 7.80 (d, 1H, J = 8.6 Hz), 7.45 (t, 1H, J = 7.6 Hz), 7.33 (d, 1H, J = 9.7 Hz), 7.27 (t, 1H, J = 7.1 Hz), 6.87 (d, 1H, J = 8.1 Hz), 3.95 (s, 3H), 3.84 (s, 3H), 2.92 (t, 1H, J = 11.9 Hz), 2.13 (d, 2H, J = 11.0 Hz), 1.89 (d, 2H, J = 13.2 Hz), 1.78 (d, 1H, J = 11.6 Hz), 1.67 (q, 2H, J = 12.1 Hz), 1.44 (q, 2H, J = 13.0 Hz), 1.32 (q, 1H, J = 13.0 Hz); ^{13}C NMR (150 MHz, 4:1 $\text{CDCl}_3/\text{CD}_3\text{OD}$) δ 149.3, 144.3, 144.1, 135.1, 127.5, 127.3, 122.7, 120.7, 112.4, 111.0, 104.5, 56.6, 56.0, 38.9, 32.2, 26.4, 26.1; HRMS (ESI) calcd for $\text{C}_{24}\text{H}_{27}\text{N}_4\text{O}_2$ ($\text{M} + \text{H}^+$) 403.2134, found 403.2100.

***N*-(Benzo[*d*][1,3]dioxol-5-yl)-2-cyclohexyl-1*H*-imidazo[4,5-*c*]quinolin-4-amine (13).** Compound **29** (13.7 mg, 47.9 μ mol) was reacted with 3,4-(methylenedioxy)aniline (16.3 mg, 115 μ mol) in DMF (100 μ L). The crude product was chromatographed on silica gel (2:3 and 1:1 hexane/EtOAc) to give 5.4 mg (14 μ mol, 29%) of **13**. R_f = 0.59 [silica gel, 1:1 hexane/EtOAc]; ^1H NMR (600 MHz, 4:1 $\text{CDCl}_3/\text{CD}_3\text{OD}$) δ 7.91 (br s, 1H), 7.82 (d, 1H, J = 8.6 Hz), 7.80 (s, 1H), 7.44 (t, 1H, J = 7.7 Hz), 7.26 (t, 1H, J = 6.8 Hz), 7.21 (d, 1H, J = 7.3 Hz), 6.78 (d, 1H, J = 8.5 Hz), 5.91 (s, 2H), 2.91 (t, 1H, J = 11.6 Hz), 2.12 (d, 2H, J = 12.0 Hz), 1.89 (d, 2H, J = 13.5 Hz), 1.77 (d, 1H, J = 13.4 Hz), 1.66 (q, 2H, J = 11.9 Hz), 1.43 (q, 2H, J = 12.7 Hz), 1.31 (q, 1H, J = 12.8 Hz); ^{13}C NMR (150 MHz, 4:1 $\text{CDCl}_3/\text{CD}_3\text{OD}$) δ 147.9, 144.0, 135.6, 127.5, 127.2, 122.8, 120.7, 115.6, 112.1, 108.5, 102.2, 101.2, 38.9, 32.2, 26.4, 26.1; HRMS (ESI) calcd for $\text{C}_{23}\text{H}_{23}\text{N}_4\text{O}_2$ ($\text{M} + \text{H}^+$) 387.1821, found 387.1814.

2-Cyclohexyl-*N*-(pyridin-3-yl)-1*H*-imidazo[4,5-*c*]quinolin-4-amine (14). Compound **29** (6.67 mg, 23.3 μ mol) was reacted with 3-aminopyridine (11.0 mg, 116 μ mol) in DMF (50 μ L). The crude product was chromatographed on normal-phase silica gel (10:1 and 3:1 $\text{CHCl}_3/\text{MeOH}$ and 90:3:1 $\text{CHCl}_3/\text{MeOH}$ /triethylamine), on reversed-phase C18 silica gel (4:1 MeOH/ H_2O), and again on normal-phase silica gel (15:1 $\text{CHCl}_3/\text{MeOH}$) to give 0.35 mg (0.99 μ mol, 4.2%) of **14**. R_f = 0.57 [silica gel, 200:10:1 $\text{CHCl}_3/\text{MeOH}$ /triethylamine]; ^1H NMR (600 MHz, $\text{DMSO}-d_6$) [major tautomer] δ 13.16 (s, 1H), 9.36 (d, 1H, J = 2.6 Hz), 9.31 (s, 1H), 8.73 (ddd, 1H, J = 8.2, 2.5, 1.3 Hz), 8.17 (dd, 1H, J = 4.7, 1.5 Hz), 8.14 (dd, 1H, J = 8.2, 1.0 Hz), 7.78 (d, 1H, J = 7.7 Hz), 7.50 (td, 1H, J = 7.7, 1.7 Hz), 7.38 (td, 1H, J = 7.4, 1.1 Hz), 7.35 (dd, 1H, J = 8.3,

4.7 Hz), 3.00 (tt, 1H, $J = 12.1, 3.5$ Hz), 2.09 (d, 2H, $J = 12.7$ Hz), 1.87 (dt, 2H, $J = 13.2, 3.4$ Hz), 1.78–1.72 (m, 3H), 1.45 (qt, 2H, $J = 13.0, 3.2$ Hz), 1.32 (qt, 1H, $J = 12.8, 3.3$ Hz); ^{13}C NMR (150 MHz, 2:1 $\text{CDCl}_3/\text{CD}_3\text{OD}$) δ 158.4, 142.0, 140.3, 139.0, 137.2, 131.5, 130.4, 130.2, 127.9, 127.8, 126.7, 124.5, 123.7, 121.1, 116.1, 39.1, 32.4, 26.6, 26.3; HRMS (ESI) calcd for $\text{C}_{21}\text{H}_{22}\text{N}_5$ ($\text{M} + \text{H}^+$) 344.1875, found 344.1873.

General Procedure for N-Substituted 1H-Imidazo[4,5-c]quinolin-4-amine (16–21, 23). A solution of the appropriate 4-chloro-1H-imidazo[4,5-c]quinoline and 3,4-dichloroaniline in DMF was heated at 140 °C overnight under a dry nitrogen atmosphere. The reaction mixture was cooled, and the solvent was evaporated. The residue obtained was subjected to preparative silica gel column chromatography (100:1 to 8:1 $\text{CHCl}_3/\text{MeOH}$).

N-(3,4-Dichlorophenyl)-2-(tetrahydro-2H-pyran-4-yl)-1H-imidazo[4,5-c]quinolin-4-amine (16). Scale: 0.14 mmol. Yield: 20 mg (35%). ^1H NMR (300 MHz, $\text{CD}_3\text{OD}/\text{CDCl}_3$) δ 8.44 (s, 1H), 7.89 (d, 2H, $J = 9.0$ Hz), 7.74 (dd, 1H, $J = 9.0, 2.5$ Hz), 7.50 (m, 1H), 7.37 (m, 2H), 4.10 (tdt, 2H, $J = 11.4, 3.3$ Hz), 3.59 (m, 2H), 3.21 (m, 1H), 2.04 (m, 4H); ^{13}C NMR (75 MHz, $\text{CD}_3\text{OD}/\text{CDCl}_3$) δ 143.4, 140.4, 132.2, 127.4, 127.2, 124.0, 123.2, 120.4, 119.9, 117.9, 115.3, 67.5, 35.5, 31.1; HRMS (ESI) calcd for $\text{C}_{21}\text{H}_{19}\text{Cl}_2\text{N}_5\text{O}^+$ ($\text{M} + \text{H}^+$) 413.0930, found 413.0952.

N-(3,4-Dichlorophenyl)-2-(exonorbornanyl)-1H-imidazo[4,5-c]quinolin-4-amine (17). Scale: 0.139 mmol. Yield: 6.4 mg (11%). ^1H NMR (300 MHz, $\text{CD}_3\text{OD}/\text{CDCl}_3$) δ 9.54 (s, 1H), 8.49 (s, 1H), 8.00 (d, 1H, $J = 8.1$ Hz), 7.83 (t, 1H, $J = 9.6$ Hz), 7.57 (t, 2H, $J = 8.4$ Hz), 7.31 (m, 1H), 7.27 (m, 12H), 3.99 (m, 1H), 3.59 (s, 1H), 3.42 (dd, 1H, $J = 11.4, 6.0$ Hz), 2.71 (s, 1H), 2.47 (s, 1H), 2.09 (m, 2H), 1.47 (m, 5H); ^{13}C NMR (75 MHz, $\text{CD}_3\text{OD}/\text{CDCl}_3$) δ 130.4, 127.8, 123.4, 120.6, 120.0, 118.4, 42.3, 41.2, 40.5, 37.3, 33.5, 29.9, 24.2; HRMS (ESI) calcd for $\text{C}_{23}\text{H}_{21}\text{Cl}_2\text{N}_4^+$ ($\text{M} + \text{H}^+$) 423.1138, found 423.1140.

N-(3,4-Dichlorophenyl)-2-(endonorbornanyl)-1H-imidazo[4,5-c]quinolin-4-amine (18). Scale: 0.139 mmol. Yield: 12.6 mg (21%). ^1H NMR (300 MHz, $\text{CD}_3\text{OD}/\text{CDCl}_3$) δ 9.81 (s, 1H), 8.44 (s, 1H), 7.98 (d, 1H, $J = 8.4$ Hz), 7.78 (m, 2H), 7.55 (t, 1H, $J = 8.5$ Hz), 7.36 (s, 1H), 2.99 (dd, 1H, $J = 8.7, 4.9$ Hz), 2.60 (s, 1H), 2.48 (s, 1H), 2.23 (m, 1H), 1.82 (m, 1H), 1.66 (m, 2H), 1.30 (m, 5H); ^{13}C NMR (75 MHz, $\text{CD}_3\text{OD}/\text{CDCl}_3$) δ 130.2, 127.4, 123.1, 120.1, 118.1, 42.7, 41.9, 36.4, 36.3, 36.1, 29.7, 28.9; HRMS (ESI) calcd for $\text{C}_{23}\text{H}_{21}\text{Cl}_2\text{N}_4^+$ ($\text{M} + \text{H}^+$) 423.1138, found 423.1151.

N-(3,4-Dichlorophenyl)-2-(3-noradamantanyl)-1H-imidazo[4,5-c]quinolin-4-amine (19). Scale: 0.186 mmol. Yield: 58.2 mg (70%). ^1H NMR (300 MHz, $\text{CD}_3\text{OD}/\text{CDCl}_3$) δ 8.23 (s, 1H), 8.06 (d, 1H, $J = 8.4$ Hz), 7.80 (d, 1H, $J = 8.4$ Hz), 7.63 (dd, 1H, $J = 8.7, 2.4$ Hz), 7.43 (m, 1H), 7.31 (m, 2H), 2.76 (t, 1H, $J = 6.3$ Hz), 2.38 (s, 2H), 2.20 (d, 2H, $J = 10.5$ Hz), 2.08 (m, 2H), 2.02 (m, 4H), 1.70 (m, 4H); ^{13}C NMR (75 MHz, $\text{CD}_3\text{OD}/\text{CDCl}_3$) δ 160.5, 145.9, 142.0, 139.9, 139.8, 132.4, 132.4, 130.3, 127.6, 125.9, 124.9, 123.5, 120.9, 118.8, 115.5, 45.8, 43.8, 37.6, 34.5; HRMS (ESI) calcd for $\text{C}_{25}\text{H}_{23}\text{Cl}_2\text{N}_4^+$ ($\text{M} + \text{H}^+$) 449.1294, found 449.1291.

N-(3,4-Dichlorophenyl)-2-(1-adamantanyl)-1H-imidazo[4,5-c]quinolin-4-amine (20). Scale: 0.115 mmol. Yield: 38 mg (71%). ^1H NMR (300 MHz, $\text{CD}_3\text{OD}/\text{CDCl}_3$) δ 8.33 (s, 1H), 8.04 (d, 1H, $J = 7.5$ Hz), 7.81 (d, 1H, $J = 8.4$ Hz), 7.66 (m, 1H), 7.42 (m, 1H), 7.31 (m, 3H), 2.09 (s, 6H), 1.78 (s, 4H); ^{13}C NMR (75 MHz, $\text{CD}_3\text{OD}/\text{CDCl}_3$) δ 161.3, 146.1, 142.4, 140.1, 132.3, 124.5, 120.6, 41.0, 36.3, 35.5, 28.6, 27.6; HRMS (ESI) calcd for $\text{C}_{26}\text{H}_{25}\text{Cl}_2\text{N}_4^+$ ($\text{M} + \text{H}^+$) 463.1451, found 463.1456.

N-(3,4-Dichlorophenyl)-2-((1s,4s)-4-(1,3-dioxoisindolin-2-yl)-cyclohexyl)-1H-imidazo[4,5-c]quinolin-4-amine (21). Scale: 0.044 mmol. Yield: 40 mg (78%). ^1H NMR (300 MHz, $\text{CD}_3\text{OD}/\text{CDCl}_3$) δ 8.47 (m, 1H), 8.23 (s, 1H), 7.78 (m, 7H), 7.39 (m, 14H), 4.28 (m, 1H), 3.09 (m, 1H), 2.52 (m, 4H), 1.91 (m, 4H); ^{13}C NMR (75 MHz, $\text{CD}_3\text{OD}/\text{CDCl}_3$) δ 168.5, 160.1, 143.3, 140.4, 134.0, 132.0, 131.6, 130.2, 129.9, 127.2, 122.9, 121.3, 120.3, 119.8, 118.9, 117.8, 100.2, 37.0, 30.9, 28.9; HRMS (ESI) calcd for $\text{C}_{30}\text{H}_{24}\text{Cl}_2\text{N}_5\text{O}_2^+$ ($\text{M} + \text{H}^+$) 556.1302, found 556.1290.

N-(3,4-Dichlorophenyl)-2-(1-benzoylpiperidin-4-yl)-1H-imidazo[4,5-c]quinolin-4-amine (23). Scale: 0.108 mmol. Yield: 25 mg (44%). ^1H NMR (300 MHz, $\text{CD}_3\text{OD}/\text{CDCl}_3$) δ 8.37 (s, 1H), 7.85 (d, 1H, $J = 7.3$ Hz), 7.82 (d, 1H, $J = 8.4$ Hz), 7.66 (dd, 1H, $J = 8.7, 2.4$ Hz), 7.43 (m, 1H), 7.31 (m, 7H), 4.69 (m, 12H), 3.82 (m, 2H), 3.17 (m, 1H), 3.00 (s, 1H), 1.92 (m, 4H); ^{13}C NMR (75 MHz, $\text{CD}_3\text{OD}/\text{CDCl}_3$) δ 171.2, 143.5, 140.4, 135.3, 132.2, 130.1, 128.6, 127.5, 127.3, 126.5, 124.0, 123.2, 120.4, 119.9, 117.9, 42.1, 36.4, 31.0, 30.5; HRMS (ESI) calcd for $\text{C}_{28}\text{H}_{24}\text{Cl}_2\text{N}_5\text{O}^+$ ($\text{M} + \text{H}^+$) 516.1352, found 516.1352.

N-Phenyl-2-(1-benzoylpiperidin-4-yl)-1H-imidazo[4,5-c]quinolin-4-amine (24). A solution of **33d** (61.4 mg, 0.156 mmol) and aniline (43 μL , 0.47 mmol) in DMF (0.8 mL) was heated at 140 °C overnight under N_2 atmosphere. The reaction mixture was cooled, and the solvent was evaporated. The residue obtained was subjected to preparative silica gel column chromatography (100:1 to 8:1 $\text{CHCl}_3/\text{MeOH}$). Yield: 40 mg (57%). ^1H NMR (300 MHz, $\text{CD}_3\text{OD}/\text{CDCl}_3$) δ 7.92 (d, 1H, $J = 7.5$ Hz), 7.81 (s, 2H), 7.75 (d, 1H, $J = 8.1$ Hz), 7.33 (dd, 1H, $J = 7.2, 1.5$ Hz), 7.21 (t, 1H, $J = 8.0$ Hz), 7.12 (d, 1H, $J = 7.3$ Hz), 6.86 (t, 1H, $J = 7.5$ Hz), 4.62 (m, 12H), 3.60 (m, 2H), 3.07 (d, 1H, $J = 11.4$ Hz), 1.92 (m, 4H); ^{13}C NMR (75 MHz, $\text{CD}_3\text{OD}/\text{CDCl}_3$) δ 170.7, 154.8, 143.6, 140.6, 135.6, 129.8, 128.7, 128.5, 127.2, 127.1, 126.6, 122.5, 121.6, 120.5, 118.7, 53.4, 20.8, 13.9; HRMS (ESI) calcd for $\text{C}_{28}\text{H}_{26}\text{N}_5\text{O}^+$ ($\text{M} + \text{H}^+$) 448.2132, found 448.2137.

2-((1s,4s)-4-Aminocyclohexyl)-N-(3,4-dichlorophenyl)-1H-imidazo[4,5-c]quinolin-4-amine (22). A 33% solution of methylamine in absolute ethanol (0.5 mL) is added to a stirred solution of **21** (10 mg, 0.017 mmol) in ethanol (0.25 mL). The solution was refluxed for 2 h. The mixture was then cooled to room temperature, and the solvent was evaporated under reduced pressure. The residue was purified by flash column chromatography using a mixture of 10:1 EtOAc/MeOH, by volume. Yield: 5 mg (70%). ^1H NMR (300 MHz, $\text{CD}_3\text{OD}/\text{CDCl}_3$) δ 8.40 (s, 1H), 8.08 (d, 1H, $J = 8.1$ Hz), 7.87 (d, 1H, 8.7 Hz), 7.75 (dd, 1H, $J = 8.7, 2.5$ Hz), 7.47 (t, 2H, $J = 6.9$ Hz), 7.29 (m, 4H), 3.40 (m, 1H), 3.12 (m, 1H), 2.27 (m, 2H), 1.95 (m, 2H), 1.89 (m, 4H); ^{13}C NMR (75 MHz, $\text{CD}_3\text{OD}/\text{CDCl}_3$) δ 143.4, 140.4, 132.1, 130.0, 127.4, 124.0, 123.2, 120.8, 119.9, 117.9, 110.7, 105.7, 31.6, 27.4, 25.6; HRMS (ESI) calcd for $\text{C}_{22}\text{H}_{22}\text{Cl}_2\text{N}_5^+$ ($\text{M} + \text{H}^+$) 426.1247, found 426.4203.

2-(1-Benzylpiperidin-4-yl)-N-(3,4-dichlorophenyl)-1H-imidazo[4,5-c]quinolin-4-amine (25). A solution of compound **23** (18 mg, 0.035 mmol) in dry THF (0.18 mL) was added dropwise to a stirred slurry of lithium aluminum hydride (0.89 mL of 0.2 M in THF, 0.175 mmol). After the mixture was heated under reflux overnight, decomposition of excess hydride was effected by cautious addition of water. The inorganic solids were removed by filtration, the organic layer was dried over anhydrous sodium sulfate and filtered, and the solvent was removed under reduced pressure. The final residue was purified by flash column chromatography using a mixture of 10:1 EtOAc/MeOH, by volume. ^1H NMR (300 MHz, $\text{CD}_3\text{OD}/\text{CDCl}_3$) δ 11.79 (s, 1H), 8.50 (s, 1H), 8.01 (m, 2H), 7.87 (dd, 1H, $J = 9.0, 2.6$ Hz), 7.78 (m, 1H), 7.53 (m, 7H), 7.37 (s, 2H), 3.59 (m, 2H), 3.09 (m, 2H), 2.06 (m, 2H), 1.76 (s, 4H); ^{13}C NMR (75 MHz, $\text{CD}_3\text{OD}/\text{CDCl}_3$) δ 140.4, 136.4, 129.4, 128.1, 127.8, 123.0, 120.3, 119.8, 117.8, 100.2, 61.6, 52.9, 30.1, 28.8; HRMS (ESI) calcd for $\text{C}_{28}\text{H}_{26}\text{Cl}_2\text{N}_5^+$ ($\text{M} + \text{H}^+$) 502.1560, found 502.1541.

N-(3,4-Dichlorophenyl)-2-(piperidin-4-yl)-1H-imidazo[4,5-c]quinolin-4-amine (26). A solution of **25** (6 mg, 0.011 mmol) in dry MeOH (0.5 mL) was hydrogenated at 1 atm in the presence of palladium black (0.011 mg). After 2 h, the catalyst was removed by filtration, and the solvent was evaporated under reduced pressure. The final residue was purified by flash column chromatography using a mixture of 10:1 EtOAc/MeOH/ NH_4OH , by volume. ^1H NMR (300 MHz, $\text{CD}_3\text{OD}/\text{CDCl}_3$) δ 8.51 (m, 1H), 8.03 (m, 2H), 7.72 (m, 1H), 7.60 (m, 1H), 7.43 (m, 2H), 3.38 (m, 2H), 2.35 (m, 2H), 1.19 (m, 12H), 0.90 (m, 12H); HRMS (ESI) calcd for $\text{C}_{21}\text{H}_{20}\text{Cl}_2\text{N}_5^+$ ($\text{M} + \text{H}^+$) 412.1090, found 412.1086.

HPLC Analysis of Compounds 5–14 and 16–26. Purity of compounds was checked using a Hewlett-Packard 1100 HPLC

equipped with a Zorbax SB-Aq 5 μ m analytical column (50 mm \times 4.6 mm; Agilent Technologies, Palo Alto, CA). System A: linear gradient solvent system; 5 mM TBAP (tetrabutylammonium dihydrogenphosphate)—CH₃CN from 50:50 to 0:100 in 13 min; the flow rate was 0.5 mL/min. System B: linear gradient solvent system; 10 mM TEAA (triethylammonium acetate)—CH₃CN from 65:35 to 0:100 in 13 min; the flow rate was 0.5 mL/min. Peaks were detected by UV absorption with a diode array detector. The compounds eluted at the following retention times: **5**, 7.0 min (system A), 9.4 min (system B); **6**, 8.9 min (system A), 10.44 min (system B); **7**, 9.2 min (system A), 10.9 min (system B); **8**, 7.5 min (system A), 9.3 min (system B); **9**, 7.3 min (system A), 10.1 min (system B); **10**, 9.4 min (system A), 11.7 min (system B); **11**, 7.1 min (system A), 9.8 min (system B); **12**, 3.3 min (system A), 8.0 min (system B); **13**, 4.7 min (system A), 8.6 min (system B); **14**, 7.2 min (system A), 4.7 min (system B); **16**, 5.7 min (system A), 8.6 min (system B); **17**, 8.4 min (system A), 10.9 min (system B); **18**, 8.7 min (system A), 11.2 min (system B); **19**, 9.3 min (system A), 11.3 min (system B); **20**, 9.7 min (system A), 12.0 min (system B); **21**, 9.2 min (system A), 11.5 min (system B); **22**, 2.4 min (system A), 10.3 min (system B); **23**, 6.5 min (system A), 9.6 min (system B); **24**, 2.9 min (system A), 6.9 min (system B); **25**, 7.06 min (system A), 10.8 min (system B); **26**, 1.93 min (system A), 10.6 min (system B).

Pharmacological Methods. [³H]**35** (47 Ci/mmol) was from GE Healthcare Bio-Sciences Corp. (Piscataway, NJ). [³H]**34** (CCPA, 42.6 Ci/mmol), [¹²⁵I]**36** (I-AB-MECA, 2000 Ci/mmol), and [³⁵S]GTP γ S (1068 Ci/mmol) were from Perkin-Elmer Life Sciences (Waltham, MA).

Cell Culture and Membrane Preparation. CHO (Chinese hamster ovary) cells expressing the recombinant human ARs (HEK-293 cells were used for the human A_{2A}AR) were cultured in DMEM and F12 (1:1) supplemented with 10% fetal bovine serum, 100 units/mL penicillin, 100 μ g/mL streptomycin, and 2 μ mol/mL glutamine. Cells were harvested by trypsinization. After homogenization and suspension, cells were centrifuged at 1000g for 10 min, and the pellet was resuspended in 50 mM Tris-HCl buffer (pH 7.4) containing 10 mM MgCl₂. The suspension was homogenized with an electric homogenizer for 10 s and was then recentrifuged at 20000g for 20 min at 4 °C. The resultant pellets were resuspended in buffer in the presence of 3 U/mL adenosine deaminase, and the suspension was stored at -80 °C until the binding experiments. The protein concentration was measured using the Bradford assay.²⁷

Binding to the Human A₁AR and the A_{2A}AR. For binding to the human A₁AR, [³H]**34** (2 nM) was incubated with membranes (40 μ g/tube) from CHO cells stably expressing the human A₁AR at 2.4 min for 60 min in 50 mM Tris-HCl buffer (pH 7.4; MgCl₂, 10 mM) in a total assay volume of 200 μ L.¹⁹ Nonspecific binding was determined using 10 μ M of N⁶-cyclopentyladenosine. For human A_{2A}AR binding, membranes (20 μ g/tube) from HEK-293 cells stably expressing the human A_{2A}AR were incubated with 15 nM [³H]**35** at 25 °C for 60 min in 200 μ L of 50 mM Tris-HCl, pH 7.4, containing 10 mM MgCl₂.²⁰ 5'-N-Ethylcarboxamidoadenosine (NECA, **37**, 10 μ M) was used to define nonspecific binding. Reaction was terminated by filtration with GF/B filters.

Binding to the Human A₃AR. Each tube in the competitive binding assay contained 100 μ L of membrane suspension (20 μ g protein), 50 μ L of [¹²⁵I]**36** (0.5 nM),²¹ and 50 μ L of increasing concentrations of the test ligands in Tris-HCl buffer (50 mM, pH 7.4) containing 10 mM MgCl₂, 1 mM EDTA. Nonspecific binding was determined using 10 μ M of **37** (NECA) in the buffer. The mixtures were incubated at 25 °C for 60 min. Dissociation was started by the addition of 10 μ M **15** in the absence or presence of 10 μ M of each allosteric modulator. Binding reactions were terminated by filtration through Whatman GF/B filters under reduced pressure using a MT-24 cell harvester (Brandell, Gaithersburg, MD). Filters were washed three times with 9 mL of ice-cold buffer. Radioactivity was determined in a Beckman 5500B γ -counter.

[³⁵S]GTP γ S Binding Assay. The preparation of membranes from CHO cells stably expressing human A₃AR was as previously

described.²² [³⁵S]GTP γ S binding was measured in 200 μ L of buffer containing 50 mM Tris-HCl (pH 7.4), 1 mM EDTA, 1 mM MgCl₂, 1 μ M GDP, 1 mM dithiothreitol, 100 mM NaCl, 3 U/mL adenosine deaminase, 0.2 nM [³⁵S]GTP γ S, 0.004% 3-[(3-cholamidopropyl)-dimethylammonio]-1-propanesulfonate (CHAPS), and 0.5% bovine serum albumin. Incubations were started by addition of the membrane suspension (10 μ g of protein/tube) to the test tubes and carried out in duplicate for 30 min at 25 °C. The reaction was stopped by rapid filtration through Whatman GF/B filters, presoaked in 50 mM Tris-HCl and 5 mM MgCl₂ (pH 7.4) containing 0.02% CHAPS. The filters were washed twice with 3 mL of the buffer mentioned before, and retained radioactivity was measured using liquid scintillation counting. Nonspecific binding of [³⁵S]GTP γ S was measured in the presence of 10 μ M unlabeled GTP γ S.

Statistical Analysis. Binding and functional parameters were calculated using Prism 5.0 software (GraphPAD, San Diego, CA). IC₅₀ values obtained from competition curves were converted to K_i values using the Cheng-Prusoff equation.²⁸ Data were expressed as the mean \pm standard error.

Acknowledgment. This research was supported in part by the Intramural Research Program of the NIH, National Institute of Diabetes and Digestive and Kidney Diseases, and by the European Union (Grant QLK3-CT-2001-51963). We thank Dr. Anikó Göblyös for preparation of compound **27**, Dr. John Lloyd for collecting MS data, and Dr. Herman Yeh and Wesley White for the helpful advice on the NMR experiments. S.d.C. thanks Ministerio de Educación y Ciencia (Spain) for financial support. Y.K. thanks the Can-Fite Biopharma for financial support.

Supporting Information Available: Selected ¹H and 2D COSY and NOESY spectra with peak assignments. This material is available free of charge via the Internet at <http://pubs.acs.org>.

References

- (1) Fredholm, B. B.; IJzerman, A. P.; Jacobson, K. A.; Klotz, K. N.; Linden, J. International Union of Pharmacology. XXV. Nomenclature and classification of adenosine receptors. *Pharmacol. Rev.* **2001**, *53*, 527–552.
- (2) Jacobson, K. A.; Gao, Z.-G. Adenosine receptors as therapeutic targets. *Nat. Rev. Drug Discovery* **2006**, *5*, 247–264.
- (3) Wan, T. C.; Ge, Z. D.; Tampo, A.; Mio, Y.; Bienengraeber, M. W.; Tracey, W. R.; Gross, G. J.; Kwok, W. M.; Auchampach, J. A. The A₃ adenosine receptor agonist CP-532,903 [N⁶-(2,5-dichlorobenzyl)-3'-aminoadenosine-5'-N-methylcarboxamide] protects against myocardial ischemia/reperfusion injury via the sarcolemmal ATP-sensitive potassium channel. *J. Pharmacol. Exp. Ther.* **2008**, *324*, 234–243.
- (4) Guzman, J.; Yu, J. G.; Suintres, Z.; Bozarov, A.; Cooke, H.; Javed, N.; Auer, H.; Palatini, J.; Hassanain, H. H.; Cardounel, A. J.; Javed, A.; Grants, I.; Wunderlich, J. E.; Christofi, F. L. ADOA3R as a therapeutic target in experimental colitis: proof by validated high-density oligonucleotide microarray analysis. *Inflammatory Bowel Dis.* **2006**, *12*, 766–789.
- (5) Fishman, P.; Jacobson, K. A.; Ochaion, A.; Cohen, S.; Bar-Yehuda, S. The anti-cancer effect of A₃ adenosine receptor agonists: a novel, targeted therapy. *Immunol., Endocr. Metab. Agents Med. Chem.* **2007**, *7*, 298–303.
- (6) Silverman, M. H.; Strand, V.; Markovits, D.; Nahir, M.; Reitblat, T.; Molad, Y.; Rosner, I.; Rozenbaum, M.; Mader, R.; Adawi, M.; Caspi, D.; Tishler, M.; Langevitz, P.; Rubinow, A.; Friedman, J.; Green, L.; Tanay, A.; Ochaion, A.; Cohen, S.; Kerns, W. D.; Cohn, I.; Fishman-Furman, S.; Farbstein, M.; Yehuda, S. B.; Fishman, P. Clinical evidence for utilization of the A₃ adenosine receptor as a target to treat rheumatoid arthritis: data from a phase II clinical trial. *J. Rheumatol.* **2008**, *35*, 41–48.
- (7) Zheng, J.; Wang, R.; Zambraski, E.; Wu, D.; Jacobson, K. A.; Liang, B. T. A novel protective action of adenosine A₃ receptors: attenuation of skeletal muscle ischemia and reperfusion injury. *Am. J. Physiol.: Heart Circ. Physiol.* **2007**, *293*, 3685–3691.
- (8) Bruns, R. F.; Fergus, J. H. Allosteric enhancement of adenosine A₁ receptor binding and function by 2-amino-3-benzoylthiophenes. *Mol. Pharmacol.* **1990**, *38*, 939–949.
- (9) Gao, Z.-G.; Kim, S. K.; Gross, A. E.; Chen, A.; Blaustein, J. B.; Jacobson, K. A. Identification of essential residues involved in the allosteric modulation of the human A₃ adenosine receptor. *Mol. Pharmacol.* **2003**, *63*, 10211031.

- (10) Bridges, T. M.; Lindsley, C. W. G-Protein-coupled receptors: from classical modes of modulation to allosteric mechanisms. *ACS Chem. Biol.* **2008**, *3*, 530–541.
- (11) Schwartz, T. W.; Holst, B. Allosteric enhancers, allosteric agonists and ago-allosteric modulators: where do they bind and how do they act? *Trends Pharmacol. Sci.* **2007**, *28*, 366–373.
- (12) Leach, K.; Sexton, P. M.; Christopoulos, A. Allosteric GPCR modulators: taking advantage of permissive receptor pharmacology. *Trends Pharmacol. Sci.* **2007**, *28*, 382–389.
- (13) Gao, Z.-G.; Jacobson, K. A. Allosterism in membrane receptors. *Drug Discovery Today* **2006**, *11*, 191–202.
- (14) Gao, Z.-G.; van Muijlwijk-Koezen, J. E.; Chen, A.; Müller, C. E.; IJzerman, A. P.; Jacobson, K. A. Allosteric modulation of A₃ adenosine receptors by a series of 3-(2-pyridinyl)isoquinoline derivatives. *Mol. Pharmacol.* **2001**, *60*, 1057–1063.
- (15) Göblyös, A.; Gao, Z.-G.; Brussee, J.; Connestari, R.; Santiago, S. N.; Ye, K.; IJzerman, A. P.; Jacobson, K. A. Structure activity relationships of 1*H*-imidazo[4,5-*c*]quinolin-4-amine derivatives new as allosteric enhancers of the A₃ adenosine receptor. *J. Med. Chem.* **2006**, *49*, 3354–3361.
- (16) Gao, Z.-G.; Kim, S. G.; Soltysiak, K. A.; Melman, N.; IJzerman, A. P.; Jacobson, K. A. Selective allosteric enhancement of agonist binding and function at human A₃ adenosine receptors by a series of imidazoquinoline derivatives. *Mol. Pharmacol.* **2002**, *62*, 81–89.
- (17) Gao, Z.-G.; Ye, K.; Göblyös, A.; IJzerman, A. P.; Jacobson, K. A. Flexible modulation of agonist efficacy at the human A₃ adenosine receptor by an imidazoquinoline allosteric enhancer LUF6000 and its analogues. *BMC Pharmacol.* **2008**, *8*, 20.
- (18) van Galen, P. J. M.; Nissen, P.; van Wijngaarden, I.; IJzerman, A. P.; Soudijn, W. 1*H*-imidazo[4,5-*c*]quinolin-4-amines: novel non-xanthine adenosine antagonists. *J. Med. Chem.* **1991**, *34*, 1202–1206.
- (19) Klotz, K. N.; Lohse, M. J.; Schwabe, U.; Cristalli, G.; Vittori, S.; Grifantini, M. 2-Chloro-*N*⁶-[³H]cyclopentyladenosine ([³H]CCPA)—a high affinity agonist radioligand for A₁ adenosine receptors. *Naunyn Schmiedeberg's Arch. Pharmacol.* **1989**, *340*, 679–683.
- (20) Jarvis, M. F.; Schutz, R.; Hutchison, A. J.; Do, E.; Sills, M. A.; Williams, M. [³H]CGS 21680, an A₂ selective adenosine receptor agonist directly labels A₂ receptors in rat brain tissue. *J. Pharmacol. Exp. Ther.* **1989**, *251*, 888–893.
- (21) Olah, M. E.; Gallo-Rodriguez, C.; Jacobson, K. A.; Stiles, G. L. ¹²⁵I-4-Aminobenzyl-5'-*N*-methylcarboxamidoadenosine, a high affinity radioligand for the rat A₃ adenosine receptor. *Mol. Pharmacol.* **1994**, *45*, 978–982.
- (22) Lorenzen, A.; Lang, H.; Schwabe, U. Activation of various subtypes of G-protein alpha subunits by partial agonists of the adenosine A₁ receptor. *Biochem. Pharmacol.* **1998**, *56*, 1287–1293.
- (23) Wolber, C.; Fozard, J. R. The receptor mechanism mediating the contractile response to adenosine on lung parenchymal strips from actively sensitised, allergen-challenged Brown Norway rats. *Naunyn Schmiedeberg's Arch. Pharmacol.* **2005**, *371*, 158–168.
- (24) Fossetta, J.; Jackson, J.; Deno, G.; Fan, X.; Du, X. K.; Bober, L.; Soude-Bermejo, A.; de Bouteiller, O.; Caux, C.; Lunn, C.; Lundell, D.; Palmer, R. K. Pharmacological analysis of calcium responses mediated by the human A₃ adenosine receptor in monocyte-derived dendritic cells and recombinant cells. *Mol. Pharmacol.* **2003**, *63*, 342–350.
- (25) Gao, Z. G.; Jacobson, K. A. Translocation of arrestin induced by human A₃ adenosine receptor ligands in an engineered cell line: comparison with G protein-dependent pathways. *Pharmacol. Res.* **2008**, *57*, 303–311.
- (26) Nordstedt, C.; Fredholm, B. B. A modification of a protein-binding method for rapid quantification of cAMP in cell-culture supernatants and body fluid. *Anal. Biochem.* **1990**, *189*, 231–234.
- (27) Bradford, M. M. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein–dye binding. *Anal. Biochem.* **1976**, *72*, 248–254.
- (28) Cheng, Y.-C.; Prusoff, W. H. Relationship between the inhibition constant (*K_i*) and the concentration of inhibitor which causes 50% inhibition (*I*₅₀) of an enzymatic reaction. *Biochem. Pharmacol.* **1973**, *22*, 3099–3108.

JM801659W