## *Endo*-Selective Enyne Ring-Closing Metathesis Promoted by Stereogenic-at-W Mono-Pyrrolide Complexes

## Yu Zhao,<sup>†</sup> Amir H. Hoveyda,<sup>\*,‡</sup> and Richard R. Schrock<sup>\*,†</sup>

Department of Chemistry, Merkert Chemistry Center, Boston College Chestnut Hill, Massachusetts 02467, United States, and Department of Chemistry, Massachusetts Institute of Technology, Cambridge, Massachusetts 02139, United States

amir.hoveyda@bc.edu; rrs@mit.edu

## Received December 16, 2010



The utility of W-alkylidene complexes for enyne ring-closing metathesis is demonstrated in a direct comparison with Mo-based analogs. Tungsten complexes lead to less alkyne oligomerization and higher levels of *endo*-selectivity and enantioselectivity.

Although high oxidation state W-based imido alkylidene bis-alkoxide complexes were the first well-defined olefin metathesis catalysts to be developed,<sup>1</sup> they have not been explored in chemical synthesis as extensively as related Mo catalysts.<sup>2</sup> Among the reasons for Mo catalysts being preferred are the relatively high stability of tungstacyclobutane intermediates toward loss of olefin, a perceived higher sensitivity of tungsten complexes to certain functional groups, and commercial availability of some Mo complexes. In virtually no olefin metathesis reactions have activities and selectivities of tungsten and molybdenum catalysts been compared directly in more than a cursory manner. Recently, we have turned to an exploration of monoaryloxide pyrrolide (MAP) imido alkylidene catalysts,<sup>3</sup> which are highly reactive and selective for a variety of olefin metathesis processes.<sup>4</sup> In the process, we discovered that W-based catalysts can be dramatically more selective than Mo-based catalysts, for example, for homocoupling of terminal olefins to generate Z internal olefins with high selectivity.<sup>5</sup> Structural and kinetic studies of tungsten and

**ORGANIC** LETTERS

2011 Vol. 13, No. 4

784-787

Massachusetts Institute of Technology.

<sup>&</sup>lt;sup>‡</sup>Boston College Chestnut Hill.

<sup>(1) (</sup>a) Schaverien, C. J.; Dewan, J. C.; Schrock, R. R. J. Am. Chem. Soc. **1986**, 108, 2771–2773. (b) Schrock, R. R. Chem. Rev. **2002**, 102, 145–180.

<sup>(2)</sup> Schrock, R. R.; Hoveyda, A. H. Angew. Chem., Int. Ed. 2003, 42, 4592–4633.

<sup>(3)</sup> Schrock, R. R. Chem. Rev. 2009, 109, 3211-3226.

<sup>(4)</sup> Enantioselective RCM: (a) Malcolmson, S. J.; Meek, S. J.; Sattely, E. S.; Schrock, R. R.; Hoveyda, A. H. Nature 2008, 456, 933–937.
(b) Sattely, E. S.; Meek, S. J.; Malcolmson, S. J.; Schrock, R. R.; Hoveyda, A. H. J. Am. Chem. Soc. 2009, 131, 943–953. (c) Meek, S. J.; Malcolmson, S. J.; Li, B.; Schrock, R. R.; Hoveyda, A. H. J. Am. Chem. Soc. 2009, 131, 943–953. (c) Meek, S. J.; Malcolmson, S. J.; Li, B.; Schrock, R. R.; Hoveyda, A. H. J. Am. Chem. Soc. 2009, 131, 16407–16409. Enantioselective ring opening-cross metathesis:(d) Ibrahem, I.; Yu, M.; Schrock, R. R.; Hoveyda, A. H. J. Am. Chem. Soc. 2009, 131, 3844–3845. Ring-opening metathesis polymerization:(e) Flook, M. M.; Jiang, A. J.; Schrock, R. R.; Hoveyda, A. H. J. Am. Chem. Soc. 2009, 131, 7962–7963. (f) Flook, M. M.; Gerber, L. C. H.; Debelouchina, G. T.; Schrock, R. R. Macromolecules 2010, 43, 7515–7522. Ethenolysis reactions:(g) Marinescu, S. C.; Schrock, R. R.; Müller, P.; Hoveyda, A. H. J. Am. Chem. Soc. 2009, 131, 10840–10841. (5) Jiang, A. J.; Zhao, Y.; Hoveyda, A. H.; Schrock, R. R. J. Am.

<sup>(3)</sup> Jiang, A. J.; Zhao, Y.; Hoveyda, A. H.; Schröck, R. K. J. Am. Chem. Soc. **2009**, 131, 16630–16631.

molybdenum MAP complexes suggest that methylidene intermediates can be relatively stable and long-lived.<sup>6</sup> Accordingly, if both Mo and W MAP methylidene species are long-lived in the appropriate circumstances, higher turnover numbers should be obtained.

Scheme 1. Various Modes of Reaction in Enyne RCM



One property of Mo-MAP complexes is their ability to catalyze envne metathesis reactions to give endo products preferentially (Scheme 1);<sup>7</sup> such cyclic dienes are virtually inaccessible through envne reactions catalyzed by Ru carbenes.<sup>8</sup> Although some envne reactions catalyzed by Ru complexes have been proposed to follow a mechanism in which the olefin binds to the metal first (Scheme 1), we have suggested that the endo product v is generated when the terminal alkyne binds initially to Mo, affording a  $\beta$ metallacyclobutane intermediate iii. Conversely, the exo product *ix* is formed through the intermediate  $\alpha$ -metallacyclobutane vii. Mixtures of endo- and exo-products form through the competitive reaction pathways shown in Scheme 1. One of the limitations in enyne metathesis by Mo-based MAP complexes might arise from multiple reactions that involve a terminal acetylene (i.e., oligomerization or polymerization).<sup>9</sup> The yields and purity of enyne metathesis products prepared through the use of Mo catalysts therefore can be limited. As a consequence of the superior selectivity of tungsten catalysts for Z-selective homocoupling of terminal olefins, presumably as a result of the better-controlled reactivity of W-complexes,<sup>5</sup> we felt compelled to explore the ability of such complexes in promoting envne RCM reactions.

(8) Diver, S. T.; Giessert, A. J. Chem. Rev. 2004, 104, 1317–1382.

**Table 1.** Enyne Metathesis of 1 with Mo- and W-based MAPCatalysts<sup>a</sup>



<sup>*a*</sup> Reactions were performed under N<sub>2</sub> atmosphere with 5 mol % catalyst in C<sub>6</sub>H<sub>6</sub> at 22 °C. <sup>*b*</sup> Conv was based on consumption of substrate. <sup>*c*</sup> Ratios were determined by analysis of 1H NMR spectra of unpurified reaction mixtures.

We began by examining enyne metathesis by Mo- and W-catalysts of tosylamide 1 (Table 1). We found that a W-based catalyst provides levels of selectivity and reactivity at least as high as the analogous Mo-based variant.

Table 2. More	Efficient a	nd Selective	Enyne	Metathesis	with
W-based Catal	ysts <sup>a</sup>				

substrate	product	catalyst	concn	time	conv (%) <sup>b</sup>	endo:exo <sup>c</sup>	yield (%)		
4	5 TsN	Mo-F <sub>6</sub> W-F <sub>6</sub>	0.05 M 0.01 M 0.05 M 0.01 M	15 min 15 min 15 min 15 min	>98 >98 >98 >98	>20:1 >20:1 >20:1 >20:1	<40 <50 71 92		
6 TSN	7 <sub>TSN</sub>	Mo-F <sub>6</sub> W-F <sub>6</sub>	0.05 M 0.01 M 0.05 M 0.01 M	15 min 15 min 15 min 15 min	>98 >98 >98 >98 >98	20:1 >20:1 >20:1 >20:1 >20:1	<40 <40 71 92		
8 N Ts		Mo-F <sub>6</sub> W-F <sub>6</sub>	0.16 M 0.16 M	1 h 1 h	>98 >98	>20:1 >20:1	68 80		
10 PhN Me Me	11 PhN	Mo-F <sub>6</sub> W-F <sub>6</sub>	0.10 M 0.10 M	1 h 1 h	>98 >98	>20:1 >20:1	<60 95		
<sup>12</sup> PhN	13 PhN	Mo-F <sub>6</sub> W-F <sub>6</sub>	0.10 M 0.10 M	1 h 1 h	>98 >98	>20:1 >20:1	60 90		
14 TsN	15 TsN	Mo-F <sub>6</sub> W-F <sub>6</sub>	0.10 M 0.10 M	10 min 10 min	>98 >98	>20:1 >20:1	70 90		
$a^{-c}$ See Table 1.									

Encouraged by these results, we turned to substrates that undergo facile oligomerization in the presence of Mocatalysts. As the data in Table 2 illustrate, in the case of enyne metathesis of **4**, where a medium ring is generated, and of **6**, where the rate of ring-closure is slower, presumably as a consequence of increased substitution at the alkene, relatively high dilution conditions proved to be necessary for Mo-catalyzed processes ( $1 \times 10^{-3}$  and  $2 \times 10^{-3}$  M, respectively).<sup>7b</sup> When the reactions were carried out at a concentration of 0.01 M, < 40% enyne metathesis products were obtained with the Mo-F<sub>6</sub> complex (Table 1) and separation of the desired product from oligomeric side products proved difficult (Table 2).

<sup>(6) (</sup>a) Jiang, A. J.; Simpson, J. H.; Müller, P.; Schrock, R. R. J. Am. Chem. Soc. **2009**, 131, 7770–7780. (b) Schrock, R. R.; King, A. J.; Marinescu, S. C.; Simpson, J. H.; Müller, P. Organometallics **2010**, 29, 5241–5251.

<sup>(7) (</sup>a) Singh, R.; Schrock, R. R.; Müller, P.; Hoveyda, A. H. J. Am. Chem. Soc. 2007, 129, 12654–12655. (b) Lee, Y.-J.; Schrock, R. R.; Hoveyda, A. H. J. Am. Chem. Soc. 2009, 131, 10652–10661.

<sup>(9) (</sup>a) Schrock, R. R. In *Metathesis Polymerization of Olefins and Polymerization of Alkynes*; Imamoglu, Y., Ed.; Kluwer: Norwell, MA, 1998; p 357. (b) Schrock, R. R.; Luo, S.; Lee, J. C., Jr.; Zanetti, N. C.; Davis, W. M. *J. Am. Chem. Soc.* **1996**, *118*, 3883–3895. (c) Fox, H. H.; Schrock, R. R. *Organometallics* **1992**, *11*, 2763–2765.

In contrast, RCM performed in the presence of  $W-F_6$  at a concentration of 0.05 M yielded *endo* products **5** and **7** in 60–70% isolated yield.<sup>10</sup> When the above transformations were carried out with  $W-F_6$  at a concentration of 0.01 M, the yield was >90%. A similar trend was observed for other substrates, such as the aryl-substituted tosylamides **8** and **14**, as well as the anilines **10** and **12**; use of  $W-F_6$  led to the enyne metathesis products in higher yields.

Table 3. Enhanced endo-Selectivity by W-based Catalysts<sup>a</sup>



<sup>*a-c*</sup> See Table 1. <sup>*d*</sup> Catalyst (10 mol %) was used.

At least in certain cases, W-based catalysts are more *endo*-selective than Mo-based alkylidenes. As shown in Table 3, enyne metathesis of internal alkyne substrates **16** with Mo–F<sub>6</sub> yielded a mixture of *endo*- and *exo*-products in a ratio of 5:1. It is plausible that the sterically less accessible internal alkyne discourages Mo-alkylidene alkyne association as depicted in  $\mathbf{ii}$  in Scheme 1, pathway 1, leading to partial reaction through pathway 3 and initiation at the alkene site. An indiscriminate  $\alpha$ - and  $\beta$ -addition of the alkylidene to the internal alkyne may also be responsible for the low selectivity. The use of W–F<sub>6</sub> resulted in a much-improved *endo*-selectivity of 10:1, albeit with a lower reaction efficiency; a 10 mol % loading was necessary to achieve full conversion to the desired product.

We have previously reported examples of enantioselective envne metathesis desymmetrization reactions performed under ethylene. Such processes likely involve cross-metathesis with ethylene, followed by enantioselective RCM of the tetraene intermediate.7b We report here that W-based catalysts provide enhanced endo- as well as enantioselectivity for desymmetrization of aniline dienvne 12. As summarized in Table 4, under otherwise identical conditions, while Mo-based catalyst 20 provided a 2.5:1 mixture of 13 and 19, with only 21% ee for the major isomer 13, the corresponding W-catalyst 21 yielded a higher ratio (8:1 favoring 13) with an ee of 70%. Both catalysts can be generated in situ from alcoholysis of the corresponding bispyrrolide complexes. We note that, unlike the reaction performed in the presence of W-F<sub>6</sub>, which cleanly yielded the desired product, the reaction catalyzed by 20 and 21 led to a significant amount of oligomerization (20-30%). When the same transformations were carried out in the presence of ethylene, the endo- and enantioselectivities of the 
 Table 4. Mo- and W-Catalyzed Enantioselective Ring-Closing

 Enyne Metathesis<sup>a</sup>



<sup>*a-c*</sup> See Table 1. <sup>*d*</sup> Determined by chiral HPLC analysis.

products remained the same, which implies that such reactions proceed through a true enyne metathesis pathway.

 Table 5. Investigation of Imido Groups for Enantioselective

 Ring-Closing Enyne Metathesis<sup>a</sup>



We carried out further structural optimization of the W-based MAP catalysts for envne metathesis of 12. Initial efforts focused on screening of various imido groups (Table 5). Out of four aryl imido groups investigated, the original 2,6-di-iso-propylphenylimido complexes (as in 21) emerged as optimal, especially in terms of enantioselectivity. The effect of different chiral phenoxides derived from monoprotected BINOL or octahydro-BINOLs were examined next (Table 6). Whereas incorporation of different substitution at the 3,3' positions of the chiral diols (different halides in entries 1, 2, 4 and 5, as well as Me in entry 3 in Table 6) resulted in less enantioselective reactions, catalyst 22 with a methyl ether unit on the chiral aryloxide was more reactive and promoted more selective reactions; cyclic diene 13 was formed in 60% isolated yield and 81% ee (entry 6, Table 6).<sup>11</sup> Further solvent screening (Table 7) identified diethyl ether as the optimal solvent in terms of endo-selectivity and enantioselectivity (91% ee), although the yield of the isolated product was only 35% as a consequence of significant competitive oligomerization.

<sup>(10)</sup> See Supporting Information for a direct comparison of crude NMR spectra of reactions performed with  $Mo-F_6$  and  $W-F_6$  complexes.

 Table 6. Investigation of Aryloxide Groups for Enantioselective

 Ring-Closing Enyne Metathesis<sup>a</sup>



In conclusion, we have demonstrated that W-based MAP complexes promote more efficient and *endo*-selective enyne RCM than the corresponding Mo-based variants; the most notable distinction is fewer side reactions with W-based system. The observations described above, along with the findings concerning Z-selective homocoupling of terminal olefins,<sup>5</sup> underline the importance of examining

 
 Table 7. Solvent Effect in W-Catalyzed Enantioselective Ring-Closing Enyne Metathesis<sup>a</sup>



W-based catalysts along with Mo-based catalysts in the context of various applications. The strategies outlined above, involving Mo- as well as W-based catalysts for olefin metathesis, will constitute a critical and unique attribute of the future studies in these laboratories.

Acknowledgment. Financial support was provided by the National Institutes of Health (GM-59426 to R.R. S. and A.H.H.). Dr. Annie J. King (MIT) and Dr. Yeon-Ju Lee (Boston College) are acknowledged for experimental assistance and helpful discussions.

**Supporting Information Available.** Experimental procedures and spectral data for products. This material is available free of charge via the Internet at http://pubs.acs.org.

<sup>(11)</sup> Other catalysts possessing a methyl ether of the other chiral diols were also tested; metathesis reactions with such complexes proved to be less enantioselective. See the Supporting Information for more screening data.