

Photoreactions of bicyclic aziridines with alkenes and alkynes: a novel synthetic methodology for 8-azabicyclo[3.2.1]octane derivatives†

Keitaro Ishii,* Motohiro Kido, Masahiro Noji and Shigeo Sugiyama

Received 7th April 2008, Accepted 6th June 2008

First published as an Advance Article on the web 10th July 2008

DOI: 10.1039/b805801f


The photochemical C–C bond cleavage of bicyclic aziridines **7** and subsequent [3 + 2] cycloaddition with electron-deficient alkenes and alkynes afforded the novel head-to-head adducts selectively and efficiently. The adducts contain the naturally occurring 8-azabicyclo[3.2.1]octane skeleton (*e.g.* tropane alkaloids). The aziridine **8** fused with a 6-membered ring also afforded the cycloadducts but in poor yields. The methylaziridine **9** reacted with an electron-deficient alkene, affording the head-to-tail adduct **23** in addition to head-to-head adducts **22a** and **22b**. The photoreactions of bicyclic aziridines with alkenes and alkynes indicate a similar behavior to that of aziridines with a linear chain.

Introduction

For the construction of pyrrolidine-containing heterocycles, the 1,3-dipolar cycloaddition of azomethine ylides with alkenes is an important and useful strategy.¹ The aziridine ring is cleaved to give the corresponding azomethine ylide on irradiation or under thermal conditions.² In general, the 1,3-dipolar cycloaddition of an azomethine ylide possessing one electron-withdrawing group (EWG) at the ylide carbon and electron-deficient alkenes affords head-to-tail adducts (2,4-disubstituted pyrrolidines).^{2b,3}

Recently, we reported that β -aziridinylacrylonitrile **1** or acrylate **2** undergo ring opening by irradiation or heating and subsequent [3 + 2] cycloaddition with electron-deficient alkenes, leading to head-to-head adducts (2,3-disubstituted pyrrolidines) selectively and efficiently (Scheme 1).⁴ One of the resonance forms, the biradical intermediate **A**, primarily contributes to the cycloaddition rather than the other resonance form, the azomethine ylide **B**. In contrast, the head-to-tail adducts are also formed only in the thermal reactions of the aziridine **2b** because the stabilization of the azomethine ylide intermediate **B** ($R^2 = \text{Ph}$) is increased by the phenyl substituent.^{4c}

We have also reported that the photochemical C–C bond cleavage of bicyclic epoxide **3** and consequent [3 + 2] cycloaddition with ethyl vinyl ether afforded 8-oxobicyclo[3.2.1]octane **4**.⁵ We have considered that the 8-azabicyclo[3.2.1]octane skeleton could be also constructed from bicyclic aziridines. This skeleton occurs naturally in the tropane alkaloids, which possess a wide range of biological activities. The tropane skeletons **6** are mainly synthesized from 3-oxopyridinium betaines **5** by dipolar cycloaddition (Fig. 1).⁶ A synthesis of **6**, involving thermal ring opening of a bicyclic aziridine and cycloaddition, has been reported by only one research group.⁷



Scheme 1




Fig. 1

Meiji Pharmaceutical University, 2-522-1 Noshio, Kiyose, Tokyo, 204-8588, Japan. E-mail: ishiikei@my-pharm.ac.jp; Fax: +81 (0)42 495 8783; Tel: +81 (0)42 495 8783

† Electronic supplementary information (ESI) available: phase-sensitive NOESY of compounds (*E*)-**15a**, (*Z*)-**15a**, (*E*)-**15b**, (*Z*)-**17a**, (*Z*)-**18a**, (*Z*)-**18b** and *c*, (*Z*)-**23** and **25a**. See DOI: 10.1039/b805801f

In this paper, we describe the scope and limitation of the cycloadditions of bicyclic aziridines **7–10** (Fig. 2) with alkenes and alkynes for the construction of the 8-azabicyclo[3.2.1]octane skeleton.




Fig. 2


Results and discussion

Preparation of aziridines

The γ,δ -epimino α,β -unsaturated esters **7** and **8** were prepared by the Horner–Emmons reaction of the ketones **11**⁸ and **12**⁸ with triethyl phosphonoacetate in 89% ($E:Z = 45 : 55$) and 68% ($E:Z = 35 : 65$) yields, respectively. In a similar manner, the δ -methyl γ,δ -epimino α,β -unsaturated ester **9** was obtained from iodoenone **13** by treatment with benzylamine, caesium carbonate and 1,10-phenanthroline (25% yield) and a subsequent Horner–Emmons reaction of the obtained aziridine **14** with triethyl phosphonoacetate in 71% ($E:Z = 49 : 51$) yield. The (*E*) and (*Z*) configurations of **7–9** were determined by means of the chemical shifts of their ¹H NMR spectra (see Experimental); they showed signals due to 1-H of the (*Z*)-isomers at a lower field (δ 3.45–3.69) than those of *E*-isomers (δ 2.06–2.46) because of the deshielding effect of the carboxyl group.⁹ The dinitrile **10** was synthesized from the ketone **11** and malononitrile by the Knoevenagel condensation¹⁰ in 69% yield (Scheme 2).

Reactions of aziridine **7** with various alkenes and alkynes

Direct irradiation of a solution of the aziridine (*Z*)-**7** with 3 equiv. methyl acrylate in acetonitrile with a low-pressure mercury lamp in a quartz test tube at room temperature afforded the head-to-head *exo*- and *endo*-adducts (*Z*)-**15a** and (*Z*)-**15b** in moderate yields, respectively (Table 1, entry 2). The 1,7-configurations of (*Z*)-**15a** and (*Z*)-**15b** were determined by means of the coupling constant of 1-H in the ¹H NMR spectra; they showed a singlet for 1,7-*cis* and a doublet (J 7.0 Hz) for 1,7-*trans*. The coupling constant of 1-H is nearly the same as those of the analogous adducts derived from carbonyl ylides and alkenes.^{5b} The aziridine (*E*)-**7** also reacted with methyl acrylate to give the adducts (*E*)-**15a** and (*E*)-**15b** (Table 1, entry 1), whose stereochemistries could not be determined by the coupling constant of 1-H owing to the overlapping with the signal



Scheme 2 Reagents and conditions: i, (EtO)₂P(O)CH₂CO₂Et, NaH, DMF or CH₂Cl₂, rt; ii, BnNH₂, Cs₂CO₃, xylene, 95 °C; iii, CH₂(CN)₂, NH₄OAc, AcOH, benzene, rt.

of 3-H. Therefore, the stereochemistries of (*E*)-**15a** and (*E*)-**15b** were primarily deduced from the phase-sensitive NOESY spectra (see ESI†).¹¹ The (*E*) and (*Z*) configurations of **15** were assigned by the chemical shifts of 1-H, as mentioned for the case of aziridines **7–9**. The photochemical reactions of aziridine **7** with *tert*-butyl acrylate (entry 3), *N*-phenylmaleimide (entries 4 and 5), dimethyl maleate (entry 6), dimethyl fumarate (entry 7), methyl propiolate (entry 8), and dimethyl acetylenedicarboxylate (entries 9 and 10) were performed, and the results are summarized in Table 1 and Fig. 3.

The *exo*-products (e.g., **15a**, **16a** and **17a**) were formed preferentially and independently of the (*E/Z*)-configuration of the exocyclic double bond. In the transition state **C** (Fig. 3) the carboxy group of the alkenes is presumably orientated to be outside the ring, because of the steric hindrance of the exocyclic double bond of aziridine ring-cleaved intermediates. (*E/Z*)-Isomerization of the exocyclic double bond was not observed during the irradiation, because the isomerization would occur mainly *via* the triplet state of aziridine compounds.^{4a,b} The reaction of aziridine **7** with dimethyl maleate afforded the 6,7-*cis* (*Z*)-**18a** and 6,7-*trans* dicarbonyl compounds (*Z*)-**18b** and (*Z*)-**18c** (entry 6). The results may indicate that the cycloaddition proceeds stepwise and/or that the maleate isomerizes to the corresponding fumarate

Table 1 Photochemical reactions of aziridine **7** with various alkenes and alkynes^a

Entry	Aziridine	Alkene/alkyne	Reaction time/h	Conversion (%)	Products (yield (%) ¹³)
1	(<i>E</i>)	Methyl acrylate	2	100	(<i>E</i>)- 15a (49) and (<i>E</i>)- 15b (22)
2	(<i>Z</i>)	Methyl acrylate	3	100	(<i>Z</i>)- 15a (38) and (<i>Z</i>)- 15b (21)
3	(<i>E</i>)	<i>t</i> -Butyl acrylate	4	94	(<i>E</i>)- 16a (62) and (<i>E</i>)- 16b (3)
4	(<i>E</i>)	<i>N</i> -Phenylmaleimide	3	80	(<i>E</i>)- 17a (59) and (<i>E</i>)- 17b (20)
5	(<i>Z</i>)	<i>N</i> -Phenylmaleimide	2	81	(<i>Z</i>)- 17a (60) and (<i>Z</i>)- 17b (12)
6	(<i>Z</i>)	Dimethyl maleate	4	99	(<i>Z</i>)- 18a (31), (<i>Z</i>)- 18b (14) and (<i>Z</i>)- 18c (3)
7	(<i>Z</i>)	Dimethyl fumarate	3	100	(<i>Z</i>)- 18b (52) and (<i>Z</i>)- 18c (21)
8	(<i>Z</i>)	Methyl propiolate	4	80	(<i>Z</i>)- 19 (59)
9	(<i>E</i>)	Dimethyl acetylenedicarboxylate	1.5	98	(<i>E</i>)- 20 (62)
10	(<i>Z</i>)	Dimethyl acetylenedicarboxylate	4	100	(<i>Z</i>)- 20 (55)

^a A 0.060 mol dm⁻³ solution of aziridine in acetonitrile with 3 equiv. of alkene or alkyne was irradiated at RT.




Fig. 3

photochemically. The reactions of aziridine **7** and alkynes also gave the adducts **19** and **20** in moderate yields (entries 8–10). In comparison with the cycloaddition of linear aziridine **1** with dimethyl maleate, dimethyl fumarate, and dimethyl acetylenedicarboxylate,^{4b} bicyclic aziridine **7** reacted smoothly with those compounds to afford the corresponding adducts.

The structures of adducts **16**–**20** were also deduced on the basis of their spectral data. In particular, the stereochemistries of **16**–**18** were determined by the chemical shift and the multiplicity of the signals at H-C(1) and H-C(7) for **16** and **18** and at H-C(6) and H-C(7) for **17** in the ¹H NMR spectra (see Table 2) in comparison with those of the compounds (*E/Z*)-**15a** and (*E/Z*)-**15b**. Furthermore, the structures of (*Z*)-**17a** and (*Z*)-**18a–c** were also confirmed by phase-sensitive NOESY spectra (see ESI).¹¹

Reactions of aziridines **8**, **9** and **10** with an alkene and an alkyne

The photochemical reactions of **8**–**10** with an alkene and an alkyne were performed, and the results are summarized in Table 3 and Fig. 4. In order to study the effects of the fused ring size in the bicyclic aziridine on the [3 + 2] cycloaddition with an alkyne, the aziridine (*Z*)-**8** was irradiated with dimethyl acetylenedicarboxylate. The yield of the adduct (*Z*)-**21** from (*Z*)-**8** drastically decreased (Table 3, entry 1) relative to that of the adduct (*Z*)-**20** from the aziridine (*Z*)-**7** (Table 1, entry 10). The same phenomenon has been also observed in the cycloaddition of 7-membered carbonyl ylide and an alkene (4% yield).^{5b,12}

The methyl substituent at the 5-position in aziridine (*Z*)-**9** had an effect on adduct distribution; the irradiation of (*Z*)-**9** with

Table 2 The chemical shift (ppm) and multiplicity of H-C(1) and H-C(7) in the ¹H NMR spectra (in CDCl₃) for **15**–**18**, **22** and **25**

<i>exo</i> -Adduct	δ _{H-C(1)}	δ _{H-C(7)}	<i>endo</i> -Adduct	δ _{H-C(1)}	δ _{H-C(7)}
(<i>E</i>)- 15a	NA ^a	2.88 dd	(<i>E</i>)- 15b	NA	3.39–3.47 br dt
(<i>Z</i>)- 15a	5.31 s	2.88 dd	(<i>Z</i>)- 15b	5.34 d	3.46–3.56 br dt
(<i>E</i>)- 16a	NA	2.78 dd	(<i>E</i>)- 16b	3.55 d	NA
(<i>E</i>)- 17a	NA	3.27 or 3.31 d ^b	(<i>E</i>)- 17b ^c	3.53–3.56 br d ^d	3.17 dd ^b
(<i>Z</i>)- 17a	5.43 s ^d	—	(<i>Z</i>)- 17b	5.72 d ^d	4.03 dd ^b
(<i>Z</i>)- 18a ^c	5.81 s	3.70 d	(<i>Z</i>)- 18b ^c	5.87 d	4.28 t
(<i>Z</i>)- 18c ^c	5.89 s	3.65–3.67 br d			
(<i>Z</i>)- 22a	5.13 s	2.78 dd	(<i>Z</i>)- 22b	5.10 d	3.30–3.39 br dt
25a	4.50 s	2.89 dd	25b	4.36 d	3.55–3.64 br dt

^a The signal overlapped with another signal. ^b Values for H-C(6). ^c Spectra taken in C₆D₆. ^d Values for H-C(7).

Table 3 Photochemical reactions of aziridines **8**, **9** and **10** with an alkene and an alkyne^a

Entry	Aziridine	Alkene/Alkyne	Reaction time/h	Conversion (%)	Products and yields (%) ¹³
1	(<i>Z</i>)- 8	Dimethyl acetylenedicarboxylate	3.5	96	(<i>Z</i>)- 21 (17)
2	(<i>Z</i>)- 9	Methyl acrylate	2	69	(<i>Z</i>)- 22a (18), (<i>Z</i>)- 22b (5) and (<i>Z</i>)- 23 (5)
3	(<i>Z</i>)- 9	Dimethyl acetylenedicarboxylate	2	100	(<i>Z</i>)- 24 (61)
4	10	Methyl acrylate	3	100	25a (33) and 25b (13)
5	10	Dimethyl acetylenedicarboxylate	2	98	26 (28)

^a A 0.060 mol dm^{−3} solution of aziridine in acetonitrile with 3 equiv. of alkene or alkyne was irradiated at RT.




Fig. 4

methyl acrylate afforded the head-to-tail adduct (Z)-23 as well as head-to-head adducts (Z)-22a and (Z)-22b (Table 3, entry 2). The formation of the head-to-tail adducts has previously been observed in the thermal reactions of phenyl-substituted aziridines 2b (Scheme 1).^{4c} The adduct (Z)-23 formed presumably *via* the azomethine ylide intermediate stabilized by the methyl substituent. The sum of yields for all adducts from (Z)-9 were reduced owing to steric reasons. The reaction of (Z)-9 with dimethyl acetylenedicarboxylate, however, proceeded smoothly, as did that of 7 (Table 1, entries 9 and 10).

The electron-withdrawing inductive effects in the side chain of the bicyclic aziridine on the [3 + 2] cycloaddition with alkenes were studied. In earlier studies of the 1,3-dipolar cycloaddition of carbonyl ylide and alkenes, the carbonyl ylides possessing a stronger electron-withdrawing substituent (*e.g.*, dinitrile) reacted with electron-rich alkenes better than weaker ones (*e.g.*, diester and mononitrile).^{5c} Linear dicyanoaziridine reacts with non-electron-deficient alkenes (vinyl acetate and isoprene) to give the adducts.^{4c} The reactions of the bicyclic dicyanoaziridine 10 with non-electron-deficient alkenes (vinyl acetate and isoprene) afforded no cycloadducts, and the yields of the reactions of 10 and electron-deficient alkene and alkyne were lower (Table 3, entries 4 and 5) relative to those of (Z)-7 (Table 1, entries 2 and 10).

The stereo- and regiochemistries of 21–25 were deduced on the basis of their spectral data. In particular, (*E/Z*)- and *endo/exo*-configurations of 22 and 25 were determined by the chemical shift and the multiplicity of the ¹H NMR signals at H-C(1) and H-C(7) (see Table 2) in comparison with those of the compounds (*E/Z*)-15a and (*E/Z*)-15b. Furthermore, the structure of 25a was also confirmed by the phase-sensitive NOESY spectrum (see ESI).¹¹ The regiochemistry of (Z)-23 was determined by the ¹H NMR spectrum; the signals at δ 1.83 (dd) and δ 2.51–2.61 (br ddd) due to 2H-C(7) and at δ 4.91 due to H-C(1) were observed. The assignment of the *exo*-configuration of (Z)-23 was deduced from the phase-sensitive NOESY spectrum (see ESI).¹¹

Conclusions

The reactions of bicyclic aziridine 7 with electron-deficient alkenes and alkynes selectively and efficiently afforded the novel head-to-head adducts, whose skeleton, 8-azabicyclo[3.2.1]octane,

is found in nature (*e.g.*, tropane alkaloids). However, the 7-azabicyclo[4.1.0]heptane 8 reacted with alkyne to give the adduct in low yields. The reactions of 5-methylaziridine 9 with electron-deficient alkenes gave head-to-tail adducts 23 in addition to head-to-head adducts. The dicyanoaziridine 10 reacted also with electron-deficient alkenes, but not with non-electron-deficient alkenes. The photoreactions of bicyclic aziridines and alkenes showed similar behavior to that of linear aziridines (*e.g.*, 1 and 2). However, the bicyclic aziridines reacted with alkynes better than linear aziridines.

Experimental

General

Melting points were measured with a Yanaco MP-3 apparatus and are uncorrected. IR spectra were recorded on a Hitachi 215 spectrometer. ¹H NMR spectra were obtained with a JEOL JNM-AL300 (300 MHz), a JEOL JNM-AL400 (400 MHz) or a JEOL JNM-LA500 (500 MHz) spectrometer, and *J* values are given in Hz. ¹³C NMR spectra were recorded on a JEOL JNM-AL300 (75 MHz), a JEOL JNM-AL400 (100 MHz) or a JEOL JNM-LA500 (125 MHz) spectrometer. Unless otherwise noted, NMR spectra were measured in CDCl₃ using tetramethylsilane as an internal standard at rt. Mass spectra (MS) and high-resolution MS (HRMS) were taken on a JEOL JMS-700 spectrometer in the EI mode at ionization potential of 70 eV. Column chromatography was performed with Merck silica gel 60 (230–400 mesh) and preparative TLC with Wakogel B-5F.

An Eikosha 60 W low-pressure mercury lamp was used for irradiation. The photolysis solutions were purged with argon before and during irradiation.

Preparations of aziridines

Ethyl (*E*)-(6-benzyl-6-azabicyclo[3.1.0]hex-2-ylidene)acetate [(*E*)-7] and ethyl (*Z*)-(6-benzyl-6-azabicyclo[3.1.0]hex-2-ylidene)acetate [(*Z*)-7]. To a suspension of NaH [197 mg, 8.2 mmol; prepared from a NaH dispersion (60%, 329 mg) by washing it twice with hexane (5 cm³)] in dry DMF (15 cm³)] was added dropwise triethyl phosphonoacetate (1.65 cm³, 8.2 mmol) at room temperature. After the mixture had been stirred for 20 min at room

temperature, ketone **11**⁸ (1.40 g, 7.48 mmol) was added dropwise, and the stirring was continued for 1 h at room temperature. Ice–water was added to the mixture, and the organic phase was extracted with diethyl ether. The ethereal extract was washed with brine, dried with MgSO₄, and concentrated *in vacuo*, giving a residue that was subjected to flash column chromatography [methylene chloride–ethyl acetate (29 : 1)] to afford (*E*)-**7** (761 mg, 40%) and (*Z*)-**7** (940 mg, 49%).

(*E*)-**7**, an oil (Found: M⁺ 257.1422. C₁₆H₁₉NO₂ requires *M*, 257.1416; *v*_{max} (film)/cm^{−1} 1690 (C=O); δ_{H} (300 MHz) 1.27 (3 H, t, *J* 7.2, CH₃), 1.73–1.86 (1 H, br dtd, *J* 13, 9 and 3, 4-H), 2.07–2.15 (1 H, br dd, *J* 13 and 8.5, 4-H), 2.24–2.38 (1 H, m, 3-H), 2.46 (1 H, d, *J* 4.4, 1-H), 2.53–2.56 (1 H, br dd, *J* 4 and 3, 5-H), 3.19 (1 H, dd, *J* 18.5 and 9.4, 3-H), 3.51 and 3.58 (2 H, each d, *J* 14.1, CH₂Ph), 4.15 (2 H, q, *J* 7.2, OCH₂), 5.96 (1 H, t, *J* 2.2, C=CH) and 7.23–7.34 (5 H, m, Ph); δ_{C} (75 MHz) 14.3 (q, CH₃), 27.5 (2 t, C-3 and -4), 47.4 and 50.3 (2 d, C-1 and -5), 59.7 and 61.5 (2 t, OCH₂ and CH₂Ph), 113.5 (d, C=CH), 126.9, 127.4 and 128.3 (3 d, 5 C in Ph), 138.8 (s, 1 C in Ph), 164.0 (s, C-2) and 166.4 (s, CO); *m/z* 257 (M⁺, 82%), 138 (35), 120 (29) and 91 (100).

(*Z*)-**7**, an oil (Found: M⁺ 257.1417. C₁₆H₁₉NO₂ requires *M*, 257.1416; *v*_{max} (film)/cm^{−1} 1695 (C=O); δ_{H} (300 MHz) 1.25 (3 H, t, *J* 7.2, CH₃), 1.66–1.78 (1 H, m, 4-H), 1.97–2.05 (1 H, br dd, *J* 13 and 8, 4-H), 2.19–2.29 (1 H, br dd, *J* 17 and 9, 3-H), 2.33–2.47 (1 H, m, 3-H), 2.56 (1 H, dd, *J* 4.2 and 3.3, 5-H), 3.40 and 3.76 (2 H, each d, *J* 13.9, CH₂Ph), 3.69 (1 H, d, *J* 4.2, 1-H), 4.17 (2 H, q, *J* 7.2, OCH₂), 5.84 (1 H, br s, C=CH) and 7.21–7.37 (5 H, m, Ph); δ_{C} (75 MHz) 14.3 (q, CH₃), 26.3 and 30.7 (2 t, C-3 and -4), 45.6 and 48.9 (2 d, C-1 and -5), 59.7 and 61.6 (2 t, OCH₂ and CH₂Ph), 114.4 (d, C=CH), 126.9, 127.6 and 128.2 (3 d, 5 C in Ph), 139.0 (s, C in Ph), 163.5 (s, C-2) and 166.5 (s, CO); *m/z* 257 (M⁺, 89%), 138 (38), 120 (29) and 91 (100).

Ethyl (*E*)-(7-benzyl-7-azabicyclo[4.1.0]hept-2-ylidene)acetate [(*E*)-8**] and ethyl (*Z*)-(7-benzyl-7-azabicyclo[4.1.0]hept-2-ylidene)acetate [(*Z*)-**8**].** By analogy with the synthesis of **7**, ketone **12**⁸ (2.60 g, 12.9 mmol) was treated with NaH (60%, 568 mg, 14.2 mmol) and triethyl phosphonoacetate (2.83 cm³, 14.2 mmol) in dry DMF (26 cm³) at room temperature, and the resulting mixture was stirred for 1 h at room temperature. Flash column chromatography [hexane–ethyl acetate (4 : 1)] of the reaction mixture afforded (*E*)-**8** (848 mg, 24%) and (*Z*)-**8** (1.53 g, 44%).

(*E*)-**8**, an oil (Found: M⁺ 271.1568. C₁₇H₂₁NO₂ requires *M*, 271.1572; *v*_{max} (film)/cm^{−1} 1710 (C=O); δ_{H} (300 MHz) 1.27 (3 H, t, *J* 7.2, CH₃), 1.35–1.46 (1 H, m, 4-H), 1.66–1.77 (2 H, m, 4- and 5-H), 1.93–2.14 (2 H, m, 5- and 6-H), 2.06 (1 H, d, *J* 6.1, 1-H), 2.60–2.72 (1 H, m, 3-H), 2.79–2.89 (1 H, br d, *J* 17, 3-H), 3.47 and 3.67 (2 H, each d, *J* 14.0, CH₂Ph), 4.14 (2 H, q, *J* 7.2, OCH₂), 5.85 (1 H, t, *J* 1.8, C=CH) and 7.21–7.33 (5 H, m, Ph); δ_{C} (100 MHz) 14.5 (q, CH₃), 19.5 (t, C-4), 23.7 and 25.9 (2 t, C-3 and -5), 42.2 and 45.0 (2 d, C-1 and -6), 59.6 and 63.9 (2 t, OCH₂ and CH₂Ph), 117.3 (d, C=CH), 126.7, 127.3 and 128.1 (3 d, 5 C in Ph), 138.9 (s, C in Ph), 158.3 (s, C-2) and 165.8 (s, CO); *m/z* 271 (M⁺, 96%), 226 (25), 198 (43), 180 (100), 152 (66), 134 (30), 124 (18), 106 (24), 91 (90) and 58 (35).

(*Z*)-**8**, an oil (Found: M⁺ 271.1577. C₁₇H₂₁NO₂ requires *M*, 271.1572; *v*_{max} (film)/cm^{−1} 1710 (C=O); δ_{H} (300 MHz) 1.27 (3 H, t, *J* 7.2, CH₃), 1.38–1.49 (1 H, m, 4-H), 1.58–1.71 and 1.73–1.84 (2 H, 2 m, 4- and 5-H), 1.94–2.06 (3 H, m, 3-, 5- and 6-H),

2.34 (1 H, dddd, *J* 14.7, 8.1, 3.5 and 1.1, 3-H), 3.27 and 3.12 (2 H, each d, *J* 14.1, CH₂Ph), 3.45 (1 H, d, *J* 6.1, 1-H), 4.14 and 4.15 (2 H, each q, *J* 7.2, OCH₂), 5.84–5.85 (1 H, br s, C=CH) and 7.20–7.37 (5 H, m, Ph); δ_{C} (100 MHz) 14.5 (q, CH₃), 20.9 (t, C-4), 24.2 and 32.2 (2 t, C-3 and -5), 39.3 and 42.0 (2 d, C-1 and C-6), 59.7 and 64.1 (2 t, OCH₂ and CH₂Ph), 117.9 (d, C=CH), 126.5, 127.5 and 128.0 (3 d, 5 C in Ph), 139.3 (s, C in Ph), 158.1 (s, C-2) and 166.4 (s, CO); *m/z* 271 (M⁺, 100%), 242 (21), 226 (18), 198 (38), 180 (93), 152 (60), 134 (27), 124 (18), 106 (19) and 91 (90).

Ethyl (*E*)-(6-benzyl-5-methyl-6-azabicyclo[3.1.0]hex-2-ylidene)acetate [(*E*)-9**] and ethyl (*Z*)-(6-benzyl-5-methyl-6-azabicyclo[3.1.0]hex-2-ylidene)acetate [(*Z*)-**9**].** To a solution of iodoenone **13** (670 mg, 3.02 mmol) in xylene (6 cm³) were added benzylamine (0.50 cm³, 4.53 mmol), caesium carbonate (1.47 g, 4.52 mmol), and 1,10-phenanthroline (543 mg, 3.02 mmol). The reaction mixture was stirred 2 h at 50 °C. After silica gel filtration, preparative TLC [hexane–ethyl acetate (2 : 1)] of the filtrate afforded 6-benzyl-5-methyl-6-azabicyclo[3.1.0]hexan-2-one **14** (150 mg, 25%), an oil (Found: M⁺ 201.1146. C₁₅H₁₅NO requires *M*, 201.1154; *v*_{max} (film)/cm^{−1} 1705 (C=O); δ_{H} (300 MHz) 1.54 (3 H, s, CH₃), 1.85 (1 H, dt, *J* 12.7 and 9.2, 4-H), 1.96–2.06 (1 H, m, 3-H), 2.04 (1 H, s, 1-H), 2.18–2.27 (1 H, m, 4-H), 2.39–2.51 (1 H, br dt, *J* 17 and 9, 3-H), 3.74 (2 H, s, CH₂Ph) and 7.22–7.35 (5 H, m, Ph); δ_{C} (100 MHz) 13.6 (q, CH₃), 31.5 and 34.3 (2 t, C-3 and -4), 50.0 (s, C-5), 52.8 (d, C-1), 55.3 (t, CH₂Ph), 126.1, 127.1 and 128.3 (3 d, 5 C in Ph), 138.8 (s, C in Ph) and 212.7 (s, C-2); *m/z* 201 (M⁺, 20%), 145 (5), 91 (100), 82 (20) and 65 (7).

By analogy with the synthesis of **7**, ketone **14** (252 mg, 1.25 mmol) was treated with NaH (60%, 55 mg, 1.38 mmol) and triethyl phosphonoacetate (0.27 cm³, 1.38 mmol) in dry methylene chloride (3 cm³) at room temperature, and the resulting mixture was stirred for 1 h at room temperature. Flash column chromatography [cyclohexane–ethyl acetate (3 : 1)] of the reaction mixture afforded (*E*)-**9** (117.8 mg, 35%) and (*Z*)-**9** (121.3 mg, 36%).

(*E*)-**9**, an oil (Found: M⁺ 271.1574. C₁₇H₂₁NO₂ requires *M*, 271.1572; *v*_{max} (film)/cm^{−1} 1705 (C=O); δ_{H} (300 MHz) 1.27 (3 H, t, *J* 7.2, CH₃), 1.50 (3 H, m, 5-CH₃), 1.72 (1 H, dt, *J* 12.8 and 9.2, 4-H), 2.08 (1 H, dd, *J* 12.8 and 9.0, 4-H), 2.22 (1 H, s, 1-H), 2.36 (1 H, dtd, *J* 18.4, 9.2 and 2.8, 3-H), 3.21 (1 H, dd, *J* 18.4 and 9.0, 3-H), 3.67 and 3.81 (2 H, each d, *J* 14.8, CH₂Ph), 4.15 (2 H, q, *J* 7.2, OCH₂), 5.88 (1 H, br s, C=CH) and 7.19–7.34 (5 H, m, Ph); δ_{C} (100 MHz) 13.4 (q, CH₃), 14.4 (q, 5-CH₃), 28.1 (t, C-4), 34.9 (t, C-3), 50.8 (s, C-5), 55.4 (t, CH₂Ph), 55.6 (d, C-1), 59.6 (t, OCH₂), 112.5 (d, C=CH), 126.4, 126.9 and 128.1 (3 d, 5 C in Ph), 139.4 (s, C in Ph), 165.6 (s, C-2) and 166.1 (s, CO); *m/z* 271 (M⁺, 100%), 226 (20), 180 (58), 152 (75), 134 (73), 106 (21) and 91 (66).

(*Z*)-**9**, an oil (Found: M⁺ 271.1575. C₁₇H₂₁NO₂ requires *M*, 271.1572; *v*_{max} (film)/cm^{−1} 1705 (C=O); δ_{H} (300 MHz) 1.25 (3 H, t, *J* 7.2, CH₃), 1.49 (3 H, m, 5-CH₃), 1.59–1.71 (1 H, m, H-4), 1.94–2.02 (1 H, br dd, *J* 13 and 8.5, 4-H), 2.18–2.27 (1 H, br dd, *J* 17 and 8.5, 3-H), 2.39–2.52 (1 H, m, 3-H), 3.49 (1 H, s, 1-H), 3.74, 3.84 (2 H, each d, *J* 14.7 Hz, CH₂Ph), 4.16 (2 H, q, *J* 7.2 Hz, OCH₂), 5.80 (1 H, br s, C=CH) and 7.18–7.38 (5 H, m, Ph); δ_{C} (100 MHz) 13.7 and 14.5 (2 q, 2 CH₃), 31.5 and 33.7 (2 t, C-3 and -4), 51.2 (d, C-1), 52.4 (s, C-5), 55.6 and 59.7 (2 t, OCH₂ and CH₂Ph), 113.4 (d, C=CH), 126.4, 127.1 and 128.0 (3 d, 5 C in Ph), 139.8 (s, C in Ph), 165.2 (s, C-2) and 166.4 (s, CO); *m/z* 271 (M⁺, 100%), 226 (18), 180 (55), 152 (74), 134 (70), 106 (20) and 91 (62).

(6-Benzyl-6-azabicyclo[3.1.0]hex-2-ylidene)malononitrile (10).

To a suspension of the ketone **11**⁸ (850 mg, 4.55 mmol) in dry benzene (1.7 cm³) was added malononitrile (0.29 cm³, 4.55 mmol), ammonium acetate (70 mg, 0.91 mmol) and acetic acid (0.26 cm³, 4.55 mmol). The reaction mixture was stirred for 5 h at room temperature. The reaction was quenched with water and the organic phase was extracted with ether. The organic extract was washed with aqueous sat. NaHCO₃ and brine, dried with MgSO₄, and concentrated *in vacuo*, giving a residue that was subjected to flash column chromatography [hexane–ether (1 : 1)] to afford **10** (733 mg, 69%) as an oil (Found: M⁺ 235.1104. C₁₅H₁₃N₃ requires M, 235.1109; $\nu_{\max}$ (CHCl₃)/cm⁻¹ 2360 (C≡N); δ_{H} (400 MHz) 1.96 (1 H, dtd, *J* 13.4, 9.0 and 2.8, 4-H), 2.20 (1 H, ddd, *J* 13.4, 8.3 and 1.1, 4-H), 2.50–2.63 (1 H, br td, *J* 19 and 9, 3-H), 2.74–2.84 (1 H, br dd, *J* 19 and 9, 3-H), 2.93–2.98 (2 H, m, 1- and 5-H), 3.49 and 3.74 (2 H, each d, *J* 13.4, CH₂Ph) and 7.27–7.40 (5 H, m, Ph); δ_{C} (100 MHz) 27.5 and 31.1 (2 t, C-3 and -4), 48.0 and 51.4 (2 d, C-1 and -5), 61.8 (t, CH₂Ph), 81.8 [s, C(CN)₂], 111.4 and 111.5 (2 s, 2 CN), 127.5, 127.6 and 128.5 (3 d, 5 C in Ph), 137.2 (s, C in Ph) and 186.0 (s, C-2); *m/z* 235 (M⁺, 7%), 104 (2), 91 (100), 77 (2) and 65 (8).

Reactions of aziridines 7–10 with alkenes and alkynes

A 0.060 mol dm⁻³ solution of aziridines **7–10** in dry acetonitrile with 3 equiv. of with alkenes and alkynes was irradiated with a low-pressure mercury lamp in a quartz test tube at room temperature. After removal of the solvent, flash column chromatography afforded the adducts. The results are summarized in Tables 1 and 3 and Fig. 3 and 4.

Ethyl (E,1RS,5RS,7RS)-(8-benzyl-7-methoxycarbonyl-8-azabicyclo[3.2.1]oct-2-ylidene)acetate [(E)-15a]. An oil (Found: M⁺ 343.1779. C₂₀H₂₅NO₄ requires M, 343.1784; $\nu_{\max}$ (CHCl₃)/cm⁻¹ 1720 sh, 1710 (C=O); δ_{H} (300 MHz) 1.28 (3 H, t, *J* 7.1, CH₃), 1.42–1.50 and 2.01–2.09 (2 H, each m, 4-H₂), 2.05 (1 H, dd, *J* 12.9 and 9.8 Hz, 6-H), 2.15–2.25 (1 H, m, 3-H), 2.55–2.63 (1 H, br quintet, *J* 6, 6-H), 2.88 (1 H, dd, *J* 9.8 and 5.1, 7-H), 3.38–3.42 (1 H, m, 5-H), 3.65–3.76 (4 H, m, 1-H, 3-H and CH₂Ph), 3.71 (3 H, s, OCH₃), 4.15 and 4.16 (2 H, each q, *J* 7.1, OCH₂), 5.54 (1 H, br d, *J* 2.2, C=CH) and 7.20–7.33 (5 H, m, Ph); δ_{C} (100 MHz) 14.4 (q, CH₃), 21.7 (t, C-4), 27.3 (t, C-3), 31.6 (t, C-6), 47.4 (d, C-7), 52.2 (q, OCH₃), 52.3 (t, CH₂Ph), 57.3 (d, C-5), 59.8 (t, OCH₂), 68.8 (d, C-1), 114.9 (d, C=CH), 126.7, 128.0 and 128.1 (3 d, 5 C in Ph), 138.7 (s, C in Ph), 157.9 (s, C-2) and 166.1 and 174.5 (2 s, 2 CO); *m/z* 343 (M⁺, 54%), 314 (11), 298 (11), 257 (100), 138 (18), 120 (12) and 91 (90).

Ethyl (E,1RS,5RS,7SR)-(8-benzyl-7-methoxycarbonyl-8-azabicyclo[3.2.1]oct-2-ylidene)acetate [(E)-15b]. An oil (Found: M⁺ 343.1776. C₂₀H₂₅NO₄ requires M, 343.1784; $\nu_{\max}$ (film)/cm⁻¹ 1720 sh and 1710 (C=O); δ_{H} (300 MHz) 1.28 (1 H, t, *J* 7.2, CH₃), 1.48–1.56 and 1.98–2.11 (2 H, each m, 4-H₂), 2.26 (1 H, dd, *J* 12.7 and 6.6, 6-H), 2.27–2.42 (2 H, m, 3- and 6-H), 3.33–3.38 (1 H, m, 5-H), 3.39–3.47 (1 H, br dt, *J* 11 and 7, 7-H), 3.55–3.66 (2 H, m, 1- and 3-H), 3.69 (2 H, s, CH₂Ph), 3.60 (3 H, s, OCH₃), 4.14 (2 H, q, *J* 7.2, OCH₂), 5.47 (1 H, d, *J* 2.6, C=CH) and 7.17–7.37 (5 H, m, Ph); δ_{C} (100 MHz) 14.4 (q, CH₃), 21.4 (t, C-4), 27.9 (t, C-3), 29.3 (t, C-6), 47.3 (d, C-7), 51.9 (q, OCH₃), 53.2 (t, CH₂Ph), 57.4 (d, C-5), 59.7 (t, OCH₂), 68.7 (d, C-1), 116.8 (d, C=CH), 126.9, 128.2

and 128.4 (3 d, 5 C in Ph), 138.5 (s, C in Ph), 156.8 (s, C-2) and 165.8 and 172.4 (2 s, 2 CO); *m/z* 343 (M⁺, 59%), 314 (11), 298 (12), 257 (100), 138 (12) and 91 (76).

Ethyl (Z,1RS,5RS,7RS)-(8-benzyl-7-methoxycarbonyl-8-azabicyclo[3.2.1]oct-2-ylidene)acetate [(Z)-15a]. An oil (Found: M⁺ 343.1777. C₂₀H₂₅NO₄ requires M, 343.1784; $\nu_{\max}$ (CHCl₃)/cm⁻¹ 1710 (C=O); δ_{H} (300 MHz) δ 1.12 (3 H, t, *J* 7.2, CH₃), 1.42–1.51 and 2.05–2.13 (2 H, each m, 4-H₂), 1.99 (1 H, dd, *J* 13.2 and 9.7, 6-H), 2.15–2.24 and 2.36–2.49 (2 H, each m, 3-H₂), 2.58–2.67 (1 H, br quintet, *J* 6.5, 6-H), 2.89 (1 H, dd, *J* 9.7 and 5.3, 7-H), 3.36–3.41 (1 H, m, 5-H), 3.70 and 3.75 (2 H, each d, *J* 14, CH₂Ph), 3.73 (3 H, s, OCH₃), 3.91 and 4.02 (2 H, each dq, *J* 10.8 and 7.2, OCH₂), 5.31 (1 H, s, 1-H), 5.66 (1 H, d, *J* 2.2, C=CH) and 7.17–7.33 (5 H, m, Ph); δ_{C} (100 MHz) 14.2 (q, CH₃), 28.5, 28.8 and 32.0 (3 t, C-3, -4 and -6), 46.9 (d, C-7), 52.2 (q, OCH₃), 52.3 (t, CH₂Ph), 57.0 (d, C-5), 59.9 (t, OCH₂), 61.2 (d, C-1), 115.1 (d, C=CH), 126.6, 127.9 and 128.1 (3 d, 5 C in Ph), 139.1 (s, C in Ph), 156.3 (s, C-2) and 165.6 and 174.8 (2 s, 2 CO); *m/z* 343 (M⁺, 65%), 314 (15), 298 (13), 257 (100), 138 (11) and 91 (78).

Ethyl (Z,1RS,5RS,7SR)-(8-benzyl-7-methoxycarbonyl-8-azabicyclo[3.2.1]oct-2-ylidene)acetate [(Z)-15b]. An oil (Found: M⁺ 343.1793. C₂₀H₂₅NO₄ requires M, 343.1784; $\nu_{\max}$ (CHCl₃)/cm⁻¹ 1720 and 1710 sh (C=O); δ_{H} (300 MHz) 1.19 (3 H, t, *J* 7.2, CH₃), 1.47–1.57 (1 H, m, 4-H), 2.02–2.23 (3 H, m, 3-, 4- and 6-H), 2.27–2.39 (1 H, m, 6-H), 2.55–2.68 (1 H, m, 3-H), 3.27–3.33 (1 H, m, 5-H), 3.46–3.56 (1 H, br dt, *J* 12 and 7, 7-H), 3.59 (3 H, s, OCH₃), 3.71 (2 H, s, CH₂Ph), 4.02, 4.08 (1 H, each dq, *J* 10.8 and 7.2, OCH₂), 5.34 (1 H, d, *J* 7.0, 1-H), 5.69 (1 H, d, *J* 2.2, C=CH) and 7.19–7.33 (5 H, m, Ph); δ_{C} (100 MHz) 14.4 (q, CH₃), 27.9, 29.1 and 29.6 (3 t, C-3, -4 and -6), 46.8 (d, C-7), 51.7 (q, OCH₃), 53.4 (t, CH₂Ph), 56.7 (d, C-5), 59.7 (t, OCH₂), 61.1 (d, C-1), 116.7 (d, C=CH), 126.8, 128.1 and 128.4 (3 d, 5 C in Ph), 138.8 (s, C in Ph), 155.9 (s, C-2) and 165.4 and 173.0 (2 s, 2 CO); *m/z* 343 (M⁺, 62%), 314 (11), 298 (12), 257 (100), 138 (11) and 91 (65).

Ethyl (E,1RS,5RS,7RS)-(8-benzyl-7-tert-butoxycarbonyl-8-azabicyclo[3.2.1]oct-2-ylidene)acetate [(E)-16a]. An oil (Found: M⁺ 385.2256. C₂₃H₃₁NO₄ requires M, 385.2253; $\nu_{\max}$ (film)/cm⁻¹ 1705 (C=O); δ_{H} (300 MHz) 1.29 (3 H, t, *J* 7.2, CH₃), 1.43 (9 H, s, O^tBu), 1.37–1.50 and 2.01–2.09 (2 H, each m, 4-H₂), 1.93 (1 H, dd, *J* 13 and 9.7, 6-H), 2.13–2.26 (1 H, m, 3-H), 2.55–2.64 (1 H, br quintet, *J* 6, 6-H), 2.78 (1 H, dd, *J* 9.7 and 5.1, 7-H), 3.39–3.43 (1 H, m, 5-H), 3.65 and 3.75 (2 H, each d, *J* 13.9, CH₂Ph), 3.62–3.72 (2 H, m, 1- and 3-H), 4.15 and 4.16 (2 H, each q, *J* 7.2, OCH₂), 5.50 (1 H, d, *J* 2.4, C=CH) and 7.20–7.33 (5 H, m, Ph); δ_{C} (100 MHz) 14.5 (q, CH₃), 21.8 (t, C-4), 27.6 (t, C-3), 28.1 (q, 3 CH₃), 31.2 (t, C-6), 48.6 (d, C-7), 52.3 (t, CH₂Ph), 57.9 (d, C-5), 59.8 (t, OCH₂), 68.9 (d, C-1), 80.4 [s, C(CH₃)₃], 114.7 (d, C=CH), 126.6, 128.0 and 128.1 (3 d, 5 C in Ph), 138.8 (s, C in Ph), 158.3 (s, C-2) and 166.1 and 173.2 (2 s, 2 CO); *m/z* 385 (M⁺, 41%), 328 (35), 312 (21), 257 (100) and 91 (68).

Ethyl (E,1RS,5RS,7SR)-(8-benzyl-7-tert-butoxycarbonyl-8-azabicyclo[3.2.1]oct-2-ylidene)acetate [(E)-16b]. An oil (Found: M⁺ 385.2258. C₂₃H₃₁NO₄ requires M, 385.2253; $\nu_{\max}$ (CHCl₃)/cm⁻¹ 1715 (C=O); δ_{H} (300 MHz) 1.26 (3 H, t, *J* 7.1, CH₃), 1.35 (9 H, s, O^tBu), 1.40–1.56 and 1.98–2.11 (2 H, each m, 4-H₂), 2.17 (1 H, dd, *J* 13.0 and 6.2, 6-H), 2.23–2.38 (2 H, m, 3- and 6-H), 3.28–3.39 (2 H, m, 5- and 7-H), 3.55 (1 H, d, *J* 6.6, 1-H),

3.54–3.65 (1 H, m, 3-H), 3.70 (2 H, s, CH_2Ph), 4.13 and 4.19 (2 H, each dq, J 11.0 and 7.1 Hz, OCH_2), 5.50 (1 H, d, J 2.6, $\text{C}=\text{CH}$) and 7.21–7.33 (5 H, m, Ph); δ_{C} (100 MHz) 14.5 (q, CH_3), 21.6 (t, C-4), 28.0 (t, C-3), 28.0 (q, 3CH_3), 29.3 (t, C-6), 47.9 (d, C-7), 53.1 (t, CH_2Ph), 57.5 (d, C-5), 59.7 (t, OCH_2), 68.6 (d, C-1), 80.7 [s, $\text{C}(\text{CH}_3)_3$], 117.1 (d, $\text{C}=\text{CH}$), 126.8, 128.2 and 128.4 (3 d, 5 C in Ph), 138.7 (s, C in Ph), 156.9 (s, C-2) and 165.8 and 171.3 (2 s, 2CO); m/z 385 (M^+ , 49%), 312 (18), 257 (100) and 91 (56).

Ethyl (E,1RS,2SR,6RS,7RS)-(11-benzyl-3,5-dioxo-4-phenyl-4,11-diazatricyclo[5.3.1.0^{2,6}]undec-8-ylidene)acetate [(E)-17a]. An oil (Found: M^+ 430.1887. $\text{C}_{26}\text{H}_{26}\text{N}_2\text{O}_4$ requires M , 430.1893); ν_{max} (CHCl_3)/ cm^{-1} 1710 ($\text{C}=\text{O}$); δ_{H} (300 MHz) 1.31 (3 H, t, J 7.1, CH_3), 1.65–1.70 (1 H, br d, J 8, 10-H), 2.13–2.39 (2 H, m, 9- and 10-H), 3.27 and 3.31 (2 H, each d, J 7.6, 2- and 6-H), 3.75–3.93 (5 H, m, 1-H, 7-H, 9-H and CH_2Ph), 4.19 and 4.22 (2 H, each dq, J 11.0 and 7.1, OCH_2), 5.72 (1 H, br s, $\text{C}=\text{CH}$) and 7.18–7.55 (10 H, m, 2 Ph); δ_{C} (100 MHz) 14.4 (q, CH_3), 22.1 (t, C-9), 25.0 (t, C-10), 48.8 and 49.5 (2 d, C-2 and -6), 50.1 (t, CH_2Ph), 60.1 (t, OCH_2), 60.7 (d, C-1), 68.0 (d, C-7), 117.3 (d, $\text{C}=\text{CH}$), 126.3, 127.2, 127.8, 128.4, 128.6 and 129.1 (6 d, 10 C in 2 Ph), 132.0 and 137.6 (2 s, 2 C in 2 Ph), 153.8 (s, C-8), 165.5 (s, CO) and 176.6 and 177.5 (2 s, C-3 and -5); m/z 430 (M^+ , 92%), 385 (14), 257 (100) and 91 (66).

Ethyl (E,1RS,2RS,6SR,7RS)-(11-benzyl-3,5-dioxo-4-phenyl-4,11-diazatricyclo[5.3.1.0^{2,6}]undec-8-ylidene)acetate [(E)-17b]. White crystals; mp 168–170 °C (hexane–ethyl acetate) (Found: C, 72.44; H, 6.29; N, 6.34. $\text{C}_{26}\text{H}_{26}\text{N}_2\text{O}_4$ requires C, 72.54; H, 6.09; N, 6.51%); ν_{max} (CHCl_3)/ cm^{-1} 1715 ($\text{C}=\text{O}$); δ_{H} (300 MHz) 1.28 (1 H, t, J 7.1, CH_3), 1.90–1.97 (1 H, br dd, J 14 and 9, 10-H), 2.05–2.16 (1 H, m, 10-H), 2.17–2.28 (1 H, m, 9-H), 3.67 (1 H, dd, J 19.0 and 8.1, 9-H), 3.76 (2 H, s, CH_2Ph), 3.74–3.80 (1 H, m, 1-H), 3.85–3.97 (3 H, m, 2-, 6- and 7-H), 4.14 (2 H, q, J 7.1, OCH_2), 5.67 (1 H, d, J 2.4, $\text{C}=\text{CH}$) and 7.21–7.52 (10 H, m, 2Ph); δ_{C} (400 MHz; C_6D_6) 0.91 (3 H, t, J 7.1, CH_3), 1.64–1.78 (2 H, m, 10-H₂), 2.32–2.43 (1 H, m, 9-H), 3.10 (1 H, dd, J 10.3 and 7.3, 2-H), 3.17 (1 H, dd, J 10.3 and 7.6, 6-H), 3.24–3.29 (1 H, m, 1-H), 3.27 (2 H, s, CH_2Ph), 3.53–3.56 (1 H, br d, J 7, 7-H), 3.77–3.84 (1 H, br dd, J 19 and 7, 9-H), 3.88 and 3.96 (2 H, each dq, J 11.0 and 7.1, OCH_2), 5.72 (1 H, d, J 2.7, $\text{C}=\text{CH}$) and 7.05–7.40 (10 H, m, 2 Ph); δ_{C} (100 MHz) 14.4 (q, CH_3), 22.1 and 22.2 (2 t, C-9 and -10), 48.6 (d, C-2), 50.3 (d, C-6), 53.5 (t, CH_2Ph), 58.8 (d, C-1), 60.0 (t, OCH_2), 67.2 (d, C-7), 119.0 (d, $\text{C}=\text{CH}$), 125.9, 127.4, 128.36, 128.41, 128.6 and 129.1 (6 d, 10 C in 2 Ph), 131.5 and 137.3 (2 s, 2 C in 2 Ph), 152.5 (s, C-8), 165.4 (s, CO) and 174.0 and 175.3 (2 s, C-3 and -5); EI-MS m/z 430 (M^+ , 97%), 385 (13), 257 (100) and 91 (52).

Ethyl (Z,1RS,2SR,6RS,7RS)-(11-benzyl-3,5-dioxo-4-phenyl-4,11-diazatricyclo[5.3.1.0^{2,6}]undec-8-ylidene)acetate [(Z)-17a]. White crystals; mp 152–154 °C (hexane–ethyl acetate) (Found: C, 72.25; H, 6.01; N, 6.46. $\text{C}_{26}\text{H}_{26}\text{N}_2\text{O}_4$ requires C, 72.54; H, 6.09; N, 6.51%); ν_{max} (CHCl_3)/ cm^{-1} 1710 ($\text{C}=\text{O}$); δ_{H} (300 MHz) 1.13 (3 H, t, J 7.1, CH_3), 1.67–1.73 (1 H, m, 10-H), 2.25–2.37 (2 H, m, 9- and 10-H), 2.42–2.49 (1 H, m, 9-H), 3.29 (2 H, s, 2- and 6-H), 3.74–3.78 (1 H, br s, 1-H), 3.75 and 3.82 (2 H, each d, J 13.7, CH_2Ph), 3.69 and 4.07 (2 H, each dq, J 10.7 and 7.1 Hz, OCH_2), 5.43 (1 H, s, 7-H), 5.85 (1 H, d, J 2.0, $\text{C}=\text{CH}$) and 7.16–7.53 (5 H, m, 2 Ph); δ_{C} (100 MHz) 14.1 (q, CH_3), 26.4 and 29.0 (2 t, C-9 and

-10), 49.1 and 49.2 (2 d, C-2 and -6), 49.9 (t, CH_2Ph), 60.2 and 60.6 (2 d, C-1 and -7), 60.4 (t, OCH_2), 118.0 (d, $\text{C}=\text{CH}$), 126.3, 127.1, 127.7, 128.4, 128.5 and 129.1 (6 d, 10 C in 2 Ph), 132.1 and 138.0 (s, 2 C in 2 Ph), 152.0 (s, C-8), 165.2 (s, CO) and 176.5 and 177.6 (2 s, C-3 and -5); m/z 430 (M^+ , 84%), 401 (14), 385 (13), 357 (11), 257 (100), 185 (10), 138 (11) and 91 (79).

Ethyl (Z,1RS,2RS,6SR,7RS)-(11-benzyl-3,5-dioxo-4-phenyl-4,11-diazatricyclo[5.3.1.0^{2,6}]undec-8-ylidene)acetate [(Z)-17b]. White crystals; mp 130–140 °C (Found: M^+ 430.1897. $\text{C}_{26}\text{H}_{26}\text{N}_2\text{O}_4$ requires M , 430.1893); ν_{max} (CHCl_3)/ cm^{-1} 1705 ($\text{C}=\text{O}$); δ_{H} (300 MHz) 1.22 (1 H, t, J 7.2, CH_3), 1.84–1.96 (1 H, m, 10-H), 2.12–2.37 (2 H, m, 9- and 10-H), 2.37–2.48 (1 H, m, 9-H), 3.67–3.72 (1 H, m, 1-H), 3.74 and 3.80 (2 H, each d, J 13.4, CH_2Ph), 3.88 (1 H, dd, J 10.3 and 7.7, 2-H), 4.05 (1 H, dd, J 10.3 and 7.9, 6-H), 4.05–4.16 (2 H, m, OCH_2), 5.72 (1 H, d, J 7.9, 7-H), 5.80 (1 H, d, J 1.8, $\text{C}=\text{CH}$) and 7.21–7.49 (10 H, m, 2 Ph); δ_{C} (100 MHz) 14.3 (q, CH_3), 23.6 and 27.3 (2 t, C-9 and -10), 48.9 (d, C-2), 49.9 (d, C-6), 53.9 (t, CH_2Ph), 58.0 (d, C-1), 60.0 (t, OCH_2), 60.2 (d, C-7), 118.6 (d, $\text{C}=\text{CH}$), 125.7, 127.3, 128.3, 128.4 and 129.0 (5 d, 10 C in 2 Ph), 131.7 and 137.6 (s, 2 C in 2 Ph), 152.3 (s, C-8), 165.1 (s, CO) and 173.9 and 175.4 (2 s, C-3 and -5); m/z 430 (M^+ , 92%), 385 (14), 257 (100), 138 (10) and 91 (61).

Dimethyl (Z,1RS,5RS,6SR,7RS)-8-benzyl-2-ethoxycarbonylmethylene-8-azabicyclo[3.2.1]octane-6,7-dicarboxylate [(Z)-18a]. An oil (Found: M^+ 401.1837. $\text{C}_{22}\text{H}_{27}\text{NO}_6$ requires M , 401.1838); ν_{max} (film)/ cm^{-1} 1740 ($\text{C}=\text{O}$); δ_{H} (C_6D_6 ; 300 MHz) 0.78–0.85 (1 H, m, 4-H), 0.90 (3 H, t, J 7.2, CH_3), 1.61–1.80 (3 H, m, 3-H₂ and 4-H), 2.91 (1 H, d, J 10.3, 6-H), 3.07 (1 H, d, J 10.3, 7-H), 3.41 and 3.43 (6 H, each s, 2 OCH_3), 3.66 (1 H, br s, 5-H), 3.75 and 3.91 (2 H, each d, J 14.1, CH_2Ph), 3.83 and 3.91 (2 H, each dq, J 10.8 and 7.2, OCH_2), 5.62 (1 H, br s, $\text{C}=\text{CH}$), 5.81 (1 H, s, 1-H), 7.07 (1 H, t, J 7.3, Ph), 7.18–7.23 (2 H, br t, J 7, Ph) and 7.52–7.56 (2 H, br d, J 7, Ph); δ_{C} (C_6D_6 ; 100 MHz) 14.0 (q, CH_3), 27.6 and 28.1 (2 t, C-3 and -4), 50.7 (2 d, C-6 and -7), 51.3 and 51.4 (2 q, 2 OMe), 51.9 (t, CH_2Ph), 59.7 (d, C-5), 59.7 (t, OCH_2), 61.3 (d, C-1), 115.9 (d, $\text{C}=\text{CH}$), 126.7, 128.0 and 128.4 (3 d, 5 C in Ph), 139.2 (s, C in Ph), 155.2 (s, C-2) and 164.9, 171.6 and 172.0 (3 s, 3 CO); m/z 401 (M^+ , 61%), 370 (18), 342 (15), 257 (100) and 91 (79).

Dimethyl (Z,1RS,5RS,6SR,7SR)-8-benzyl-2-ethoxycarbonylmethylene-8-azabicyclo[3.2.1]octane-6,7-dicarboxylate [(Z)-18b]. An oil contaminated with ca. 30% of (Z)-18c; δ_{H} (C_6D_6 ; 500 MHz) 0.93 (1 H, t, J 7.0, CH_3), 1.19–1.25 (1 H, m, 4-H), 1.72–1.82 (2 H, m, 3- and 4-H), 2.48–2.59 (1 H, m, 3-H), 3.29 and 3.31 (6 H, each s, 2 OCH_3), 3.58 (1 H, d, J 6.7, 6-H), 3.59 and 3.70 (2 H, each d, J 13.7, CH_2Ph), 3.63 (1 H, br s, 5-H), 3.91 and 3.95 (2 H, each dq, J 11.0 and 7.0, OCH_2), 4.28 (1 H, t, J 7, 7-H), 5.70 (1 H, d, J 2.4, $\text{C}=\text{CH}$), 5.87 (1 H, d, J 7.0, 1-H), 7.04–7.09 (1 H, m, Ph), 7.12–7.16 (2 H, m, Ph) and 7.29–7.33 (2 H, m, Ph); δ_{C} (C_6D_6 ; 125 MHz) 14.2 (q, CH_3), 28.1 and 29.0 (2 t, C-3 and -4), 49.1 and 50.6 (2 d, C-6 and -7), 51.6 and 51.7 (2 q, 2 OMe), 53.6 (t, CH_2Ph), 59.6 (t, OCH_2), 61.4 (d, C-5), 62.3 (d, C-1), 117.6 (d, $\text{C}=\text{CH}$), 127.3, 128.5 and 128.8 (3 d, 5 C in Ph), 139.0 (s, C in Ph), 154.8 (s, C-2) and 165.3, 172.0 and 174.3 (3 s, 3CO).

Dimethyl (Z,1RS,5RS,6RS,7RS)-8-benzyl-2-ethoxycarbonylmethylene-8-azabicyclo[3.2.1]octane-6,7-dicarboxylate [(Z)-18c]. An oil contaminated with ca. 70% of (Z)-18b; δ_{H} (C_6D_6 ; 500 MHz) δ 0.93 (3 H, t, J 7.0, CH_3), 1.25–1.31 (1 H, m, 4-H), 1.72–1.82

(2 H, m, 3- and 4-H), 2.48–2.59 (1 H, m, 3-H), 3.24 and 3.33 (6 H, each s, 2 OCH₃), 3.46–3.50 (1 H, m, 5-H), 3.67 and 3.77 (2 H, each d, *J* 13.4, CH₂Ph), 3.65–3.67 (1 H, br d, *J* 7, 7-H), 3.84–4.00 (2 H, m, OCH₂), 4.18 (1 H, t, *J* 6.7, 6-H), 5.64 (1 H, d, *J* 2.1, C=CH), 5.89 (1 H, s, 1-H), 7.04–7.09 (1 H, m, Ph), 7.12–7.16 (2 H, m, Ph), and 7.29–7.33 (2 H, m, Ph); δ_{C} (C₆D₆; 125 MHz) 14.2 (q, CH₃), 25.7 and 28.2 (2 t, C-3 and -4), 49.8 and 50.0 (2 d, C-6 and -7), 51.5 and 51.9 (2 q, 2 OMe), 53.4 (t, CH₂Ph), 59.7 (t, OCH₂), 60.1 (d, C-5), 62.4 (d, C-1), 115.3 (d, C=CH), 128.3, 128.5 and 128.7 (3 d, 5 C in Ph), 139.0 (s, C in Ph), 156.5 (s, C-2) and 165.4, 172.4 and 174.0 (3 s, 3 CO).

Ethyl (Z,1*RS*,5*SR*)-(8-benzyl-7-methoxycarbonyl-8-azabicyclo[3.2.1]oct-6-en-2-ylidene)acetate [(Z)-19]. An oil (Found: *M*⁺ 341.1631. C₂₀H₂₃NO₄ requires *M*, 341.1627); ν_{max} (film)/cm⁻¹ 1715 (C=O); δ_{H} (300 MHz) 1.20 (1 H, t, *J* 7.2, CH₃), 1.49 (1 H, ddd, *J* 13.2, 7.5 and 2.2, 4-H), 2.03 (1 H, dddd, *J* 13.2, 10.6, 6.8 and 3.9, 4-H), 2.13–2.21 (1 H, br dd, *J* 16 and 7, 3-H), 2.47 (1 H, dddd, *J* 16.1, 10.6, 8 and 2.5, 3-H), 3.63 and 3.71 (2 H, each d, *J* 13.4, CH₂Ph), 3.71–3.76 (1 H, m, 5-H), 3.73 (3 H, s, OCH₃), 4.05 and 4.13 (2 H, each dq, *J* 10.8 and 7.2, OCH₂), 5.64 (1 H, br d, *J* 2, C=CH), 5.75 (1 H, s, H-1), 7.05 (1 H, d, *J* 2.4, 6-H) and 7.19–7.34 (5 H, m, Ph); δ_{C} (C₆D₆; 100 MHz) 14.3 (q, CH₃), 24.0 and 27.2 (2 t, C-3 and -4), 51.7 (q, OMe), 55.5 (t, CH₂Ph), 59.8 (t, OCH₂), 63.6 (d, C-5), 64.2 (d, C-1), 115.9 (d, C=CH), 126.8, 128.1 and 128.6 (3 d, 5 C in Ph), 137.0 and 138.2 (2 s, C-7 and C in Ph), 143.8 (d, C-6), 152.3 (s, C-2) and 163.9 and 165.4 (2 s, 2 CO); *m/z* 341 (*M*⁺, 15%), 282 (32), 268 (14), 250 (64), 236 (32), 204 (24) and 91 (100).

Dimethyl (E,1*RS*,5*SR*)-(8-benzyl-2-ethoxycarbonylmethylene-8-azabicyclo[3.2.1]oct-6-ene-6,7-dicarboxylate [(E)-20]. An oil (Found: *M*⁺ 399.1683. C₂₂H₂₅NO₆ requires *M*, 399.1682); ν_{max} (film)/cm⁻¹ 1720 (C=O); δ_{H} (300 MHz) 1.27 (3 H, t, *J* 7.2, CH₃), 1.72–1.81 (1 H, br dd, *J* 13 and 8, 4-H), 1.89–2.01 (1 H, m, 4-H), 2.30–2.45 and 3.59–3.69 (2 H, each m, 3-H₂), 3.66 (2 H, s, CH₂Ph), 3.77 and 3.83 (6 H, each s, 2 OCH₃), 3.92–3.95 (1 H, br t, *J* 3, 5-H), 4.06 (1 H, s, 1-H), 4.14 (2 H, q, *J* 7.2, OCH₂), 5.62 (1 H, d, *J* 2.4, C=CH) and 7.23–7.35 (5 H, m, Ph); δ_{C} (100 MHz) 14.4 (q, CH₃), 21.2 and 23.0 (2 t, C-3 and -4), 52.3 and 52.4 (2 q, 2 OMe), 56.2 (t, CH₂Ph), 59.9 (t, OCH₂), 67.5 (d, C-5), 74.2 (d, C-1), 116.1 (d, C=CH), 127.2, 128.3 and 128.6 (3 d, 5 C in Ph), 137.4, 137.9 and 142.4 (3 s, C-6, C-7 and C in Ph), 151.9 (s, C-2) and 163.3, 164.7 and 165.9 (3 s, 3 CO); *m/z* 399 (*M*⁺, 10%), 367 (9), 340 (65), 308 (55), 294 (21), 276 (25), 262 (10) and 91 (100).

Dimethyl (Z,1*RS*,5*SR*)-(8-benzyl-2-ethoxycarbonylmethylene-8-azabicyclo[3.2.1]oct-6-ene-6,7-dicarboxylate [(Z)-20]. An oil (Found: *M*⁺ 399.1683. C₂₂H₂₅NO₆ requires *M*, 399.1682); ν_{max} (CHCl₃)/cm⁻¹ 1705 (C=O); δ_{H} (300 MHz) 1.19 (3 H, t, *J* 7.1, CH₃), 1.72–1.81 (1 H, br dd, *J* 13 and 9 Hz, 4-H), 1.95–2.07 (1 H, m, 4-H), 2.29 (1 H, dd, *J* 16 and 6 Hz, 3-H), 2.55 (1 H, dddd, *J* 16.3, 10.4, 8.1 and 2.2, 3-H), 3.64 and 3.73 (2 H, each d, *J* 13.3, CH₂Ph), 3.78 and 3.80 (6 H, each s, 2 OCH₃), 3.90–3.93 (1 H, br t, *J* 3, 5-H), 4.03 and 4.10 (2 H, each dq, *J* 10.8 and 7.1, OCH₂), 5.65 (1 H, d, *J* 2.2, C=CH), 5.82 (1 H, s, 1-H) and 7.21–7.34 (5 H, m, Ph); δ_{C} (100 MHz) 14.3 (q, CH₃), 25.1 and 27.2 (2 t, C-3 and -4), 52.28 and 52.31 (2 q, 2 OMe), 56.4 (t, CH₂Ph), 59.9 (t, OCH₂), 66.2 (d, C-5), 67.8 (d, C-1), 116.0 (d, C=CH), 127.0, 128.2 and 128.6 (3 d, 5 C in Ph), 137.8 (s, C in Ph), 140.1 and 140.4 (2 s, C-6 and -7), 150.9 (s, C-2) and 164.0, 164.3 and 165.3 (3 s, 3 CO); *m/z*

399 (*M*⁺, 10%), 367 (13), 340 (48), 308 (60), 294 (42), 276 (33), 262 (17) and 91 (100).

Dimethyl (Z,1*RS*,6*SR*)-9-benzyl-2-ethoxycarbonylmethylene-9-azabicyclo[4.2.1]non-7-ene-7,8-dicarboxylate [(Z)-21]. An oil (Found: *M*⁺ 413.1836. C₂₃H₂₇NO₆ requires *M*, 413.1838); ν_{max} (CHCl₃)/cm⁻¹ 1720 (C=O); δ_{H} (300 MHz) 1.22 (3 H, t, *J* 7.2, CH₃), 1.37–1.54 (1 H, m, 4-H), 1.69–1.85 (1 H, m, 4- and 5-H), 1.88–1.98 (1 H, m, 5-H), 2.30–2.39 (1 H, br dd, *J* 14 and 7, 3-H), 2.96–3.07 (1 H, br t, *J* 14, 3-H), 3.71 and 3.77 (6 H, each s, 2 OCH₃), 3.75 and 3.83 (2 H, each d, *J* 13.4, CH₂Ph), 4.02 and 4.10 (2 H, each dq, *J* 10.8 and 7.2, OCH₂), 4.06–4.12 (1 H, m, 6-H), 5.71–5.74 (1 H, m, C=CH), 5.76 (1 H, br s, 1-H) and 7.21–7.38 (5 H, m, Ph); δ_{C} (100 MHz) 14.4 (q, CH₃), 23.0 (t, C-4), 34.2 and 38.7 (2 t, C-3 and -5), 52.1 and 52.3 (2 q, 2 OMe), 59.9 (t, OCH₂), 61.0 (t, CH₂Ph), 70.7 (d, C-6), 73.6 (d, C-1), 115.3 (d, C=CH), 126.9, 128.2 and 128.3 (3 d, 5 C in Ph), 132.5, 139.0 and 145.9 (3 s, C-7, C-8 and C in Ph) and 160.4, 163.0, 165.8 and 166.0 (4 s, C-2 and 3 CO); *m/z* 413 (*M*⁺, 5%), 381 (47), 354 (23), 322 (22), 308 (53), 280 (53) and 91 (100).

Ethyl (Z,1*RS*,5*RS*,7*RS*)-(8-benzyl-7-methoxycarbonyl-5-methyl-8-azatricyclo[3.2.1]oct-2-ylidene)acetate [(Z)-22a]. An oil (Found: *M*⁺ 357.1940. C₂₁H₂₇NO₄ requires *M*, 357.1940); ν_{max} (film)/cm⁻¹ 1710 (C=O); δ_{H} (300 MHz) 0.97 (3 H, t, *J* 7.2, CH₃), 1.21 (3 H, s, 5-CH₃), 1.51 (1 H, dd, *J* 13.4 and 7, 4-H), 1.86–1.99 (1 H, m, 4-H), 2.14 (1 H, dd, *J* 13.0 and 9.7, 6-H), 2.15–2.24 (1 H, br dd, *J* 15 and 6, 3-H), 2.34 (1 H, ddd, *J* 13.0, 5.0 and 1.7, 6-H), 2.42–2.55 (1 H, m, 3-H), 2.78 (1 H, dd, *J* 9.7 and 5.0, 7-H), 3.42 and 3.76 (2 H, each d, *J* 14.2, CH₂Ph), 3.68 (3 H, s, OCH₃), 3.72 and 3.90 (2 H, each dq, *J* 10.8 and 7.2, OCH₂), 5.13 (1 H, s, 1-H), 5.73 (1 H, d, *J* 2.2, C=CH) and 7.21–7.29 (5 H, m, Ph); δ_{C} (100 MHz) 13.9 (q, CH₂CH₃), 24.7 (q, 5-CH₃), 29.7, 33.9 and 39.9 (3 t, C-3, -4 and -6), 44.7 (d, C-7), 47.3 (t, CH₂Ph), 52.1 (q, OMe), 59.9 (s, C-5), 60.0 (t, OCH₂), 60.6 (d, C-1), 116.9 (d, C=CH), 126.4, 127.8 and 127.9 (3 d, 5 C in Ph), 139.8 (s, C in Ph), 154.3 (s, C-2) and 165.5 and 174.6 (2 s, 2 CO); *m/z* 357 (*M*⁺, 53%), 298 (60), 284 (38), 271 (27), 252 (56), 234 (49), 220 (30), 188 (21) and 91 (100).

Ethyl (Z,1*RS*,5*RS*,7*SR*)-(8-benzyl-7-methoxycarbonyl-5-methyl-8-azatricyclo[3.2.1]oct-2-ylidene)acetate [(Z)-22b]. An oil (Found: *M*⁺ 357.1944. C₂₁H₂₇NO₄ requires *M*, 357.1940); ν_{max} (CHCl₃)/cm⁻¹ 1720 (C=O); δ_{H} (300 MHz) 1.10 (3 H, t, *J* 7.2, CH₃), 1.22 (s, 3H, 5-CH₃), 1.49–1.58 (1 H, br dd, *J* 13 and 8, 4-H), 1.88–1.96 (1 H, br dd, *J* 13 and 7, 4-H), 2.00 (1 H, ddd, *J* 13.2, 12 and 1.5, 6-H), 2.11–2.21 (1 H, br dd, *J* 16 and 7, 3-H), 2.37 (1 H, dd, *J* 13.2 and 6.4, 6-H), 2.59–2.73 (1 H, m, 3-H), 3.30–3.39 (1 H, br dt, *J* 12 and 7, 7-H), 3.44 and 3.70 (2 H, each d, *J* 13.6, CH₂Ph), 3.55 (3 H, s, OCH₃), 3.89 and 3.99 (2 H, each dq, *J* 10.8 and 7.2, OCH₂), 5.10 (1 H, d, *J* 7.0, 1-H), 5.78 (1 H, d, *J* 2.4, C=CH) and 7.21–7.27 (5 H, m, Ph); δ_{C} (100 MHz) 14.3 (q, CH₂CH₃), 25.3 (q, 5-CH₃), 29.4 (t, C-3), 33.8 (t, C-4), 38.1 (t, C-6), 45.4 (d, C-7), 48.0 (t, CH₂Ph), 51.6 (q, OMe), 59.7 (t, OCH₂), 59.9 (s, C-5), 60.0 (d, C-1), 118.5 (d, C=CH), 126.5, 127.9 and 128.3 (3 d, 5 C in Ph), 139.8 (s, C in Ph), 154.5 (s, C-2) and 165.2 and 173.0 (2 s, 2 CO); *m/z* 357 (*M*⁺, 77%), 298 (39), 284 (66), 271 (61), 252 (33), 234 (46), 220 (27), 188 (25) and 91 (100).

Ethyl (Z,1*RS*,5*RS*,6*SR*)-(8-benzyl-6-methoxycarbonyl-5-methyl-8-azatricyclo[3.2.1]oct-2-ylidene)acetate [(Z)-23]. An oil

(Found: M^+ 357.1947. $C_{21}H_{27}NO_4$ requires M , 357.1940); $\nu_{\max}$ ($CHCl_3$)/ cm^{-1} 1710 (C=O); δ_H (300 MHz) 1.04 (3 H, t, J 7.2, CH_3), 1.12 (3 H, s, 5- CH_3), 1.60 (1 H, dd, J 13.2 and 6.8, 4-H), 1.83 (1 H, dd, J 13.2 and 9.4, 7-H), 1.91–2.02 (1 H, br td, J 13 and 6 Hz, 4-H), 2.15–2.23 (1 H, br dd, J 16 and 6, 3-H), 2.45–2.52 (1 H, m, 3-H), 2.51–2.61 (1 H, br ddd, J 13, 7 and 6, 7-H), 2.97 (1 H, dd, J 9.4 and 5.8, 6-H), 3.50 and 3.75 (2 H, each d, J 14.1, CH_2Ph), 3.73 (3 H, s, OCH_3), 3.82 and 3.92 (2 H, each dq, J 10.8 and 7.2, OCH_2), 4.91 (1 H, d, J 7.0, 1-H), 5.71 (1 H, d, J 2.0, C=CH) and 7.15–7.32 (5 H, m, Ph); δ_C (100 MHz) 14.1 (q, CH_2CH_3), 21.7 (q, 5- CH_3), 29.3 (t, C-3), 31.5 (t, C-7), 36.1 (t, C-4), 47.7 (t, CH_2Ph), 51.4 (d, C-6), 51.6 (q, OMe), 57.1 (d, C-1), 59.9 (t, OCH_2), 63.3 (s, C-5), 116.1 (d, C=CH), 126.4 (d, C in Ph), 127.9 (d, 4 C in Ph), 140.1 (s, C in Ph), 156.5 (s, C-2) and 165.6 and 175.2 (2 s, 2 CO); m/z 357 (M^+ , 39%), 312 (21), 298 (22), 284 (41), 271 (54), 266 (40), 220 (100), 188 (28) and 91 (89).

Dimethyl (Z,1RS,5RS)-8-benzyl-2-ethoxycarbonylmethylene-5-methyl-8-azabicyclo[3.2.1]oct-6-ene-6,7-dicarboxylate [(Z)-24]. An oil (Found: M^+ 413.1843. $C_{23}H_{27}NO_6$ requires M , 413.1838); $\nu_{\max}$ (film)/ cm^{-1} 1720 (C=O); δ_H (300 MHz) 1.05 (3 H, t, J 7.2, CH_3), 1.32 (3 H, s, 5- CH_3), 1.77–1.86 (1 H, br dd, J 13 and 7, 4-H), 1.95 (1 H, ddd, J 13.4, 10.5 and 6.6, 4-H), 2.22–2.30 (1 H, br dd, J 16 and 6, 3-H), 2.53–2.66 (1 H, m, 3-H), 3.59 and 3.66 (2 H, each d, J 13.6, CH_2Ph), 3.70 and 3.86 (6 H, each s, 2 OCH_3), 3.81 and 3.97 (2 H, each dq, J 10.8 and 7.2, OCH_2), 5.55 (1 H, d, J 7.0, 1-H), 5.74 (1 H, d, J 2.4, C=CH) and 7.18–7.31 (5 H, m, Ph); δ_C (100 MHz) 14.1 (q, CH_2CH_3), 20.0 (q, 5- CH_3), 28.3 (t, C-3), 29.5 (t, C-4), 49.0 (t, CH_2Ph), 52.0 and 52.3 (2 q, 2 OMe), 59.8 (t, OCH_2), 62.9 (d, C-1), 69.1 (s, C-5), 118.5 (d, C=CH), 126.7, 128.0 and 128.4 (3 d, 5 C in Ph), 136.4 and 138.2 (2 s, C-7 and C in Ph), 147.5 and 150.0 (2 s, C-2 and -6) and 162.7, 164.9 and 165.4 (3 s, 3 CO); m/z 413 (M^+ , 24%), 354 (82), 322 (39), 308 (36), 262 (22) and 91 (100).

Methyl (1RS,5RS,7RS)-8-benzyl-2-dicyanomethylene-8-azatricyclo[3.2.1]octane-7-carboxylate (25a). An oil (Found: M^+ 321.1473. $C_{19}H_{19}N_3O_2$ requires M , 321.1477); $\nu_{\max}$ ($CHCl_3$)/ cm^{-1} 2240 (C≡N) and 1730 (C=O); δ_H (300 MHz) 1.68–1.77 (1 H, br dd, J 13 and 8, 4-H), 2.05 (1 H, dd, J 13.6 and 9.7, 6-H), 2.12–2.25 (1 H, m, 4-H), 2.49 (1 H, ddd, J 16.7, 11.4 and 7.9, 3-H), 2.65–2.74 (1 H, m, 6-H), 2.89 (1 H, dd, J 9.7 and 5.3, 7-H), 2.90–3.00 (1 H, br ddd, J 17, 8 and 1, 3-H), 3.44–3.49 (1 H, m, 5-H), 3.72 (2 H, s, CH_2Ph), 3.76 (3 H, s, OCH_3), 4.50 (1 H, s, 1-H) and 7.24–7.37 (5 H, m, Ph); δ_C (100 MHz) 26.6 (t, C-3), 29.8 (t, C-4), 31.4 (t, C-6), 47.0 (d, C-7), 52.7 (q, OMe), 53.1 (t, CH_2Ph), 56.7 (d, C-5), 65.4 (d, C-1), 82.9 [s, $C(CN)_2$], 111.0 and 111.1 (2 s, 2 CN), 127.5, 128.1 and 128.4 (3 d, 5 C in Ph), 137.0 (s, C in Ph), 172.8 (s, CO) and 180.0 (s, C-2); m/z 321 (M^+ , 52%), 290 (13), 262 (15), 235 (27) and 91 (100).

Methyl (1RS,5RS,7SR)-8-benzyl-2-dicyanomethylene-8-azatricyclo[3.2.1]octane-7-carboxylate (25b). An oil (Found: M^+ 321.1479. $C_{19}H_{19}N_3O_2$ requires M , 321.1477); $\nu_{\max}$ ($CHCl_3$)/ cm^{-1} 2230 (C≡N) and 1730 (C=O); δ_H (300 MHz) 1.71–1.80 (1 H, br dd, J 13 and 8, 4-H), 2.14–2.26 (2 H, m, 4- and 6-H), 2.65–2.74 (1 H, br td, J 13 and 7, 6-H), 2.54–2.67 (1 H, br ddd, J 17, 11 and 8, 3-H), 2.86–2.96 (1 H, br dd, J 17 and 7, 3-H), 3.38–3.44 (1 H, m, 5-H), 3.55–3.64 (1 H, br dt, J 12 and 6, 7-H), 3.70 (3 H, s, OCH_3), 3.71 (2 H, s, CH_2Ph), 4.36 (1 H, d, J 6.4, 1-H) and 7.23–7.38

(5 H, m, Ph); δ_C (100 MHz) 26.3 (t, C-3), 29.5 and 29.8 (2 t, C-4 and -6), 47.1 (d, C-7), 52.4 (q, OMe), 53.4 (t, CH_2Ph), 56.4 (d, C-5), 65.7 (d, C-1), 84.5 [s, $C(CN)_2$], 110.5 and 111.1 (2 s, 2 CN), 127.6, 128.3 and 128.5 (3 d, 5 C in Ph), 136.9 (s, C in Ph), 171.5 (s, CO) and 179.6 (s, C-2); m/z 321 (M^+ , 40%), 290 (9), 235 (28) and 91 (100).

Dimethyl (1RS,5SR)-8-benzyl-2-dicyanomethylene-8-azatricyclo[3.2.1]oct-6-ene-6,7-dicarboxylate (26). Pale yellow crystals; mp 131–133 °C (hexane–ethyl acetate) (Found: C, 66.96; H, 5.24; N, 10.95. $C_{21}H_{19}N_3O_4$ requires C, 66.83; H, 5.07; N, 11.13%); $\nu_{\max}$ ($CHCl_3$)/ cm^{-1} 2240 (C≡N) and 1730 (C=O); δ_H (300 MHz) 1.88–1.95 and 2.14–2.26 (2 H, 2 m, 4- H_2), 2.56–2.66 (1 H, br quintet, J 9, 3-H), 2.94–3.01 (1 H, m, 3-H), 3.72 (2 H, br s, CH_2Ph), 3.83 and 3.87 (6 H, 2 s, 2 OCH_3), 4.03 (1 H, br s, 5-H), 4.78 (1 H, s, 1-H) and 7.26–7.38 (5 H, m, Ph); δ_C (100 MHz) 23.4 and 25.4 (2 t, C-3 and -4), 52.8, (2 q, 2 OMe), 56.4 (t, CH_2Ph), 66.9 and 70.0 (2 d, C-1 and -5), 84.4 [s, $C(CN)_2$], 110.2 and 110.8 (2 s, 2 CN), 127.8, 128.5 and 128.6 (3 d, 5 C in Ph), 135.1, 136.0 and 144.6 (3 s, C-6, C-7 and C in Ph), 161.9 and 163.8 (2 s, 2 CO) and 172.9 (s, C-2); m/z 377 (M^+ , 18%), 318 (22), 286 (26), 254 (13), 242 (34) and 91 (100).

Acknowledgements

This work was partially supported by a Grant-in-Aid for Scientific Research (C) (No. 13672239) administered by the Japan Society for the Promotion of Science. The authors wish to thank the staff of the Analysis Center of Meiji Pharmaceutical University for performing the elemental analysis and measurements of LA-NMR (NOESY spectra) (Ms S. Kubota) and mass spectra (Ms T. Koseki). We are also grateful to Mr T. Ujike and Mr T. Okuda for their technical assistance.

References and notes

- For reviews, see: W. J. Lown, in *1,3-Dipolar Cycloaddition Chemistry*, ed. A. Padwa, Wiley, New York, 1984, vol. 1; W. H. Pearson and P. Stoy, *Synlett*, 2003, 903–921; I. Coldham and R. Hufton, *Chem. Rev.*, 2005, **105**, 2765–2809; G. Pandey, P. Banerjee and S. R. Gadre, *Chem. Rev.*, 2006, **106**, 4484–4517; H. Pellissier, *Tetrahedron*, 2007, **63**, 3235–3285.
- (a) R. Huisgen, W. Scheer and H. Huber, *J. Am. Chem. Soc.*, 1967, **89**, 1753–1755; (b) P. DeShong, D. A. Kell and D. R. Sidler, *J. Org. Chem.*, 1985, **50**, 2309–2315; (c) W. Eberbach, I. Heinze, K. Knoll, H. Fritz and F. Borle, *Helv. Chim. Acta*, 1988, **71**, 404–418; (d) B. R. Henke, A. J. Kouklis and C. H. Heathcock, *J. Org. Chem.*, 1992, **57**, 7056–7066; (e) P. Garner, P. B. Cox, J. T. Anderson, J. Protasiewicz and R. Zaniwski, *J. Org. Chem.*, 1997, **62**, 493–498; (f) C. Gaebert and J. Mattay, *Tetrahedron*, 1997, **53**, 14297–14316.
- C. Chen, X. Li and S. L. Schreiber, *J. Am. Chem. Soc.*, 2003, **125**, 10174–10175; W. Gao, X. Zhang and M. Raghunath, *Org. Lett.*, 2005, **7**, 4241–4244.
- (a) K. Ishii, Y. Shimada, S. Sugiyama and M. Noji, *J. Chem. Soc., Perkin Trans. 1*, 2000, 3022–3024; (b) K. Ishii, T. Sone, Y. Shimada, T. Shigeyama, M. Noji and S. Sugiyama, *Tetrahedron*, 2004, **60**, 10887–10898; (c) K. Ishii, T. Sone, T. Shigeyama, M. Noji and S. Sugiyama, *Tetrahedron*, 2006, **62**, 10865–10878.
- (a) M. Kotera, K. Ishii, O. Tamura and M. Sakamoto, *J. Chem. Soc., Perkin Trans. 1*, 1994, 2353–2354; (b) M. Kotera, K. Ishii, O. Tamura and M. Sakamoto, *J. Chem. Soc., Perkin Trans. 1*, 1998, 313–318; (c) M. Kotera, K. Ishii, M. Hiraga and M. Sakamoto, *Heterocycles*, 1999, **51**, 2147–2157.
- For recent examples, see: V. K. Aggarwal, R. S. Grainger, G. K. Newton, P. L. Spargo, A. D. Hobson and H. Adams, *Org. Biomol. Chem.*, 2003, **1**, 1884–1893; C. G. Thomson, E. Carlson, G. G. Chicchi, J. J. Kulagowski, M. M. Kurtz, C. J. Swain, K.-L. C. Tsao and A. Wheeldon, *Bioorg.*

- Med. Chem. Lett.*, 2006, **16**, 811–814; I. T. Huscroft, E. J. Carlson, G. G. Chicchi, M. M. Kurtz, C. London, P. Raubo, A. Wheeldon and J. J. Kulagowski, *Bioorg. Med. Chem. Lett.*, 2006, **16**, 2008–2012; N. R. Curtis, R. G. Ball and J. J. Kulagowski, *Tetrahedron Lett.*, 2006, **47**, 2635–2638; K. M. Peese and D. Y. Gin, *J. Am. Chem. Soc.*, 2006, **128**, 8734–8735.
- 7 J. W. Lown and K. Matsumoto, *J. Chem. Soc. D*, 1970, 692–693.
- 8 M. T. Barros, C. D. Maycock and M. R. Ventura, *Tetrahedron Lett.*, 2002, **43**, 4329–4331.
- 9 The deshielding effect of the carboxyl group was also observed in the bicyclic epoxide system (see ref. 5c).
- 10 V. Bardakos and C. Sandris, *Org. Magn. Reson.*, 1981, **15**, 339–342.
- 11 The characteristic cross-peaks between 3-H and 7-H in compounds (*E*)-**15a**, (*Z*)-**15a** and **25a** and between 6-H and 9-H in (*Z*)-**17a** indicate an *exo*-configuration. In contrast, in (*E*)-**15b** the cross-peak between 1-H and 7-H indicates an *endo*-configuration. The cross-peaks between 3-H and 7-H and 6-H and 7-H in compound (*Z*)-**18a** indicate a 6,7-*cis* and *exo*-configuration. The cross-peaks between 1-H and 7-H and 3-H and 7-H in (*Z*)-**18b** indicate a 6,7-*trans* and *endo*-configuration. The cross-peaks between 3-H and 7-H and 5-H and 6-H in (*Z*)-**18c** indicate a 6,7-*trans* and *exo*-configuration. The cross-peak between 3-H and 6-H in compound (*Z*)-**23** characteristically indicates an *exo*-configuration.
- 12 The cycloaddition of the 6-membered carbonyl ylide derived from the epoxide **3** and ethyl vinyl ether afforded the adduct **4** in 44% yield (see ref. 5b).
- 13 Yields for compounds are based on converted starting material.